ERDC/EL TR-00-15

Environmental Laboratory

**US Army Corps
of Engineers®**
Engineer Research and
Development Center

An Analysis of Freshwater Mussels (Unionidae) in the Upper Ohio River near Huntington, West Virginia, 1998 Studies

Andrew C. Miller and Barry S. Payne

October 2000

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

PRINTED ON RECYCLED PAPER

An Analysis of Freshwater Mussels (Unionidae) in the Upper Ohio River near Huntington, West Virginia, 1998 Studies

by Andrew C. Miller, Barry S. Payne
Environmental Laboratory
U.S. Army Engineer Research and Development Center
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Final report

Approved for public release; distribution is unlimited

Engineer Research and Development Center Cataloging-in-Publication Data

Miller, Andrew C.

An analysis of freshwater mussels (Unionidae) in the upper Ohio River near Huntington, West Virginia, 1998 studies / by Andrew C. Miller, Barry S. Payne ; prepared for U.S. Army Engineer District, Huntington.

45 p. : ill. ; 28 cm. -- (ERDC/EL ; TR-00-15)

Includes bibliographic references.

1. Unionidae -- Ohio River. 2. Freshwater mussels -- Ohio River. 3. Freshwater mussels -- Field studies. I. Payne, Barry S. II. United States. Army. Corps of Engineers. Huntington District. III. Engineer Research and Development Center (U.S.) IV. Environmental Laboratory (U.S.) V. Title. VI. Series: ERDC/EL TR ; 00-15.
TA7 E8 no.ERDC/EL TR-00-15

Contents

Preface	iv
1—Introduction	1
Background	1
Purpose and Scope	2
2—Study Area and Methods	3
Study Area	3
Methods	3
Preliminary reconnaissance	3
Qualitative mussel samples	3
Quantitative mussel samples	4
Data analysis	4
3—The Bivalve Community	5
Bivalve Community Characteristics	5
Evidence of Recent Recruitment	6
Mean Density of Unionidae	6
Relationships Between Unionidae and <i>C. fluminea</i>	7
Relationship Between Unionidae and <i>Dreissena polymorpha</i>	7
Changes in Species Richness and Species Diversity	8
Size Demography of Dominant Populations	8
4—Discussion	10
References	13
Tables 1-4	
Figures 1-18	
Appendix A: Freshwater Mussels Collected from Upper Ohio River Using Quantitative Methods	A1

SF 298

Preface

The work reported herein was conducted by the Environmental Laboratory (EL), Vicksburg, MS, U.S. Army Engineer Research and Development Center (ERDC), in July 1993 for the U.S. Army Engineer District, Huntington, Huntington, WV. The purpose was to use qualitative and quantitative methods to collect freshwater mussels (Unionidae) from known mussel beds in a reach of the Ohio River near Huntington. Data on density, size demography of dominant populations, species diversity, and community composition will be used to analyze the environmental effects of water resource developments along the upper Ohio River.

Divers were Messrs. Larry Neill and Robert T. James, Muscle Shoals, AL, and Dennis Baxter and Johnny Buchanan, Norris, TN, from the Tennessee Valley Authority (TVA). Assistance in the field and advice on location of sampling sites was provided by Mr. Bill Tolin, U.S. Fish and Wildlife Service, and Mr. Barry Passmore, Huntington District. Mr. John Campbell and Ms. Mindy Hamilton, Marshall University, Huntington, WV, assisted in the field.

During the conduct of this study Dr. John Harrison was Chief, EL; Dr. C. J. Kirby was Chief, Ecological Research Division, EL; and Dr. E. A. Theriot was Chief of the Aquatic Ecology Branch, Ecological Research Division. Safety equipment and foul weather gear were provided by TVA. Authors of this report were Drs. Andrew C. Miller and Barry S. Payne, Aquatic Ecology Branch. Design of the study and conduct of all related activities in the field and laboratory were the sole responsibility of Drs. Miller and Payne.

At the time of publication of this report, Dr. James R. Houston was Director of ERDC, and COL James S. Weller, EN, was Commander.

Miller, A. C., and Payne, B. S. (2000). "An analysis of freshwater mussels (Unionidae) in the upper Ohio River near Huntington, West Virginia; 1998 studies," ERDC/EL TR-00-15, U.S. Army Engineer Research and Development Center, Vicksburg, MS.

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

1 Introduction

Background

Freshwater mussels are long-lived (30 or more years in some species), rely on particulate organic matter for nutrition, and because they are relatively nonmotile, are unable to change their surroundings if conditions become unsuitable. Their habitat is affected by local and upriver changes in climate, season, land use, edaphic conditions, and water level. Habitat conditions are also affected by source and nonsource pollution, commodity movement, and commercial developments in the watershed. It is clear that man-made and natural disturbances in large rivers are long term, complex, and often subtle and episodic. The characteristics of freshwater mussels can make them susceptible to these disturbances. Thus these organisms and their habitats are particularly appropriate for long-term ecological studies.

The biological consequences of man-made and natural disturbances to habitat (specific physical and chemical conditions) can be measured on organisms held in the laboratory. However, caution must be used when extrapolating results of laboratory experiments to the field (Payne, Miller, and Aldridge 1987). Physiological responses that occur in a laboratory often do not take place under natural conditions. Planners and biologists should evaluate the effects of natural and man-made disturbances on naturally occurring populations rather than on organisms held in the laboratory. Field studies are the best means of understanding effects of physical disturbances on natural populations. Studies can be designed to evaluate physical effects of water resource development on recruitment, rate of growth, density, species richness, and diversity. These parameters provide the most useful measures of the overall health and ultimate survival of a mussel community. Long-term studies on freshwater mussels in large rivers should not be discounted as "mere monitoring" (Taylor 1987). Instead, they provide an opportunity to investigate the effects of complex, episodic events on a resource with ecological, economic, and cultural value.

Purpose and Scope

The purpose of this study was to characterize important biotic variables (mussel density, evidence of recent recruitment, community structure, and spatial distribution) of mussel beds downriver of the Robert Byrd Lock and Dam in the Ohio River, West Virginia. Results will be used to evaluate effects of water resource developments on freshwater mussels. In addition, data will be used to assess effects of the exotic zebra mussel (*Dreissena polymorpha*) on native freshwater mussels.

2 Study Area and Methods

Study Area

The Ohio River originates in Pittsburgh, PA, at the confluence of the Allegheny and Monongahela Rivers. It flows 1,578 km (981 miles) to the northwest and then the southwest before it joins the Mississippi River near Cairo, IL. The river drains 528,101 sq km (203,900 square miles) and falls 137 m (450 ft) before it joins the Mississippi River.

For this project, study sites were located in the upper Ohio River (UOR) between River Miles (RM) 292 and 284 (Table 1, Figures 1-3). Quantitative and qualitative methods were used to collect bivalves at RM 287, 292, 289, and 284 (Table 1, Figures 2, 3). The majority of samples were collected about 61 m (200 ft) from shore in water between 3.7 and 7.6 m (12 and 25 ft) deep. Global Positioning System (GPS) coordinates for each mussel bed are listed in Table 1.

Methods

Preliminary reconnaissance

All underwater work was accomplished by a dive crew equipped with surface-supplied air and communication equipment. Before intensive sampling was initiated, a single diver conducted a preliminary reconnaissance of the immediate area. He obtained information on substratum type, water velocity, and presence of mussels to determine if he was on the mussel bed and would be able to collect specimens using quantitative methods. Since this river reach was studied in 1992 and 1993, some knowledge was available on the approximate location of mussel beds (Miller and Payne 1993, 1995).

Qualitative mussel samples

The majority of the qualitative samples were obtained by having two divers collect simultaneously. Each diver placed a specific number of live mussels in each of six nylon bags; 5 mussels were placed in three bags and 20 were placed

in each of the other nine bags. At each site 12 samples and approximately 200 individuals were collected. Divers were instructed to obtain mussels without bias toward size or type. They attempted to exclude the Asiatic clam, *Corbicula fluminea*. If *C. fluminea* was inadvertently collected, it was later eliminated. A total of 84 qualitative samples (bags of mussels) were taken at four mussel beds (Table 1).

All mussels were brought to the surface, counted, and identified. Data were recorded on standard data sheets and returned to the laboratory for analysis and plotting. Shells of voucher specimens for each species were placed in plastic zip-lock bags and labeled with high-rag-content paper. Mussels not needed for voucher were returned to the river. Methods for sampling mussels are based on techniques described in Miller and Nelson (1983); Isom and Gooch (1986); Kovalak, Dennis, and Bates (1986); Miller and Payne (1988); and Miller et al. (1993). Mussel identification was based on taxonomic keys and descriptive information in Burch (1975), Cummings and Mayer (1992), and Parmalee and Bogan (1998). Taxonomy is consistent with Williams et al. (1993).

Quantitative mussel samples

Quantitative samples (that included unionids as well as *C. fluminea*) were obtained at RM 287, 292, 290, and 284. At each mussel bed, bivalves were collected at each of two to five sites that were separated by 5-10 m. At each site, 10 quadrats (0.25 sq m) were positioned approximately 1 m apart. A diver excavated all sand, gravel, shells, and live bivalves to a depth of 100-150 mm. Material was sent to the surface in a 20-L bucket and transported to shore. Sediment was screened through a sieve series. All live mussels and *C. fluminea* removed from samples were placed in 4-L zip-lock bags. Each bivalve was then identified and total shell length (SL) measured to the nearest 0.1 mm with digital calipers. Bivalves were processed in the field and returned to the river unharmed.

Data analysis

Species diversity H' was determined with the following formula:

$$H' = - (p_j \log p_j) \quad (1)$$

where p_j is the proportion of the population that is of the j^{th} species (Shannon and Weaver 1949). Evenness was calculated with the modified Hill's ratio (Ludwig and Reynolds 1988). All calculations were done with programs written in BASIC or SAS (Statistical Analytical System) on a personal computer. Discussion of statistical procedures that were used can be found in Green (1979) and Hurlbert (1984). Species area curves and dominance-diversity curves were constructed from qualitative and quantitative biological data. More information on methods used for this survey can be found in McNaughton and Wolf (1973), Isom and Gooch (1986), Kovalak, Dennis, and Bates (1986), Miller and Payne (1988), and Miller et al. (1993).

3 The Bivalve Community

Bivalve Community Characteristics

Over 1,200 freshwater mussels representing 22 species were collected in the study area using qualitative methods (Table 2). Three more species were collected using qualitative than quantitative methods (a total of 19). Qualitative sampling typically generates a longer species list simply because more organisms are collected. However, the quantitative collecting has the advantage of obtaining the smaller species and individuals that divers can miss searching by hand when visibility is low.

A total of 30 species of freshwater mussels, not counting the nonindigenous zebra mussel (*D. polymorpha*) or Asian clam (*C. fluminea*), have been collected in this river reach since the beginning of this project (Table 2). Differences in the total species count each year are caused by slight variations in the sampling locations or simply chance. For species that represent less than 0.01 percent of the community, it is obvious that several thousand mussels might have to be obtained before one individual of that species is collected.

Based upon qualitative collecting at three sites, the fauna was dominated by two thick-shelled species (*Elliptio crassidens* and *Quadrula pustulosa*), which composed 44.18 and 13.76 percent of the assemblage, respectively. *Elliptio crassidens* was taken in more than 90 percent of the samples and *Q. p. pustulosa*, which was much less abundant, was taken in slightly more than 80 percent of the samples. Eleven species represented from 1 to 10 percent of the fauna, and ten species made up less than 1 percent of the fauna. With the exception of two species, the majority of the mussels were either fairly common or rare. Virtually all of the mussels have massively thick shells such as those belonging to the genera *Quadrula*, *Elliptio*, *Fusconaia*, *Amblema*, or *Megaloniaia*s. Species that did not belong to this group included *Leptodea fragilis*, which has a very thin shell, and *Potamilus alatus*, which has a slightly thicker shell. Thin-shelled species were uncommon in the study area.

A total of 188 individuals and 19 species were collected in 120 quantitative samples (Tables 2 and 3 and Appendix A). *Quadrula p. pustulosa*, *E. crassidens*, and *Obovaria reflexa* composed 26.6, 18.1, and 15.4 percent of the Unionidae (Table 3). Fourteen species were between 1 and 7 percent of the

fauna and two species (*Lasmigonia complanata* and *Lampsilis teres*) composed less than 1 percent of the fauna. As stated previously, fewer species (19) were taken using quantitative rather than qualitative methods (22 species, Table 2). This was the result of the large number of individuals collected using qualitative methods (1,279) compared with quantitative methods (188).

Evidence of Recent Recruitment

An indicator of recent recruitment is the percentage of individuals and species less than 30 mm total shell length. This will include all Unionidae less than 2-3 years old regardless of species. Based upon data obtained in 1992 and 1993, recent recruitment was low in this river reach (Figures 4 and 5). In 1992 and 1993, less than 10 percent of all native mussels collected were less than 30 mm total shell length. In 1998 the percentage of individuals less than 30 mm ranged from 14 to 41 percent at three locations, a notable increase. In 1992 less than 20 percent of the species showed evidence of recent recruitment. In 1993 recent recruitment was evident only at RM 287; none was found at the other locations. In 1998 more than 20 percent of the mussel species collected at each of three locations (RM 284, 287, and 290) showed evidence of recent recruitment.

The only species showing evidence of recent recruitment in 1992 were *Truncilla truncata* and *Q. p. pustulosa*. The only species besides *Q. p. pustulosa* showing evidence of recent recruitment in 1993 were *T. truncata* and *O. reflexa*. In 1998 a total of nine species showed evidence of recent recruitment: *A. p. plicata*, *E. lineolata*, *L. fragilis*, *O. reflexa*, *P. alatus*, *Q. p. pustulosa*, *Q. quadrula*, *T. truncata*, and *T. donaciformis*.

Mean Density of Unionidae

Mean densities were low in 1998 in this reach of the river; maximum density was less than 8 individuals/sq m at RM 284 and 287 (Figure 6). Mean unionid densities have always been low in this river reach (Figure 7). Mean density ranged from 7.9 to 13.7 in 1992, from 3.6 to 13.4 in 1993, and from 0.8 to 11.2 in 1998. Although densities were variable among years and sites during any one year, there appears to be no specific trend with respect to native mussels. Mean densities were either similar to or greater than previous years in the beds at RM 284, 287, and 290. The extreme reduction in density at RM 292 in 1998 is difficult to explain; however, it could be due to loss of an older cohort or simply spatial variation in mussel abundance.

Relationships Between Unionidae and *C. fluminea*

Mean density of *C. fluminea* was moderately high, 80 to 100 individuals/sq m at RM 287 and 290 (Figure 8). High-density populations of *C. fluminea* were found in 1992 and 1993 (Figure 9). In 1992 mean densities at RM 284 exceeded 3,000 individuals/sq m. In 1998 mean density of all *C. fluminea* populations at all locations was less than 500 individuals/sq m. At two locations (RM 284 and 290) mean densities were less than 100 individuals/sq m. Density of this species has declined dramatically in this reach of the UOR since these surveys began.

Various authors have reported that high densities of *C. fluminea* could negatively affect native freshwater mussels by competing for food and space. After many years of detailed study in the lower Ohio River, Miller and Payne (1994) were unable to demonstrate any positive or negative relationship between nonindigenous and native bivalves. It appears that densities of *Corbicula* are gradually decreasing, and obviously there will be only limited interactions between these species in the future.

Relationship Between Unionidae and *Dreissena polymorpha*

The zebra mussel, *D. polymorpha*, has now become part of the bivalve fauna throughout the Mississippi River from Wisconsin to New Orleans and virtually all of the Ohio River. Only eight zebra mussels were collected at these sites during 1993. All were located at RM 292.0 and were obtained in quantitative samples; estimated density was 1.6/sq m. *Dreissena polymorpha*, unlike *C. fluminea*, can byssally attach to unionids in high numbers. Therefore, there is potentially much more direct competition between zebra mussels and native mussels than between zebra mussels and Asian clams.

Based upon results of quantitative sampling in 1998, mean density of zebra mussels in the substratum was estimated at less than 100 individuals/sq m. Zebra mussels were attached to cobble and gravel, dead shells, and live mussels. The number of attached zebra mussels on each native mussel was counted to determine infestation rate. Fifteen percent of the native mussels harbored no attached zebra mussels, and 41.5 percent had from one to nine attached zebra mussels. Only 18.5 percent were considered to be heavily infested, and had more than 50 attached zebra mussels, as shown in the following tabulation.

Infestation Level	Percentage of Unionids
Not infested	41.5
Less than 10 zebra mussels	25.0
Between 10 and 50 zebra mussels	18.5
More than 50 zebra mussels	15.0

In the early 1990s extremely high densities of zebra mussels were found at a dense and diverse mussel bed in the lower Ohio River near Olmsted, IL (Miller and Payne 1997). In August 1994, mean densities at some locations on the mussel bed were as high as 300,000 individuals/sq m. In July 1995, this cohort ranged from 11.3 to nearly 27.3 mm total shell length. Mean density at most sites was 20,000 to 50,000 individuals/sq mm. Between July and October of 1995 the majority of the mussels in that cohort died. Mean density of the zebra mussels at that site was less than 100 individuals/sq m, and shell length ranged from 1.4 to 20.2 mm. Quantitative samples have been taken at that bed every year since 1993; although zebra mussels are still at that site, their numbers are typically less than 1,000/sq m.

Changes in Species Richness and Species Diversity

Overall richness and diversity of species collected in this reach of the Ohio River (qualitative and quantitative samples combined) have gradually increased during the study period (Figure 10). In 1992, 21 species were collected; in 1993, 25 species were found; and in 1998, a total of 26 species were collected. So far, 30 species have been found at this location. Shannon's diversity index (calculated for all the mussels collected using qualitative methods) was higher in 1998 than in either of the previous 2 years. This slight increase was probably the result of a more even distribution of species in these latter collections.

Size Demography of Dominant Populations

Unionid density was low at RM 284 and 287 and extremely low at RM 290 and 292 (Figure 6). Mean density was approximately 7.2 and 6.2 individuals/sq m at sites with moderate density, and less than 1.0 individual/sq m at sites with low density. Despite such low densities, unionids showed substantial evidence of recent recruitment, evident in the population size structure of the most abundant species. For example, 13 of 50 *Q. p. pustulosa* were less than 30 mm long; the smallest was only 14 mm long (Figure 11). The estimated age of *Q. p. pustulosa* ranging from 15 to 30 mm is 2 to 4 years. The largest *Q. p. pustulosa* obtained was 65 mm long—a very large adult of this species in the UOR.

In contrast, *E. crassidens* showed no evidence of recent recruitment (Figure 12). All individuals of this species were greater than 90 mm long. The largest individual obtained was 170 mm long—a very large adult of this species. The moderate sample size ($n = 34$) did not allow a conclusion that absolutely no recent recruitment of this species had occurred. However, if moderate recent recruitment had occurred, a few small mussels would have been obtained. Thus, at best, recent recruitment of this species has been low, and no recruitment has occurred recently.

Recent recruitment was clearly evident in the *A. p. plicata* population, although no large adults were present (Figure 13). Only 13 individuals of this species were obtained. Nearly all of these were recent recruits 12 to 30 mm long that were likely 1 to 3 years old. The largest individual was only 47 mm long; individuals of this species commonly often attain maximum lengths of more than 90 mm.

The *O. reflexa* population was similar to that of *Q. p. pustulosa* in that there was evidence both of recent recruitment and occurrence of large adults (Figure 14). Six of fifteen individuals obtained of this species were less than 30 mm long and represented recent recruitment. Four individuals were 40 to 50 mm long. Although 50 mm is not an absolute maximum length in the Ohio River, this length is certainly a large adult size for this species.

Corbicula fluminea density was greatest at RM 287 and 290, averaging 108 and 85 individuals/sq m, respectively (Figure 8). Density at RM 284 and 292 averaged less than 40 individuals/sq m. All of these density values are low for this species, which can often be more than 1,000 individual/sq m in suitable habitats.

Size structure of *C. fluminea* population varied among sites. At RM 284, individuals ranged from 5 to 20 mm, and two cohorts were evident (Figure 15). One was centered at 10 mm and the second at approximately 15 mm. At RM 287, individuals ranged from 7 to 27 mm, but only one cohort was evident between 7 and 20 mm (Figure 16). At RM 290 (Figure 17), individuals ranged from 5 to 21 mm, and two cohorts were present. The major cohort was centered at 10 mm, and a minor cohort was centered at approximately 14 mm. At RM 292 individuals ranged from 4 to 18 mm, and two cohorts were present (Figure 18). The major intersite differences probably reflect combined effects of variable cohort strength and individual growth rates among sites.

4 Discussion

The fauna in this reach of the UOR consisted almost entirely of thick-shelled species such as *E. crassidens*, *Q. p. pustulosa*, with lesser numbers of *Q. metanevra* and *Pleurobema cordatum*. Thin-shelled species such as *pyganodon* spp. and *L. fragilis* were either absent or very uncommon. Each year slightly more species have been collected (Figure 10, Table 4). There is simply no indication that species are being lost from this reach of the river, or that community composition is changing greatly. Conditions actually appear to be improving.

Total species richness in the study area (30 species based on quantitative and qualitative sampling since 1992 (Table 2) is similar to that at other mussel beds in large rivers. At a mussel bed in the lower Ohio River near Olmsted, 26 species of freshwater mussels were collected using qualitative and quantitative methods (Miller, Payne, and Siemsen 1986). In a survey of the lower Tennessee River, Miller, Payne, and Tippit (1992) collected 4,768 individuals and identified 23 species.

In 1993 a single specimen of *Lampsilis abrupta* (Say, 1831), listed as endangered by the U.S. Fish and Wildlife Service (1991), was collected during qualitative searches by divers at RM 287.2 (Table 2). This species was not collected in 1992 or 1998. Although this species has been collected at other locations in the UOR (Tolin, Schmidt, and Zeto 1987), *L. abrupta* is obviously very uncommon. Density of this species must be less than 0.0168/sq m (less than 1 individual/ 60 sq m). It should be noted that even at extremely low densities (1/100 sq m) there could be many tens of *L. abrupta* in this river reach.

Another uncommon species of some concern is *Plethobasus cyphus*, listed as endangered by the Commonwealth of Kentucky (Branson et al. 1981). This species was collected in all 3 years. It has been found in a dense and diverse bed in the lower Ohio River near Olmsted (Payne and Miller 1994), and composed 0.19 percent of the fauna at a site stabilized by wing dams in Pool 10 of the upper Mississippi River (Miller 1988).

These mussel beds in this reach of the Ohio River can be characterized as having extremely low density (Figure 7). Mean density each year was less than 14 individuals/sq m. Spatial and temporal density differences have been notable.

At RM 290, density in 1998 was almost double that in 1993. However, at RM 292 mean density was substantially less than in 1992 and 1993. Mean density at RM 284 and 287 was approximately the same each study year. These density data are less than values often found at other large river mussel beds. In the lower Tennessee River, Miller, Payne, and Tippit (1992) estimated total density to range from 9.2 to 128.0 individuals/sq m at six closely placed sites. In a survey of the upper Mississippi River, Miller et al. (1990) reported that total mussel density ranged from 5.2 to 333.2 individuals/sq m at 16 sites. At half of those sites, total density was greater than 50 individuals/sq m and at four sites it was greater than 100 individuals/sq m. At an inshore and offshore site sampled in 1986 at RM 18.6 in the lower Tennessee River, total mussel density was 187.7 and 79.7 individuals/sq m, respectively (Way, Miller, and Payne 1989).

More evidence of recent recruitment was noted in this reach of the Ohio River than during previous years (Figures 4 and 5). Good evidence of recruitment was noted in 1998 for *Q. p. pustulosa* (Figure 11), *A. p. plicata* (Figure 13), and *O. reflexa* (Figure 14). However, the size demography of *E. crassidens* was similar in 1992, 1993, and 1998; during all years this population was composed of mainly old, large individuals. *Elliptio crassidens* requires the skipjack herring, *Alosa chrysochloris*, as the host fish for its glochidia (Fuller 1974). High-lift dams prevent annual upstream migrations by this fish species (Fuller 1974). Ten such dams, constructed during this century, occur between these study sites and the confluence of the Ohio River with the Mississippi River. It is likely that numbers of this species in the Ohio River have declined as a result of reduction of skipjack herring.

Demography of *Q. p. pustulosa* was virtually identical all study years, and showed good evidence of recent recruitment. This species, which uses some relatively common fish hosts (the black and brown bullhead, channel and flathead catfish, and white crappie (Fuller 1974)), has consistently shown substantial recent recruitment. This illustrates that for some species, conditions for successful recruitment are present.

Mean density of *C. fluminea* declined in this reach of the UOR during the study period (Figure 9). Convincing evidence was not found that this species competes with native mussels (Miller and Payne 1994). It was also noted that numbers of this species have declined in the lower Ohio River since the early 1980s (Payne and Miller 1994).

Mackie (1991) described impacts of *D. polymorpha* on Unionidae in Lake St. Clair during 1991. He noted that zebra mussels can affect the following activities of native species: normal burrowing and locomotion; balance and equilibrium; ability to close the shell; function of the siphons and therefore respiration and feeding; growth, reproduction, and feeding; food supplies; shell growth; and occlusion of the shell gape and ultimately smothering or death. Ricciardi, Whoriskey, and Rasmussen (1995) examined patterns of unionid infestations and predicted that severe unionid mortality, greater than 90 percent, occurs when *D. polymorpha* density is greater than 6,000 individuals/m² and

more than 100 zebra mussels per unionid. These conditions do not currently exist in the UOR.

Although zebra mussels were collected in this reach of the UOR, their numbers are still too low to negatively affect the native species. Zebra mussels were first collected in the Illinois River in 1992 (Thiel, Blodgett, and Miller 1993). By 1993 density had increased dramatically; at RM 5.5 at Grafton, IL, values as high as 61,126/m² were reported (Whitney, Blodgett, and Sparks 1995). However, by fall 1993, it was estimated that mortality at RM 5.5 and 66.8 was 22 percent and 41 percent, respectively. By late 1994 the investigators estimated that there had been a 99 percent reduction in zebra mussels at both RM 5.5 and 66.8. In 1995, investigators collected 34 1-m quadrats in the lower 193 km (120 miles) and found only 109 adult zebra mussels greater than 20 mm long (Whitney, Blodgett, and Sparks 1996).

The continued use of inland waterways to transport bulk commodities (Dietz et al. 1983) has caused planners and biologists in government agencies to express concern over the possible negative effects of commercial use of waterways on freshwater mussels. Quantitative and qualitative techniques are being used to obtain data on mussel density, relative species abundance, community composition, population demography, and presence of nonindigenous species. These parameters provide the most useful measures of the overall health and ultimate survival of a mussel community. They provide an opportunity to investigate the effects of complex, episodic events on a resource with ecological, economic, and cultural value.

Based upon information gathered in 1998, the condition at the mussel beds is good and is certainly no worse than in previous years. With respect to recent recruitment and continued existence of uncommon native species of freshwater mussels, it appears to be improving slightly.

References

- Branson, B. A., Harker, D. F., Baskin, J. M., Medley, M. E., Batch, D. L., Warren, M. L., Davis, W. H., Houtcooper, W. C., Monore, B., Phillippe, L. R., and Cupp, P. (1981). "Endangered, threatened, and rare animals and plants of Kentucky," *Transactions of the Kentucky Academy of Sciences* 42, 77-89.
- Burch, J. B. (1975). *Freshwater unionacean clams (Mollusca: Pelecypoda) of North America*. Malacological Publications, Hamburg, MI.
- Cummings, K., and Mayer, C. A. (1992). *Field guide to freshwater mussels of the Midwest*. Illinois Natural History Survey, Manual 5, Illinois Natural History Survey, Champaign, IL.
- Dietz, A. R., Harrison, R. W., Olson, H. E., Grier, D., and C. Simpkins. (1983). "National waterways study--A framework for decision making--Final Report," Report NWS-83-1, U.S. Army Engineer Institute for Water Resources, Water Resources Support Center, Fort Belvoir, VA.
- Fuller, S. L. H. (1974). "Clams and mussels (Mollusca:Bivalvia)." *Pollution ecology of fresh water invertebrates*. C. Hart and S. L. H. Fuller, ed., Academic Press, New York, 215-273.
- Green, R. H. (1979). *Sampling design and statistical methods for environmental biologists*. John Wiley & Sons, New York.
- Hurlbert, S. H. (1984). "Pseudoreplication and the design of ecological field experiments," *Ecological Monographs*, 54, 187-211.
- Isom, B. G., and Gooch, C. (1986). "Rationale and sampling design for fresh-water mussels, unionidae, in streams, large rivers, impoundments, and lakes." *Rationale for sampling and interpretation of ecological data in the assessment of freshwater ecosystems*. ASTM STP 894, B. G. Isom, ed., American Society for Testing and Materials, Philadelphia, PA, 46-59.

- Kovalak, W. P., Dennis, S. D., and Bates, J. M. (1986). "Sampling effort required to find rare species of freshwater mussels." *Rationale for sampling and interpretation of ecological data in the assessment of freshwater ecosystems*. ASTM STP 894, B. G. Isom, ed., American Society for Testing and Materials, Philadelphia, PA, 34-45.
- Ludwig, J. A., and Reynolds, J. F. (1988). *Statistical ecology; A primer on methods and computing*. John Wiley & Sons, New York.
- Mackie, G. L. (1991). "Biology of the exotic zebra mussel, *Dreissena polymorpha* in relation to native bivalves and its potential to impact in Lake St. Clair," *Hydrobiologia* 219, 251-268.
- McNaughton, S. J., and Wolf, L. L. (1973). *General ecology*. Holt, Rinehart, and Winston, New York.
- Miller, A. C. (1988). "Mussel fauna associated with wing dams in Pool 7 of the upper Mississippi River," *Freshwater Ecology* 4, 299-302.
- Miller, A. C., and Nelson, D. A. (1983). "An instruction report on freshwater mussels," Instruction Report EI-83-2, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Miller, A. C., and Payne, B. S. (1988). "The need for quantitative sampling to characterize size demography and density of freshwater mussel communities," *American Malacological Bulletin* 6, 49-54.
- Miller, A. C., and Payne, B. S. (1993). "An analysis of freshwater mussels (Unionidae) in the upper Ohio River near Huntington, West Virginia; 1992 studies," Technical Report EL-93-27, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Miller, A. C., and Payne, B. S. (1994). "Co-occurrence of native freshwater mussels (Unionidae) and the nonindigenous *Corbicula fluminea* at two stable shoals in the Ohio River," *Malacological Review* 27, 87-97.
- Miller, A. C., and Payne, B. S. (1995). "An analysis of freshwater mussels (Unionidae) in the upper Ohio River near Huntington, West Virginia; 1993 studies," Technical Report EL-95-2, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Miller, A. C., Payne, B. S., and Siemsen, T. (1986). "Description of the habitat of the endangered mussel, *Plethobasus cooperianus*," *The Nautilus* 100, 14-18.
- Miller, A. C., Payne, B. S., and Tippit, R. (1992). "Characterization of a freshwater mussel community immediately downriver of Kentucky Lock and Dam in the Tennessee River," *Transactions of the Kentucky Academy of Sciences* 53, 154-161.

- Miller, A. C., Payne, B. S., Hornbach, D. J., and Ragland, D. V. (1990). "Physical effects of increased commercial navigation traffic in the Upper Mississippi River; Phase I studies," Technical Report EL-90-3, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Miller, A. C., Payne, B. S., Shafer, D. S., and Neill, L. T. (1993). "Techniques for monitoring bivalve communities and populations in large rivers." *Conservation and management of freshwater mussels*, Proceedings of an Upper Mississippi River Conservation Committee symposium, St. Louis, MO, 12-14 October 1992. K. S. Cummings, A. C. Buchanan, and L. M. Koch, ed., Rock Island, IL, 147-158.
- Parmalee, P. W., and Bogan, A. E. (1998). *The freshwater mussels of Tennessee*. The University of Tennessee Press, Knoxville, TN.
- Payne, B. S., and Miller, A. C. (1994). "Freshwater mussels at two beds in the lower Ohio River near Olmsted, Illinois; 1992 studies," Technical Report EL-94-2, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Payne, B. S., and Miller, Andrew C. (1997). "Spatial distribution of mussels at a bed in the lower Ohio River near Olmsted, Illinois," Technical Report EL-97-3, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Payne, B. S., Miller, A. C., and Aldridge, D. W. (1987). "Environmental effects of navigation traffic: Laboratory studies of the effects on mussels of intermittent exposure to turbulence and suspended solids," Technical Report EL-87-14, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Ricciardi, A., Whoriskey, F. G., and Rasmussen, J. B. (1995). "Predicting the intensity and impact of *Dreissena* infestation on native unionid bivalves from *Dreissena* field density," *Canadian Journal of Fisheries and Aquatic Sciences* 52, 1449-1461.
- Shannon, C. E., and Weaver, W. (1949). *The mathematical theory of communication*. University of Illinois Press, Urbana IL.
- Taylor, L. R. (1987). "Objective and experiment in long-term research." *Long-term studies in ecology*. G. E. Likens, ed., Springer-Verlag, New York, 20-70.
- Thiel, P. A., Blodgett, K. D., and Miller, A. C. (1993). "Zebra mussel invasion of the Illinois and Mississippi Rivers," *Dreissena polymorpha Information Review* 4(1), 1.

- Tolin, W. A., Schmidt, J., and Zeto, M. (1987). "A new location for the Federally-listed endangered unionid *Lampsilis abrupta* (Say 1821) (= *Lampsilis orbiculata* (Hildreth 1828)), the pink mucket, in the upper Ohio River bordering West Virginia," *Malacology Data Net* 2, 18.
- U.S. Fish and Wildlife Service. (1991). "Endangered and threatened wildlife and plants," *Federal Register*, July 15, 1991. 50 CFR 17.11 & 17.12.
- Way, C. M., Miller, A. C., and Payne, B. S. (1989). "The influence of physical factors on the distribution and abundance of freshwater mussels (Bivalvia: Unionidae) in the lower Tennessee River," *The Nautilus* 103, 96-98.
- Whitney, S. D., Blodgett, K. D., and Sparks, R. E. (1995). "Update on zebra mussels and native unionids in the Illinois River," *Illinois Natural History Survey, Zebra Mussel Research Team*, January, 1.
- Whitney, S. D., Blodgett, K. D., and Sparks, R. E. (1996). "Update on zebra mussels and native unionids in the Illinois River; Part III," *Illinois Natural History Survey, Zebra Mussel Research Team*, April, 1.
- Williams, J. D., Warren, M. L., Jr., Cummins, K. S., Harris, J. L., and R. J. Neves. (1993). "Conservation status of freshwater mussels of the United States and Canada," *Fisheries* 18(9), 6-22.

Figure 1. The study area in the upper Ohio River (to convert miles to kilometers, multiply by 1.609)

Figure 2. Sample sites located between RM 287 and 284 in the upper Ohio River, 1998 (to convert inches to millimeters, multiply by 25.4; to convert feet to meters, multiply by 0.3048)

Figure 3. Sample sites located between RM 292 and 288 in the upper Ohio River, 1998

Figure 4. Evidence of recent mussel recruitment, percent individuals less than 30 mm total shell length

Figure 5. Evidence of recent mussel recruitment, percent species less than 30 mm total shell length

Figure 6. Mean unionid density at four mussel beds in the upper Ohio River, 1998

Figure 7. Mean unionid density, 1992, 1993, and 1998, at four locations in the upper Ohio River

Figure 8. Mean density of the Asian clam *C. fluminea* at four locations in the upper Ohio River, 1998

Figure 9. Mean *C. fluminea* density, 1992, 1993, and 1998, at four locations in the upper Ohio River

Figure 10. Changes in species richness and species diversity, all sites combined, qualitative samples only, upper Ohio River for years 1992, 1993, and 1998

Figure 11. Size demography for *Q. p. pustulosa*, all locations combined, in the upper Ohio River, 1998

Figure 12. Size demography for *E. crassidens*, all locations combined, in the upper Ohio River, 1998

Figure 13. Size demography for *A. p. plicata*, all locations combined, in the upper Ohio River, 1998

Figure 14. Size demography for *O. reflexa*, all locations combined, in the upper Ohio River, 1998

Figure 15. Size demography for *C. fluminea* at upper Ohio River Mile 284, 1998

Figure 16. Size demography for *C. fluminea* at upper Ohio River Mile 287, 1998

Figure 17. Size demography for *C. fluminea* at upper Ohio River Mile 290, 1998

Figure 18. Size demography for *C. fluminea* at upper Ohio River Mile 292, 1998

Location and Number of Samples Collected Using Qualitative and Quantitative Methods in the Upper Ohio River, September 1998, 61 m (200 ft) from Shore, Depth of Water 3.7-7.6 m (12-25 ft)

Date	Approx. River Mile	Global Positioning System	Bank ¹	Qualitative		Quantiative	
				No. of Sites	Total	No. of Subsites	No. of Samples
18 Sep	287	38 ° 35.302 82 ° 12.677	RDB	2	24	4	40
19 Sep	292	38 ° 25.210 82 ° 12.811	RDB			2	20
19 Sep	290	38 ° 35.913 82 ° 15.592	RDB	1	12	1	10
20 Sep	284	38 ° 36.703 82 ° 10.249	LDB			2	20
20 Sep	284		LDB	2	24	1	10
21 Sep	284		LDB	2	24	2	20
Total Samples					84		120

¹ RDB = Right downbound; LDB = Left downbound.

Table 2
Species of Freshwater Mussels Collected in the upper Ohio River, 1992, 1993, and 1998

Scientific Name	Common Name	1992		1993		1999	
		Qual	Quant	Qual	Quant	Qual	Quant
<i>Actinonaias ligamentina</i>	Mucket	X	X	X	X	X	X
<i>Amblema plicata plicata</i>	Threeeridge	X	X	X	X	X	X
<i>Ellipsaria lineolata</i>	Butterfly	X	X	X	X	X	X
<i>Elliptio crassidens</i>	Elephant ear	X	X	X	X	X	X
<i>Elliptio dilatata</i>	Spike					X	
<i>Fusconaia ebena</i>	Ebony shell	X	X	X	X	X	X
<i>Fusconaia flava</i>	Wabash pigtoe			X	X		
<i>Fusconaia subrotunda</i>	Long-solid	X		X		X	
<i>Lampsilis abrupta</i>	Pink mucket			X			
<i>Lampsilis cardium</i>	Pocketbook	X	X	X	X	X	X
<i>Lampsilis siliquoidea</i>	Fatmucket					X	
<i>Lampsilis teres</i>	Yellow sandshell						X
<i>Lasmigonia complanata</i>	White heelsplitter			X			X
<i>Lasmigonia costata</i>	Fluted shell			X	X	X	
<i>Leptodea fragilis</i>	Fragile papershell	X		X		X	X
<i>Ligumia recta</i>	Black sandshell	X	X	X		X	X
<i>Megalonaias nervosa</i>	Washboard	X	X	X		X	X
<i>Obovaria reflexa</i>	Threehorn wartyback	X	X	X	X	X	X
<i>Obovaria retusa</i>	Ring pink		X				
<i>Obovaria subrotunda</i>	Round hickorynut			X		X	
<i>Plethobasus cyphus</i>	Sheepnose	X	X	X		X	
<i>Pleurobema coccineum</i>	Round pigtoe	X		X		X	
<i>Pleurobema cordatum</i>	Ohio pigtoe	X	X	X	X	X	X
<i>Potamilus alatus</i>	Pink heelsplitter	X	X	X	X	X	X
<i>Quadrula metanevra</i>	Monkeyface	X	X	X	X	X	X
<i>Quadrula nodulata</i>	Wartyback			X			
<i>Quadrula pustulosa pustulosa</i>	Pimpleback	X	X	X	X	X	X
<i>Quadrula quadrula</i>	Mapleleaf	X	X	X	X	X	X
<i>Truncilla truncata</i>	Deertoe	X	X		X		X
<i>Truncilla donaciformis</i>	Fawnsfoot	X					X
Total species		20	17	24	15	22	19
Total individuals		1477	328	4700	191	1279	188
Total sites		5	4	25	4	3	3
Total samples		97	120	293	118	84	120

Table 3**Percent Abundance and Frequency of Occurrence of Unionid Species Collected in the Upper Ohio River, September 1998 (All individuals were collected using quantitative methods)**

Species	Percent Abundance	Frequency of Occurrence
<i>Q. p. pustulosa</i>	26.60	29.17
<i>E. crassidens</i>	18.09	22.50
<i>O. reflexa</i>	15.43	22.50
<i>A. p. plicata</i>	6.91	9.17
<i>E. lineolata</i>	6.38	10.00
<i>A. ligamentina</i>	4.26	5.00
<i>Q. metanevra</i>	3.19	5.00
<i>P. alatus</i>	3.19	11.67
<i>L. recta</i>	2.66	4.17
<i>T. truncata</i>	2.66	3.33
<i>P. cordatum</i>	2.13	10.00
<i>Q. quadrula</i>	1.60	2.50
<i>L. fragilis</i>	1.60	2.50
<i>L. cardium</i>	1.06	1.67
<i>T. donaciformis</i>	1.06	2.50
<i>F. ebena</i>	1.06	1.67
<i>M. nervosa</i>	1.06	1.67
<i>L. complanata</i>	0.53	0.83
<i>L. teres</i>	0.53	0.83
Note: Total individuals = 188; total species = 19; total quadrats = 120.		

Table 4
Summary Information on Freshwater Mussels Collected in Qualitative Collections,
September 1998

Species	RM 287		RM 290		RM 284		Total	
	% Abun	% Occur	% Abun	% Occur	% Abun	% Occur	% Abun	% Occur
<i>Elliptio crassidens</i>	38.32	83.33	29.35	83.33	50.89	100.00	44.18	92.86
<i>Quadrula pustulosa</i> <i>pustulosa</i>	13.59	83.33	29.89	100.00	9.77	75.00	13.76	80.95
<i>Quadrula metanevra</i>	8.70	54.17	10.33	75.00	9.63	72.92	9.46	67.86
<i>Pleurobema cordatum</i>	11.14	75.00	6.52	58.33	3.58	41.67	6.18	53.57
<i>Actinonaias ligamentina</i>	2.72	33.33	1.09	8.33	8.12	70.83	5.55	51.19
<i>Potamilus alatus</i>	3.26	33.33	2.17	16.67	4.95	48.92	4.07	39.29
<i>Elliptio dilatata</i>	10.05	12.50	0.00	0.00	0.00	0.00	2.89	3.57
<i>Quadrula quadrula</i>	0.00	0.00	5.98	75.00	2.61	33.33	2.35	29.76
<i>Obovaria reflexa</i>	1.63	25.00	4.89	41.67	1.65	18.75	2.11	23.81
<i>Amblesma plicata plicata</i>	2.45	37.50	3.26	41.67	1.51	20.83	2.03	28.57
<i>Fusconaia ebena</i>	1.90	20.83	3.80	25.00	1.10	14.58	1.72	17.86
<i>Ligumia recta</i>	1.09	16.67	0.00	0.00	2.48	29.17	1.72	21.43
<i>Ellipsaria lineolata</i>	0.27	4.17	2.72	33.33	1.38	16.67	1.25	15.48
<i>Lampsilis cardium</i>	1.36	20.83	0.00	0.00	0.83	12.50	0.86	13.10
<i>Megaloniais nervosa</i>	1.63	20.83	0.00	0.00	0.55	8.33	0.78	10.71
<i>Fusconaia subrotunda</i>	0.82	8.33	0.00	0.00	0.28	4.17	0.39	4.76
<i>Obovaria subrotunda</i>	0.82	12.50	0.00	0.00	0.00	0.00	0.23	3.57
<i>Leptodea fragilis</i>	0.00	0.00	0.00	0.00	0.28	4.17	0.16	2.38
<i>Pleurobema coccineum</i>	0.00	0.00	0.00	0.00	0.14	2.08	0.08	1.19
<i>Lampsilis siliquioidea</i>	0.00	0.00	0.00	0.00	0.14	2.08	0.08	1.19
<i>Lasmigonia costata</i>	0.27	4.17	0.00	0.00	0.00	0.00	0.08	1.19
<i>Plethobasus cyphus</i>	0.00	0.00	0.00	0.00	0.14	2.08	0.08	1.19
Total individuals	368		184		727		1279	
Total species	17		11		19		22	
Total samples	24		12		48		84	

Appendix A

Freshwater Mussels Collected from Upper Ohio River Using Quantitative Methods

Table A1
Percent Species Abundance of Native Mussels Collected at
Ohio River Mile 284, September 1998

Species	Subsite Number				Total
	1	2	3	4	
<i>E. crassidens</i>	0.00	0.00	25.00	48.15	31.03
<i>Q. p. pustulosa</i>	0.00	0.00	10.00	22.22	13.79
<i>O. reflexa</i>	41.67	0.00	15.00	7.41	6.90
<i>E. lineolata</i>	8.33	20.00	10.00	3.70	6.90
<i>A. ligamentina</i>	0.00	0.00	10.00	7.41	6.90
<i>T. truncata</i>	8.33	0.00	5.00	0.00	6.90
<i>Q. metanevra</i>	0.00	0.00	0.00	3.70	10.34
<i>A. p. plicata</i>	16.67	0.00	5.00	0.00	0.00
<i>P. alatus</i>	8.33	20.00	0.00	0.00	3.45
<i>L. recta</i>	0.00	20.00	10.00	0.00	0.00
<i>T. donaciformis</i>	16.67	20.00	0.00	0.00	0.00
<i>L. fragilis</i>	0.00	0.00	5.00	3.70	0.00
<i>F. ebena</i>	0.00	0.00	0.00	0.00	6.90
<i>L. cardium</i>	0.00	20.00	0.00	0.00	3.45
<i>L. complanata</i>	0.00	0.00	5.00	0.00	0.00
<i>Q. quadrula</i>	0.00	0.00	0.00	0.00	3.45
<i>P. cordatum</i>	0.00	0.00	0.00	3.70	0.00
Total individuals	12	5	20	27	29
Total species	6	5	10	8	11
% Individuals < 30 mm	41.66	20	25	7.41	10.71
% Species < 30 mm	50	20	50	25	27.27
Menhinick's index	1.73	2.24	2.23	1.59	2.04
Species diversity H'	1.58	1.61	2.15	1.56	2.14
Evenness	1.16	0.00	1.34	0.72	0.95

Table A2 Percent Species Occurrence of Native Mussels Collected at Ohio River Mile 284, September 1998					
Species	Subsite Number				Total
	1	2	3	4	
<i>E. crassidens</i>	0.00	0.00	40.00	90.00	70.00
<i>O. reflexa</i>	50.00	0.00	30.00	10.00	20.00
<i>Q. p. pustulosa</i>	0.00	0.00	20.00	40.00	30.00
<i>E. lineolata</i>	10.00	10.00	20.00	10.00	20.00
<i>A. ligamentina</i>	0.00	0.00	20.00	10.00	10.00
<i>Q. metanevra</i>	0.00	0.00	0.00	10.00	30.00
<i>T. truncata</i>	10.00	0.00	10.00	0.00	20.00
<i>T. donaciformis</i>	20.00	10.00	0.00	0.00	0.00
<i>P. alatus</i>	10.00	10.00	0.00	0.00	10.00
<i>L. recta</i>	0.00	10.00	20.00	0.00	0.00
<i>L. fragilis</i>	0.00	0.00	10.00	10.00	0.00
<i>L. cardium</i>	0.00	10.00	0.00	0.00	10.00
<i>A. p. plicata</i>	10.00	0.00	10.00	0.00	0.00
<i>F. ebena</i>	0.00	0.00	0.00	0.00	20.00
<i>Q. quadrula</i>	0.00	0.00	0.00	0.00	10.00
<i>P. cordatum</i>	0.00	0.00	0.00	10.00	0.00
<i>L. complanata</i>	0.00	0.00	10.00	0.00	0.00
Total samples	10	10	10	10	10

Table A3
Percent Species Abundance of Native Mussels Collected at
Ohio River Mile 287, September 1998

Species	Subsite Number				Total
	1	2	3	4	
<i>Q. p. pustulosa</i>	33.33	35.29	30.00	52.94	38.10
<i>O. reflexa</i>	22.22	29.41	30.00	17.65	25.40
<i>E. crassidens</i>	0.00	17.65	5.00	11.76	9.52
<i>A. p. plicata</i>	22.22	0.00	5.00	5.88	6.35
<i>E. lineolata</i>	0.00	17.65	0.00	0.00	4.76
<i>M. nervosa</i>	0.00	0.00	5.00	5.88	3.17
<i>P. cordatum</i>	11.11	0.00	5.00	0.00	3.17
<i>L. recta</i>	11.11	0.00	0.00	5.88	3.17
<i>L. fragilis</i>	0.00	0.00	5.00	0.00	1.59
<i>P. alatus</i>	0.00	0.00	5.00	0.00	1.59
<i>Q. quadrula</i>	0.00	0.00	5.00	0.00	1.59
<i>L. teres</i>	0.00	0.00	5.00	0.00	1.59
Total individuals	9	17	20	17	63
Total species	5	4	10	6	12
% Individuals < 30 mm	33.30	47.06	40.00	41.17	41.27
% Species < 30 mm	60.00	75.00	50.00	50.00	50.00
Menhinick's index	1.67	0.97	2.23	1.45	1.51
Species diversity <i>H'</i>	1.52	1.34	1.92	1.39	1.85
Evenness	1.73	1.20	0.92	0.79	0.67

Table A4 Percent Species Abundance of Native Mussels Collected at Ohio River Mile 287, September 1998					
Species	Subsite Number				Total
	1	2	3	4	
<i>Q. p. pustulosa</i>	30.00	40.00	50.00	70.00	38.00
<i>O. reflexa</i>	20.00	40.00	50.00	30.00	28.00
<i>E. crassidens</i>	0.00	30.00	10.00	20.00	12.00
<i>A. p. plicata</i>	20.00	0.00	10.00	10.00	8.00
<i>E. lineolata</i>	0.00	30.00	0.00	0.00	6.00
<i>M. nervosa</i>	0.00	0.00	10.00	10.00	4.00
<i>P. cordatum</i>	10.00	0.00	10.00	0.00	4.00
<i>L. recta</i>	10.00	0.00	0.00	10.00	4.00
<i>L. fragilis</i>	0.00	0.00	10.00	0.00	2.00
<i>P. alatus</i>	0.00	0.00	10.00	0.00	2.00
<i>Q. quadrula</i>	0.00	0.00	10.00	0.00	2.00
<i>L. teres</i>	0.00	0.00	10.00	0.00	2.00
Total quadrats	10	10	10	10	40

Table A5 Percent Species Abundance of Native Mussels Collected at Ohio River Mile 290, September 1998		
Species	Percent Abundance	Percent Occurrence
<i>Q. p. pustulosa</i>	50.00	70.00
<i>A. p. plicata</i>	10.71	20.00
<i>A. ligamentina</i>	7.14	20.00
<i>Q. metanevra</i>	7.14	20.00
<i>E. lineolata</i>	7.14	20.00
<i>P. alatus</i>	7.14	20.00
<i>P. cordatum</i>	3.57	10.00
<i>E. crassidens</i>	3.57	10.00
<i>Q. quadrula</i>	3.57	10.00
Total individuals	28	
Total species	9	
Total quadrats	10	
Individuals < 30 mm	14.28	
Species < 30 mm	22.22	
Menhinick's index	1.51	
Species diversity <i>H'</i>	1.85	
Evenness	0.67	

Table A6 Percent Species Abundance of Native Mussels Collected at Ohio River Mile 292, September 1998					
Species	Subsite 1		Subsite 2		Total
	Abundance	Frequency	Abundance	Frequency	
<i>A. p. plicata</i>	0.00	0.00	100.00	30.00	75.00
<i>O. reflexa</i>	100.00	10.00	0.00	0.00	25.00
Total individuals	1		3		4
Total species	1		1		2
Total samples		10		10	20
% Individuals < 30 mm	0.00		66.67		50.00
% Species < 30 mm	0.00		100.00		50.00

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) October 2000		2. REPORT TYPE Final Report		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE An Analysis of Freshwater Mussels (Unionidae) in the Upper Ohio River near Huntington, West Virginia, 1998 Studies				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Andrew C. Miller, Barry S. Payne				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Engineer Research and Development Center Environmental Laboratory 3909 Halls Ferry Road Vicksburg, MS 39180-6199				8. PERFORMING ORGANIZATION REPORT NUMBER ERDC/EL TR-00-15	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Engineer District, Huntington 502 Eighth Street Huntington, WV 25701-2070				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <p>A survey to assess community characteristics, density, population demography of dominant species, and the likelihood of finding endangered species of freshwater mussels (Unionidae), was conducted in the upper Ohio River (approximate River Miles (RM) 292-284) near Huntington, WV, in September 1998. This was the third study year to assess impacts of commercial navigation traffic in the Ohio River. Similar studies were conducted in 1992 and 1993.</p> <p>In 1998 over 1,200 freshwater mussels representing 22 species were collected using qualitative methods. Mean densities of native unionids were low in 1998; maximum density was less than 8 individuals/sq m at RM 284 and 287. Although densities have been low and variable among years and sites, there are no specific temporal or spatial trends at this location. In 1998 the percentage of individuals less than 30 mm total shell length ranged from 14 to 41 percent, a substantial increase over the earlier two study years. The number of species identified in this reach of the Ohio River (qualitative and quantitative samples combined) has gradually increased during the study period. In 1992, 21 species were collected; in 1993, 25 species were found; and in 1998, a total of 26 species were collected. A total of 30 species have been found in this river reach during the three study years.</p> <p style="text-align: right;">(Continued)</p>					
15. SUBJECT TERMS Freshwater mussels Ohio River Navigation traffic impacts					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT UNCLASSIFIED	b. ABSTRACT	c. THIS PAGE UNCLASSIFIED			19b. TELEPHONE NUMBER (include area code)

14. (Concluded)

Although density of zebra mussels (*Dreissena polymorpha*) has increased in the study area, mean density was estimated at less than 100 individuals/sq m. Fifteen percent of the native mussels harbored no attached zebra mussels, and 41.5 percent had from one to nine attached zebra mussels. Only 18.5 percent were heavily infested. Density of *Corbicula fluminea* has declined considerably. Maximum density in 1992 was over 3,000 individuals/sq m; at the same site in 1998 the density was less than 100 individuals/sq m. Based on evidence of recent recruitment and continued existence of uncommon species, conditions at this bed have improved since the early 1990s.