

TECHNICAL REPORT HL-82-15

### THE ATCHAFALAYA RIVER DELTA

Report 9

WIND CLIMATOLOGY

by

Bruce A. Ebersole

Coastal Engineering Research Center

DEPARTMENT OF THE ARMY
Waterways Experiment Station, Corps of Engineers
PO Box 631
Vicksburg, Mississippi 39180-0631



January 1985 Report 9 of a Series

Approved For Public Release; Distribution Unlimited

Prepared for US Army Engineer District, New Orleans New Orleans, Louisiana 70160

Monitored by Hydraulics Laboratory
US Army Engineer Waterways Experiment Station
PO Box 631, Vicksburg, Mississippi 39180-0631

Destroy this report when no longer needed. Do not return it to the originator.

The findings in this report are not to be construed as an official Department of the Army position unless so designated.

by other authorized documents.

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Et	ntered)	
REPORT DOCUMENTATION P	READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER 2 Technical Report HL-82-15	. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE ATCHAFALAYA RIVER DELTA; ReporCLIMATOLOGY	Report 9 of a series  Report 9 of a series	
7. AUTHOR(0) Bruce A. Ebersole		8. CONTRACT OR GRANT NUMBER(8)
<ol> <li>PERFORMING ORGANIZATION NAME AND ADDRESS</li> <li>US Army Engineer Waterways Experim Coastal Engineering Research Cente PO Box 631, Vicksburg, Mississippi</li> </ol>	r	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Engineer District, New Orl PO Box 60267 New Orleans, Louisiana 70160 14. MONITORING AGENCY NAME & ADDRESS(II different	eans from Controlling Office)	12. REPORT DATE January 1985  13. NUMBER OF PAGES 122  15. SECURITY CLASS. (of this report)
US Army Engineer Waterways Experim Hydraulics Laboratory PO Box 631, Vicksburg, Mississippi	Unclassified  15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distr	ibution unlimit	ed.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

#### 18. SUPPLEMENTARY NOTES

Available from National Technical Information Service, 5285 Port Royal Road, Springfield, Virginia 22161.

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Atchfalaya Bay (La.) (LC)

Atmospheric circulation (LC)

Climatology--Louisiana (LC)

Winds--Louisiana--Atchafalaya Bay--Statistics (LC)

### 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

This report contains a description of the wind climate typical of the Atchafalaya Bay area. Frequencies of occurrence of various synoptic circulation patterns as well as statistical estimates of wind speed and direction for each pattern are given. The seasonal variation of the wind climate is investigated. Prototype wind data used in this study are comprised of 15 years of long-term data from New Orleans and Lake Charles and 1 year of data measured

(Continued)

### Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)
20. ABSTRACT (Continued).
within the Atchafalaya Bay. Some shipboard observations are included as well. Statistical information includes frequencies of occurrence, estimates of duration, and historical extremes.

#### PREFACE

This report presents an analysis of the wind climate within the Atchafalaya Bay, Louisiana. This study was supported as a reimbursable project within the Estuaries Division, Hydraulics Laboratory, US Army Engineer Waterways Experiment Station (WES), Vicksburg, Mississippi, for the US Army Engineer District, New Orleans, Louisiana.

The study was conducted under the direction of the following WES personnel: Messrs. H. B. Simmons, Chief of the Hydraulics Laboratory (Retired);

F. A. Herrmann, Jr., Chief of the Hydraulics Laboratory; R. A. Sager, Assistant Chief of the Hydraulics Laboratory; W. H. McAnally, Jr., Project Manager; and J. V. Letter, Jr., Task Coordinator. This study was performed by Mr. Bruce A. Ebersole, formerly of the Wave Dynamics Division (WDD), Hydraulics Laboratory, under the direct supervision of Dr. R. W. Whalin and Mr. C. E. Chatham, Jr., former and acting Chiefs of WDD. The WDD and its personnel were transferred to the Coastal Engineering Research Center (CERC), WES, on 1 July 1983, under the direction of Dr. R. W. Whalin, Chief of the Coastal Engineering Research Center.

Commanders and Directors of WES during the study and the preparation and publication of this report were COL John L. Cannon, CE, COL Nelson P. Conover, CE, COL Tilford C. Creel, CE, and COL Robert C. Lee, CE. Technical Director was Mr. F. R. Brown.

### CONTENTS

<u>P.</u>	age
PREFACE	1
CONVERSION FACTORS, US CUSTOMARY TO METRIC (SI)	
UNITS OF MEASUREMENT	3
PART I: INTRODUCTION	4
PART II: DATA BASE	6
PART III: DATA ANALYSIS AND MODEL FORMULATION	9
PART IV: SUMMARY	19
REFERENCES	20
TABLES 1-44	
FIGURES 1-57	

# CONVERSION FACTORS, US CUSTOMARY TO METRIC (SI) UNITS OF MEASUREMENT

US customary units of measurement used in this report can be converted to metric (SI) units as follows:

Multiply	By	To Obtain
feet	0.3048	metres
knots (international)	0.514444	metres per second

### THE ATCHAFALAYA RIVER DELTA

### WIND CLIMATOLOGY

#### PART I: INTRODUCTION

- 1. This task of the Atchafalaya Bay investigation is an investigation of the different wind conditions that typically influence the Atchafalaya Bay system. Wind fields play an important role in determining the hydrodynamics not only within the bay itself but also in those portions of the Gulf of Mexico that surround the bay. Along with astronomically induced currents and riverine discharges, wind-induced currents, as a result of shallow depths existing in the bay, dictate the response of many physical processes within this system. These processes include: (a) hydrodynamic circulation patterns, (b) flooding and drying of marshes, (c) dispersion and deposition of sediments, (d) distribution of fresh and salt water, and (e) dispersion of any pollutants that enter the system. In addition, winds dictate the wave climate in this area. Both short-period, locally generated waves as well as longer period waves, propagating into this region from the Gulf, have the capability to redistribute sediments. There is a definite need to both quantitatively and qualitatively describe the characteristics of the wind fields within this bay system. The role of the wind climatology task in the Atchafalaya Bay investigation is described by McAnally and Heltzel (1978).
- 2. Most conventional wind analyses rely only on an estimation of statistical parameters to describe the wind fields. These statistical estimates are usually derived from an analysis of a set of prototype wind data, normally speed and direction, under the assumption that all wind conditions belong to one homogeneous population of events. This kind of model can yield very useful information; however, additional insight can be gained by categorizing the circulation patterns associated with daily weather conditions into different synoptic scale events and then examining the wind fields associated with each type of event. In this way, a more homogeneous population of statistical events is assured. Byrne et al. (1977) used this approach in an analysis of the precipitation within this same Atchafalaya Bay system. Wax (1977) studied the variation of water-surface elevations as a function of the sequence of occurrence of different synoptic scale circulation patterns. This study combines

wind-field statistics with information on weather pattern variations in order to form a statistical-climatological model to describe wind conditions within the Atchafalaya Bay area.

3. In addition to a breakdown by synoptic type, the wind analysis is categorized into four seasons. The winter season is chosen to include December through February; the spring season, March through May; the summer season, June through August; and the fall season, September through November. The purpose of this additional discretization is to examine whether or not the frequency of occurrence of each synoptic type and its associated winds changes during a typical year. Evidence of such a seasonal variation was presented in the work by Byrne et al. (1977) and Wax (1977). It is important to note that any seasonal change indicates a climatic variation with a time scale which is larger than that of the synoptic events. The causes of this larger scale of change will not be examined in this study.

### PART II: DATA BASE

- 4. An extensive set of prototype data is required to relate certain sets of wind conditions to particular synoptic scale events within a seasonal framework. A minimum of 1 year of continuous data is essential for this type of study with additional years being optimal. Prior to October of 1981, sporadic wind data were recorded in the Atchafalaya Bay area, primarily for use in testing the weather stations and for use in other aspects of the overall project objectives. Beginning in October 1981 and extending through September 1982, a comprehensive plan to collect wind information within the bay was undertaken.
- 5. Weather stations were installed at the three locations shown in Figure 1. Exact elevations and locations of the anemometers, in terms of latitude and longitude, are shown in Table 1. Two anemometers were installed at each site, one at approximately a 5-m elevation and the other as close to 10 m as possible. The lower anemometer was installed primarily as a backup for the upper one. These three locations were chosen in order to obtain reasonable spatial coverage of the wind information gathered within the bay. The exact anemometer sites selected were dependent upon the availability of suitable structures upon which the weather stations could be installed. It was anticipated that this configuration could lend insight into the spatial variability of the winds within the bay. However, it also was anticipated that continuous data could not be recorded at all three locations. Therefore a second priority was to record as nearly a continuous record as possible at weather station E, since this station was more or less centrally located.
- 6. As expected, due to instrument failure, servicing difficulty resulting from logistics problems, and "unkind" weather, continuous data were not recorded at all three locations. Figure 2 shows, in a calendar format, all the data that were collected at each of the six anemometer positions. It should be noted that the secondary objective, continuous data at station E, was essentially met. During the winter and spring seasons, the availability of data was quite good; but for the other two seasons the amount of data taken was considerably less. By using some data taken during the summer of 1981, sufficient records were obtained for both of these seasons as illustrated by the following percentages.

Season	Percentage of Season During Which Usable Data Exist
Winter	96
Spring	92
Summer	52
Fall	69

- 7. Data taken at each anemometer included wind speed, wind direction, and the standard deviation of direction. All three of these measurements were computed using a 20-min average and were recorded every 20 min. Some of the data taken were not of a usable quality because the anemometers experienced a few mechanical problems during the course of the data-gathering experiment, particularly during electrical storms. Figure 2 also shows the times for which data were recorded but subsequently found to be erroneous. These times are designated by crosshatched areas.
- 8. Other sources of data were used in the formulation of the statistical-climatological model. In order to determine the long-term wind characteristics of this region, wind data recorded by the National Weather Service at Lake Charles and New Orleans, Louisiana, were used, since many years of data recorded at these two sites are available. Prior to the early 1960's, the locations and elevations of these anemometers changed frequently and it was decided to only use data recorded since that time; the anemometer at Lake Charles remained fixed at an elevation of 22 ft\* after May of 1962 and the anemometer at New Orleans remained at 20 ft after December of 1960. These data consisted of measurements of wind speed and direction every 1 or 3 hr and were available from the starting dates listed above through 1975. It was believed that 10 to 15 years of data was sufficient to determine long-term wind characteristics.
- 9. One other data source was used to illustrate the long-term trends of the wind fields within this area. These data consisted of wind observations reported by ships in the Atchafalaya Bay area and northern Gulf of Mexico adjacent to the bay. More precisely, they included observations within a 1-deglatitude by 1-deg-longitude block extending from 28 deg north to 29 deg north

 $<sup>\</sup>mbox{\ensuremath{\bigstar}}$  A table of factors for converting US customary units of measurements to metric (SI) units is presented on page 3.

and from 91 deg west to 92 deg west that were collected during the years 1956 to 1975. These observations were also compiled by the National Weather Service and were used in the study since they were readily available. It is important to note that these measurements, since many were visual observations, are far less accurate than those recorded by anemometers.

10. The final source of data used in this study was the publication "Daily Weather Maps" issued by the Environmental Data and Information Service of the National Oceanic and Atmospheric Administration (US Department of Commerce). This information was used to define the type of synoptic weather pattern causing the circulation of air within the study area on a daily basis. These maps were available for every day during the prototype data collection effort.

### PART III: DATA ANALYSIS AND MODEL FORMULATION

- 11. The first step in the formulation of the statistical-climatological model was to devise a method of categorizing day-to-day circulation into various synoptic scale types based on certain dynamical characteristics particular to each type. The classification criteria that were adopted follow those used by Muller (1977) and Byrne et al. (1977) but with some exceptions. They stratified the weather into eight all-inclusive categories. In this study, two of their types were combined and two others were modified, resulting in a total of seven.
- 12. Two of the synoptic types deal with the passage of fronts through the area. The first situation, called Frontal Overrunning (FOR), is associated with the passage of a southerly or southeasterly moving cold front. This event usually precedes the west to east progression of a high pressure cell across the continental United States and is characterized by a period of steady winds, usually from southerly directions, that is suddenly interrupted by a rapid increase in wind speed and a switch in direction to winds from the north (Figure 3). Figure 4 shows the time-history of winds encountered during the passage of that cold front over the Atchafalaya Bay.
- 13. A second synoptic event associated with frontal activity is the Frontal Gulf Return (FGR) situation and is characterized by a northerly moving warm front or by the progression of a weak-to-moderate strength low pressure cell from southern Texas toward the northeast. Steady wind flow prior to the frontal passage is broken by a change in wind direction to winds from southeast through southwest directions. This change in wind direction and speed can often be quite abrupt especially in the case of a low pressure cell passage. An example of a more gentle shift in winds associated with a warm front passage is shown in Figures 5 and 6.
- 14. Both of these passages also cause noticeable changes in the quality of the air itself. Overrunning of a cold front brings colder, dryer air from the north into the area while a warm front brings with it warmer, moister air from the northern Gulf. The time scales associated with these two phenomena are about the same; each usually lasts between 6 and 24 hr depending upon the forward speed of the front.
- 15. Another type of circulation that can have high wind conditions with widely changing direction is the infrequent passage of an intense low pressure

center called a Gulf Tropical Depression (GTD). This weather type is comprised of a small low pressure center with tightly spaced isobaric contours and high winds and occasionally affects this area, especially during the late summer months. Hurricanes and intense tropical storms fall into this category. These events are considered separately due to their intensity and infrequent occurrence.

- 16. The remaining synoptic types are associated with the movement and position of high pressure cells relative to the continental United States and have time scales ranging from hours to days. The first of these types to be considered is called a Continental High (CH) situation. In this study, this type will include both the CH and Pacific High (PH) patterns classified by Byrne et al. (1977). Airflow during these conditions is steady and directed from northwest through northeast directions. Fair, dry, cooler air generally characterizes this circulation which is associated with a clockwise rotation around a high pressure center located over the central and western United States or southern Canada. Figure 7 shows this synoptic pattern, and Figure 8 shows the associated time-history of wind speed and direction measured in the Atchafalaya Bay during this event.
- 17. A second type, Coastal Return (CR) flow, occurs when these same high pressure systems reach the eastern coast of North America or even move out into the western reaches of the Atlantic Ocean. With this progression, the winds shift from northerly directions to those with primarily eastern components. During winter and spring, this circulation pattern represents colder air modified by short passages over the Atlantic Ocean and northern Gulf as shown in Figure 9. During summer and fall, this includes the Bermuda High situation in which a band of tropical air extends eastward from the southeastern United States far into the Atlantic Ocean; and again, flow is essentially from the east. Figure 10 shows typical wind conditions during a winter coastal return flow of air.
- 18. When a high pressure system moves still farther eastward out over the Atlantic Ocean, the winds shift from the east to the south resulting in what is called Gulf Return (GR) airflow. These winds bring moist tropical air from the Gulf of Mexico into the area. GR circulation is illustrated in Figure 11 and the accompanying prototype observations are shown in Figure 12. This particular air circulation also can occur after the passage of a warm front or low pressure center once the winds have become fairly steady.

- 19. The final synoptic scale weather pattern to be considered is the Gulf High (GH) condition in which a high pressure cell is displaced over the northern Gulf. During the eastward passage of this "high," winds are usually rather weak and change in direction from northwest to west to southwest and they bring warmer, dryer air from southern Texas and Mexico. An example of this weather pattern is shown in Figure 13 and the measured winds are shown in Figure 14.
- 20. Having broken the weather patterns into separate types based on dynamical considerations, the next step was to determine which synoptic pattern was influencing the Atchafalaya Bay area during the times that data were recorded. This was done using daily weather maps in conjunction with measured data. Sudden changes in wind speed and/or direction as well as changes in the standard deviation of wind direction were used to define the transition from one weather type to another. Using all available prototype data, the frequency of occurrence of each synoptic scale type, within the seasonal framework mentioned earlier, was computed. These estimates are shown in Table 2. Results found by Byrne et al. (1977) for New Orleans and Lake Charles can be grouped into this seasonal framework for comparison. Their findings also are shown in Table 2.
- 21. A few points should be remembered when comparing the results of Bryne et al. (1977) with those of this study. Differences in definitions of the synoptic types (defining the beginning and end of FOR and FGR frontal weather patterns) make comparisons difficult. In addition, the PH circulation type described by Byrne et al. (1977) has been combined with the CH type. Despite these differences, many conclusions about the variation in circulation affecting this area can be formulated and are shown to be substantiated by both sets of results.
- 22. Winter is dominated by the CH and FOR circulation types which are both characterized by northerly winds. During the 1981-82 winter season, there seemed to be about 10 percent less of these northerly wind conditions than "usual" (49 percent in the Atchafalaya Bay versus 60 percent and 59 percent using the long-term data at Lake Charles and New Orleans). The difference was made up by an increase in CR and GR circulation. Note the sum of the frequencies of FGR and GR for all three data sets. They are nearly identical and indicate that the occurrence of this kind of airflow was about normal for this season. The discrepancies in the individual frequencies of

occurrence of FGR and GR are due to differences in determining the start and end of individual weather patterns.

- 23. During the spring season, the difference in northerly winds between 1982 and typical years is even more obvious and is again compensated by a large increase in CR circulation and a slight increase in the GR-FGR type of airflow. As with the winter season, the GH and GTD activity during the spring is very small.
- 24. This trend of decreasing northerly winds continued into the summer of 1982. Once again, the CR circulation has increased as has the airflow into the area from the Gulf. Results during the summer of 1982 may be biased because of the lack of data taken over the entire season. Compared with the long-term results, there seems to be about a 15 percent increase in winds from the south as well as a 5 percent decrease in winds associated with the GH circulation. In addition, GH airflow only becomes important during the summer and the frequency of occurrence of GTD's during this one summer is less than usual.
- 25. The fall season of 1981-82 was very typical in terms of the distribution of the occurrence of the different synoptic weather patterns. The percentages of CH and FOR wind conditions as well as CR and GR-FGR circulation are almost identical with the percentages found using the results of Byrne et al. (1977). The only major difference is a decrease in the frequency of occurrence of the GTD events.
- 26. In general, the 1981-82 year in the Atchafalaya Bay area experienced fewer than normal northerly wind conditions. This decrease was accompanied by slight increases in both easterly and southerly winds, and it appears that high pressure systems were able to move across the continent faster. The other major difference, mentioned earlier, is the lack of occurrence of already rare events—the intense tropical storms.
- 27. Up to this point the "climatological" aspects of the model have been discussed. These include the categorization of weather circulation into separate synoptic types and an analysis of the frequency of occurrence of each type during the year as a whole and during the four seasons that comprise the year. The remainder of the report will deal with wind data that were recorded, manipulation of the data, statistics that were computed, and relations between statistical findings and "climatological" information. One point must be addressed that deals with a key assumption of the model.

- 28. Due to the size of the Atchafalaya Bay, in relation to the spatial extent of all the synoptic weather types except tropical storms, which will not be discussed very much due to their infrequent occurrence, wind conditions over the entire area were assumed to be relatively constant in terms of both wind speed and direction. Figures 15 and 16, and 17 and 18, show a comparison of the measured wind speeds and directions at weather stations E and B, and E and D, respectively. Winds at each site were recorded on the upper anemometers, which are at elevations within 1 ft of each other. In both instances, wind speeds are within 2 m/sec or 4 knots of each other. Wind directions at E and D are within 20 deg of each other and the direction measurements at E and B are within about 40 deg of each other. There is some evidence that these directional differences may represent a bias. Nevertheless, differences were assumed to be negligible and the supposition of constant wind fields over the entire Atchafalaya Bay was assumed to be true.
- 29. A vast amount of data was used to generate wind statistics for the Atchafalaya Bay area. Table 3 shows a breakdown, by season, of the number of paired observations of wind speed and direction that were utilized. Statistical information derived from data recorded at Lake Charles, New Orleans, and from ships observations can only be used in a gross sense, such as testing whether or not the distributions of wind speed and direction measured during the 1 year in the Atchafalaya Bay reflect "typical" conditions for this area. Without relating the long-term data directly to the atmospheric circulation on a day-to-day basis, the long-term data can only be used to make inferences about the relations between gross wind statistics measured during 1981-82 and the long term.
- 30. Before analysis, the data were adjusted to specified formats. Long-term data at the National Weather Service stations were recorded in whole knots and in direction bands for wind speed and direction, respectively. Sixteen bands were used with 22.5 deg in each band. Band number 1, which is centered about 0.0 deg, indicates winds blowing from due north; and increasing bands in the clockwise direction denote increasing wind angles. For example, band number 5 indicates winds blowing from the east at 90 deg and band number 9, winds from the south at 180 deg.
- 31. Data recorded in the Atchafalaya Bay during 1981-82 were measured in decimal metres per second and decimal degrees. They also were recorded at various elevations dependent upon the anemometer location. Therefore, for

comparison purposes, wind speed data were all converted to a common elevation (10 m) using the following expression given in the Shore Protection Manual (US Army Coastal Engineering Research Center 1977);

$$U_{10} = \left(\frac{10}{Z}\right)^{1/7} U_{Z} \tag{1}$$

where  ${\rm U}_{10}$  is the equivalent 10-m wind speed for a measured wind speed,  ${\rm U}_{\rm Z}$ , recorded at an elevation Z (where Z is in metres). The equivalent 10-m wind speed was then converted to knots. The measured wind directions were assumed to be invariant between the measured elevation and the 10-m elevation and were converted to the 16-band scale mentioned above. In this way, comparisons between statistics computed from the measured data set and the long-term results could be made more easily.

- 32. The data at Lake Charles were measured at an elevation of 22 ft (6.7 m), and the data at New Orleans, at 20 ft (6.1 m). According to Equation 1, wind speeds at Lake Charles are 6 percent less than their equivalent 10-m values and at New Orleans they are 7 percent less. In the statistical tables for these two sites (Tables 4-11), no compensation for elevation was made in order to avoid difficulties in the presentation of results. However, in the plot comparisons (Figures 19-28), long-term results were properly shifted along the velocity axis. Wind directions at Lake Charles and New Orleans were again assumed not to be a function of elevation. Ships observations, since they had no vertical control, were analyzed and presented as recorded.
- 33. Tables 4-15 show the bivariate distribution of wind speed and direction, by season at New Orleans, Lake Charles, and Atchafalaya Bay. Figures 19-26 show comparison plots of the one-dimensional frequency of occurrence of both speed and direction, again by season, and include analysis results using ships observations.
- 34. The frequency of occurrence of wind speeds from all three long-term sources show spikes at multiples of 5 knots. This is obviously due to the fact that a number of recorded observations (such as most of the ships observations) were made by visual inspection or guesswork and not from instrument recordings. For these same ships observations, it is apparent that for wind direction a rougher 8-point scale was utilized most of the time. Therefore results inferred from this particular data source should be used only to

describe generalized trends in the data and not for any detailed analysis.

- 35. There also appear to be different thresholds for lower values of wind speed recorded at the National Weather Service stations. Nothing less than 1 knot was recorded at New Orleans and nothing less than 2 knots was measured at Lake Charles. These differences, since they occur at the lower end of the wind speed spectrum, are not of great importance. A problem also appears to exist with the anemometers used in the Atchafalaya Bay. They do not seem to be able to differentiate wind directions from about 5 to 10 deg on either side of due north. This is apparent in the data analysis from the lack of observations in direction band 1. This discrepancy must be recognized since it does pose a problem.
- 36. Referring to the comparison plots, Figures 19 and 20 in particular, it appears that the 1981-82 winter season was quite typical in terms of the wind speeds encountered; however, in terms of direction, there are some differences. There is a decrease in winds from the northernmost directions and an increase in the frequency of winds from the east through the southeast directions. This supports earlier observations concerning the differences in the occurrence of the synoptic weather types between the 1981-82 measurement year and the long-term results.
- 37. During the spring season of 1982, winds in the Atchafalaya Bay area were not as strong as typically expected except for very large wind speeds, above 20 knots. Again in the wind direction plot, there is a decrease in northerly winds and an increase in easterly winds over "typical" conditions. The large "spike" of winds from the south measured at New Orleans and Lake Charles is not explainable. This also was apparent during the winter season. It might be a remnant of visual observations. Also, note the shift in dominant wind directions from northerly winds to southeasterly winds which again is supported by the findings concerning changes in the frequency of occurrence of the synoptic scale events from winter to spring.
- 38. During the summer of 1982, wind speeds again seem to be quite normal although their magnitude, in general, has substantially decreased compared with those typical of winter and spring seasons. Winds also are fairly normal in terms of their directional distribution and are shifting from the southeast to the south, which is accompanied by a decrease in winds from the north. Note the effect on wind direction as a result of GH circulation (increased westerly winds).

- 39. Wind speeds measured during the fall of 1981 and 1982 are slightly larger than those to be normally expected, as indicated by an increase in winds in the 10- to 15-knot range. The distribution of wind direction is very typical for this season which again agrees with the information presented in Table 2. Winds, in general, also are shifting back to northeasterly directions as a result of an increase in CH and FOR circulations.
- 40. Figures 27 and 28 show the one-dimensional frequency of occurrence of both speed and direction for the entire year in Atchafalaya Bay as well as "average years" in this region, as represented by the cumulative longer term statistics computed from New Orleans and Lake Charles data. Wind speeds measured in the bay during 1981-82 are typical but the distribution of wind direction is slightly different. These cumulative statistics, without regard for synoptic variation, can be very useful in: (a) understanding the overall effects of the seasonal variation in wind conditions, (b) substantiating findings in the Atchafalaya Bay using longer term data, and (c) estimating wind conditions that can be expected in the future.
- 41. Tables 16-40 contain the bivariate distributions of wind speed and direction, by season and by synoptic weather type, for data recorded in the Atchafalaya Bay during the 1981-82 measurement year. These tables are designed to show wind conditions associated with each of the synoptic circulation patterns defined earlier. They also can be used to examine the seasonal variation of wind conditions for a particular synoptic event. Care must be exercised when applying conclusions drawn from these tables to times other than the measurement year. Caution should be used concerning the frequency of occurrence of that synoptic type during a particular season and how that frequency compares with the expected frequency during the same season in other years. In addition, especially during the summer and spring seasons, it is important to note the number of wind speed-direction pairs that were considered in a particular bivariate distribution. Results presented can be very useful; however, their limitations should be understood. The tables are grouped in the following manner. There are four tables for each synoptic type except the Gulf Tropical Depression (there is only one table). The order of synoptic types is: Continental High, Frontal Overruning, Coastal Return, Gulf Return, Frontal Gulf Return, Gulf High, and Gulf Tropical Depression. Four tables for each type, one for each season, are ordered in this way: winter, spring, summer, and fall.

- 42. Tables 16-19 are for the CH circulation pattern. During all four seasons this airflow is characterized by potentially high winds out of the northwest through the northeast directions. The highest winds occur during winter and spring and they drop off in magnitude during the summer. During all four seasons, the dominant direction is from the northeast. Tables 20-23 show the bivariate distributions for FOR conditions that are again characterized by high winds which can be quite variable in both speed and direction. Dominant directions also are from the north or northeast and are associated with winds during the latter stages of the frontal passage. Again, like the CH situation, during the summer the magnitudes of the winds during this weather type are smaller.
- 43. Circulation during the next type considered, the CR flow, is from the northeast through the southeast directions with the dominant direction being from due east. Accompanying this type of weather pattern are winds that are steady in both speed and direction. Winds can be quite large. A decrease in wind magnitude also is apparent during the summer and fall months. The seasonal variation in this synoptic type is shown in Tables 24-27.
- 44. The GR flow, shown in Tables 28-31, is characterized by steady winds from the east through the southwest directions. In general, wind speeds are less than those found during coastal return airflows. There does not appear to be a decrease in wind speed during the summer for this type, which was the case for previous types.
- 45. Wind conditions during FGR flow, in Tables 32-35, are characterized by variable winds changing to steady winds blowing from the southeasterly through the southwesterly directions. These steady winds appear to be of low to moderate speeds except during the winter when they can become quite strong. In addition, variable winds often experienced during the initial stages of a warm front passage can be large.
- 46. GH circulation, shown in Tables 36-39, typically involves a rotation of winds from north to west to the south as the high pressure cell, which is centered in the northern Gulf, migrates eastward. The dominant wind direction is out of the southwest and winds are usually light. As mentioned earlier, this type of circulation only becomes prevalent during the summer months.
- 47. The remaining circulation pattern is the GTD. Table 40 shows the winds that accompanied the single intense low pressure system which passed

through this general area and was recorded. In this instance, winds were not as variable as they might have been had the center of the system passed directly over the bay. Here winds were predominantly from the south, which indicates passage of the system to the west of the study area. The important feature of this synoptic type is the potentially high winds that can accompany these intense storms.

- 48. The series of Figures 29-53 are in the same order as the previous series of tables. These plots contain estimates of the duration of wind speed by synoptic type and by season where duration is defined to be the persistence of a certain wind speed once that wind speed has been exceeded. There are two curves on each plot, one for the average duration and one for the maximum duration encountered. The same cautions should be remembered when using these results as were given for the bivariate information. These duration statistics can be very helpful in determining estimates of the average and maximum durations of a particular synoptic type during a certain season as well as the durations of different wind speeds during a particular type of circulation. Figures 54-57 illustrate the duration statistics computed for each season, but in this case all the synoptic weather patterns are combined.
- 49. The final information presented is a list of wind speed maxima by direction band and by season using the long-term data at New Orleans and Lake Charles (shown in Tables 41-44). Extreme tropical storms have been deleted from this data set; however, this information is useful in determining the expected extreme wind conditions during a year in terms of direction for the other synoptic types. These can be compared with those extremes measured during the data-gathering experiment in the Atchafalaya Bay to determine whether or not typical extreme wind conditions associated with the synoptic weather types were measured. A frequency of occurrence analysis also could be performed on these wind speeds in order to determine future extreme values.

### PART IV: SUMMARY

- 50. Wind data were gathered extensively within the confines of Atchafalaya Bay for a period of 1 year, which started in October 1981 and continued until October 1982. The amount of usable data was extremely large during the winter and spring seasons and sufficient during the summer and fall of 1982. A large amount of data from other sources, namely the National Weather Service stations in Lake Charles and New Orleans and a compilation of observed-fromships wind conditions in this immediate area, was assembled and analyzed. Statistical estimates were used to express wind conditions in terms of speed and direction. These statistics were then related to the different kinds of weather patterns that influence this region by categorizing these patterns into seven all-inclusive types with each defined by a particular set of dynamical characteristics. In this way, the variations in wind conditions were better related to the climatological forcing that induced these variations.
- 51. Information was provided on the frequency of occurrence of wind conditions both with and without regard to their causative factors. The latter results were used to substantiate those results found using the statistical-climatological approach. Estimates of the duration of these different weather patterns as well as estimates of the persistence of winds common to each pattern were computed. Finally, information concerning extreme wind conditions that can be expected to occur in any given year was presented using the long-term historical data that were available. Using this information, the variations in the physical processes within the bay that are strongly dependent on the local wind conditions can be better understood and predicted.

#### REFERENCES

- Byrne, P., et al. 1977. "Climatology, Hydrology and Hydrography of the Vermilion Basin," Final Report to the State Planning Office, Baton Rouge, La.
- McAnally, W. H., Jr., and Heltzel, S. B. 1978. "A Plan for Predicting the Evolution of Atchafalaya Bay, Louisiana" (unpublished), US Army Engineer Waterways Experiment Station, Vicksburg, Miss.
- Muller, R. A. 1977. "A Synoptic Climatology for Environmental Baseline Analysis: New Orleans." Journal Applied Meteorology, Vol 16, No. 1, pp 20-23.
- US Army Coastal Engineering Research Center. 1977. <u>Shore Protection Manual</u>, Department of the Army, Corps of Engineers, Fort Belvoir, Va.
- US Department of Commerce. "Daily Weather Maps" (published weekly), National Oceanic and Atmospheric Administration, Environmental Data and Information Service, Washington, DC.
- Wax, C. L. 1977. An Analysis of the Relationships Between Water Level Fluctuations and Climate of Coastal Louisiana, Ph. D. Dissertation, Louisiana State University, Baton Rouge, La.

Table 1

Exact Locations and Elevations of the Atchafalaya Bay Anemometers

	Elev	ation		***************************************
Station	ft	m	Latitude	Longitude
WS-B(U)	29.0	8.85	29°36'N	91°42'W
WS-B(L)	19.0	5.80	29°36'N	91°42'W
WS-D(U)	28.0	8.55	29°27'N	92°25'W
WS-D(L)	16.5	5.05	29°27'N	91°25'W
WS-E(U)	28.5	8.70	29°23'N	91°32'W
WS-E(L)	23.0	7.00	29°23'N	91°32'W

Table 2
Frequency of Occurrence, in percent
by Synoptic Weather Type

Location	Season	PH	СН	FOR	CR	GR	FGR	GH	GTD
Atchafalaya Bay	Winter		29	20	15	22	9	4	0
(1981 <b>-</b> 1982)	Spring		18	10	22	44	5	1	0
	Summer		7	11	14	39	7	23	0
	Fall		33	22	16	19	3	5	3
	Year		22	16	17	31	6	8	1
				···					
Lake Charles (1971-1974)	Winter	11	22	27	10	15	16	0	0
	Spring	9	22	14	10	26	16	2	0
	Summer	0	17	8	11	24	5	32	4
	Fall	7	28	19	15	15	9	2	8
	Year	7	22	16	12	20	11	9	3
New Orleans	Winter	6	25	28	9	13	17	2	0
(1966-1975)	Spring	6	20	15	11	27	18	3	0
	Summer	0	17	6	12	23	8	27	7
	Fall	4	34	15	16	12	10	4	6
	Year	4	25	16	12	18	13	9	3

Table 3

Number of Wind Observations

(Speed and Direction Pairs)

Location	Winter	Spring	Summer	<u>Fall</u>	Total
Lake Charles* (1962-1975)	12,391	13,325	13,069	13,237	52,022
New Orleans* (1960-1975)	20,174	20,921	18,671	17,709	77,475
Atchafalaya Bay**					
(1981-1982)	12,170	11,344	3,847	6,371	33,732
Ship observations	4,438	4,713	5,274	4,708	19,133

<sup>\*</sup> Hourly and 3-hourly observations.

<sup>\*\*</sup> Twenty-min observations.

Table 4

**NEW ORLEANS** 

# BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASO	ON C	DEC T	O FEB	)							20	174 H	OURLY	OBSE	RVATIO	N PAIRS
WIND SPEED							DIF	RECTIO	N BAN	4D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
1 2 3 4 5 6 7 8 9 10 11 12 11 14 15 17 18 19 22 22 24 25 26 27 28 29 30 31 33 34 35 36 37 38 39 40 30 31 31 31 31 31 31 31 31 31 31 31 31 31	149632444522442057220000000000000000000000000	0666207824616179860500000000000000000000000000000000000	130 57635 1998388666 111613010000000000000000000000000000	001118675101205331000000000000000000000000000000	057955397686868541000000000000000000000000000000000000	01113681690794624110000000000000000000000000000000000	102875175639129620010000000000000000000000000000000	08710813914342860724100000000000000000000000000000000000	03121573766443221 03121573766443221	043754458774342969110000200000000000000000000000000000	155485412556316621010000000000000000000000000000000	06657853061005323100000000000000000000000000000000	1360497702205174541200000000000000000000000000000000000	0982067664228096540200000000000000000000000000000000000	0784284184150490120000000000000000000000000000000000	0749574545716952224601000000000000000000000000000000000	14 299 541 787 787 932 1061 1018 1007 766 875 538 655 433 303 270 169 113 99 114 55 30 0 0 0 0 0 0 0 0 0 0 0 0 0 0

TOTAL 1213 900 874 881 664 521 510 593 722 429 372 304 424 348 451 786

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 5

### **NEW ORLEANS**

# BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEASO	1 ) אכ	MAR TO	D MAY	)							20	921 H	URLY	OBSE	RVATIO	N PAIRS
WIND SPEED							DI	RECTI	ON BAI	ND							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
1 23 45 67 89 111 115 117 119 112 114 117 119 112 114 117 119 119 119 119 119 119 119 119 119	01222724811125574320100000000000000000000000000000000000	02503553021898561100000000000000000000000000000000000	0606623030042944010000000000000000000000000000000	0177412468595131322000000000000000000000000000000000	000448845539586512100000000000000000000000000000000000	09291212487264731111100000000000000000000000000000000	063425279252401046020200000000000000000000000000000	0684693334781223016450200000000000000000000000000000000000	02239223151991112864835120822100000000000000000000000000000000	008344535370111786502010010000000000000000000000000000000	0105426504180763110000000000000000000000000000000000	060409696144198432000000000000000000000000000000000000	0980726045647075452001000000000000000000000000000000000	0478998500119096450000000000000000000000000000000000	07884975257189956971010000000000000000000000000000000000	0384084414854118820000000000000000000000000000000000	4257578810359971616182319736971322400000000000000000000000000000000000

TUTAL 780 602 592 595 519 684 808 1048 1287 645 543 386 394 314 322 474

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.6 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 6

### **NEW ORLEANS**

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEAS	ON (	JUN T	O AUG	• )							18	671 F	OURLY	OBSE	RVATI	ON PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DI 7	RECTI 8	ON BA	ND 10	11	12	13	14	15	16	TOTAL
1 2 3 4 5 6 7 8 9 10 112 12 13 14 15 16 17 18 19 12 22 22 22 22 22 22 23 33 33 33 33 34 35 40 37 38 40 37 37 37 37 37 37 37 37 37 37 37 37 37	0669011197198320010000000000000000000000000000000000	1067867987165322010000000000000000000000000000000000	18660 8107720010000000000000000000000000000000	282319506690652000000000000000000000000000000	18278613856128062112000100000000000000000000000000	077306755441588833010000000000000000000000000000000	0092462673991632100000000000000000000000000000000000	01387671011622111000000000000000000000000000	0 26 5 38 6 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	02256549808795321000010000000000000000000000000000000	121191514435847165533102000000000000000000000000000000000	0831113988648471686311101000000000000000000000000000000	00377000000000000000000000000000000000	0 9 9 1 0 9 9 1 0 9 9 1 0 9 9 1 0 9 9 1 0 9 9 1 0 9 9 9 1 0 9 9 9 9	060811742805100000000000000000000000000000000000	13381 77199056431 156431 156431 156431 156431 156431 156431 156431 156431 156431 16643	9 693 1221 1557 1958 4113 996 4113 1996 1000 0000 0000 0000 0000 0000
IOIAL	117	220	0/2	221	400	441	454	56/	840	758	919	784	831	554	359	507	

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 7

### NEW ORLEANS

# BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SEASON ( SEP TO NOV ) 17709 HOURLY OBSERVATION PAIRS																	
WIND SPEED							DIE	RECTIO	IN RAI	ND							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 40 40 40 40 40 40 40 40 40 40 40 40	1597 1130 1130 1130 1130 1130 1130 1130 113	1257954209 1135597113209 11320 1132	153992428796808711010000000000000000000000000000000	196555872532420000000000000000000000000000000000	1449903878075251087310000000000000000000000000000000000	05198430270255701000000000000000000000000000000	0139606197195451120000000000000000000000000000000000	09574834253222200381101000000000000000000000000000	082638563800458231120001000000000000000000000000000000	021200541376352000000000000000000000000000000000000	133321187370111110000000000000000000000000000	3442113295552200000000000000000000000000000000	159977688084235200000000000000000000000000000000000	311515896687652000000000000000000000000000000000000	43222212111 432222212111	231967768785396653000000000000000000000000000000000	160 5530 10933 11758 11058 6776 3816 2184 1134 1258 1200 1200 1200 1200 1200 1200 1200 120

TOTAL 1194 1037 1336 1279 789 656 539 460 425 239 214 227 388 300 308 600

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 8

#### LAKE CHARLES

# BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASO	) N ( I	DEC T	O FEB	)							12	391 H	OURLY	OBSE	RVATIO	N PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	D18	RECTI 8	ON BAI	10	11	12	13	14	15	16	TOTAL
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 120 221 223 224 225 227 228 230 311 323 334 335 336 337 338 340	0 0 4 6 9 3 1 1 0 3 8 1 1 0 3 8 1 1 0 3 8 1 1 0 9 7 1 8 3 2 2 2 3 6 4 6 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	006788619210463650552422200000000000000000	001579936499996459812000000000000000000000000000000000000	00045758066471184224021000000000000000000000000000000	00197721776632111000000000000000000000000000000000	00 165 890 177 462 1352 1200 000 000 000 000 000 000	008978223579414485420200000000000000000000000000000000	009150303580100290140101000000000000000000000000000	00 196661239082100010000000000000000000000000000000	0007501224498438521200000000000000000000000000000000000	0140603971634584111100000000000000000000000000000000	00239519954174644000000000000000000000000000000000	0080777950362878200100000000000000000000000000000000	0083138911028885321000000000000000000000000000000000000	005697339932843748502200000000000000000000	0 0 8 8 6 4 4 2 1 7 4 9 8 0 3 6 6 6 9 0 4 6 1 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	01 26055 1092244572258227796597442100000000000000000000000000000000000

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

THE WIND SPEED IS THAT MEASURED AT A 6.7 METER ELEVATION.

TOTAL 1229 774 787 995 753 689 794 594 1046 418 364 233 219 229 310 557

Table 9

LAKE CHARLES

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEAS	ON (	MAR T	O MAY	)							13	325 H	IOURLY	OBSE	RVATI	ON PAIRS
WIND SPEED							ומ	PECTI	ON BA	מא							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	0	0	0	. 0	0	0	0	0	0	0	_ 0	0	0	0	0	0	0
3	21	16	16	16	24	16	14	11	19	. 8	17	- 9	14	- 9	15	- 9	240
4 5	61 73	36 44	56 64	81 99	60 99	57 75	76 115	57 99	103 128	41 54	40 57	33 32	35 38	27 29	29	25	824
6	42	36	53	84	94	72	93	91	144	60	63	32 21	21	18	33 26	33 19	1078 943
7	54	32	51	95	86	59	105	89	165	90	63	25	15	15	19	23	992
8	47	46	42	69	63	52 52	72	96	182	<b>82</b>	51	15	15	12	23	24	898
9	30	21	20	45	42	36	40	60	152	72	43	21	-8	7	14	17	636
10	58	47	48	74	58	48	79	92	249	112	62	15	12	12	18	26	1017
11	30	18	27	29	29	15	29	70	126	60	26	10	4	3	4	14	501
12	41	33	28	38	21	33	45	85	190	90	41	6	5	2	11	18	692
13 14	33 24	24 17	26 14	30 18	19	16 16	45 28	68 71	170 115	75 55	33	9 3	3 3	2 2	15	15	587
15	19	8	14	19	6 8	15	27	64	126	60	30 17	4	3	4	10 7	12 18	432 420
16	iź	8	6	12	4	5	12	30	57	34	5	i	3	i	ıí	9	216
Ĩ7	îī	7	ĭ	-6	3	3	12	29	47	28	7	ī	ŏ	ī	- 6	6	173
18	6	8	7	5	2	2	15	24	40	10	8	ī	Ŏ	ī	3	6	145
19	0	1	0	2	1	0	3	5	9	3	2	0	0	0	1	Ö	33
20	1	4	1	Ō	1	3	10	9	27	8	3	0	0	2	6	3	84
21	0	0	0	0	Ō	0	1	3	6	2	0	0	0	0	2	0	17
22 23	0	0	0	0	1 0	0	3	6	3	Q	0	0	0	0	0	0	18
24	ŏ	0	Ů	1 0	ŏ	ĭ	2	1 2	3 3	0	Ů	0	0	0	0	0	13 9
25	ň	ŏ	ŏ	ŏ	ň	Ď	ŏ	3	2	1	ŏ	Û	Ů	0	ň	Ü	9
26	ŏ	ŏ	ŏ	ŏ	ŏ	ĭ	ĭ	2	ī	ō	ŏ	ŏ	ŏ	Ö	ň	ň	6
27	Ó	0	Ö	Ŏ	Ō	ō	ō	ō	ō	Ŏ	ŏ	Ŏ	Ŏ	ŏ	ŏ	ŏ	ž
28	0	0	0	0	0	0	0	1	1	G	0	0	0	Ó	0	Ö	3
29	0	0	Ō	0	Ō	Ō	0	0	0	0	0	0	0	0	0	0	0
30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
31 32	U	0	0	0	0	0	0	0	0	0	Ŏ	0	0	0	0	0	0
33	Ŭ	Ü	0	ů	Ü	Û	0	0	0	0	0	0	0	0	0	0	0
34	ň	ů	ŏ	ŏ	ñ	ñ	ŭ	ů	ñ	Ů	Ô	Ô	Ü	n	n	0	Ů
35	ŏ	ŏ	ŏ	ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ň	ñ	ŏ	ŏ	Õ	ñ	ů
36	Ó	Ó	Ó	Ō	Ŏ	Ō	Ŏ	Ŏ	Ŏ	Ŏ	ŏ	ŏ	Ö	ŏ	ŏ	ŏ	ŏ
37	0	0	0	0	0	0	0	0	0	Ó	Ó	0	0	Ō	Ŏ	Ŏ	Ö
38	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
39	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
40	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL	572	413	482	730	631	532	841	1079	2081	956	574	214	185	157	260	285	

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 10

### LAKE CHARLES

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEAS	ON (	JUN T	O AUG	)							13	069 H	OURLY	OBSE	RVATIO	N PAIRS
WIND SPEED							DI	RECTI	ON BA	AND							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
	1 0 0 499 1035 408 226 136 6 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 0082153431200001000000000000000000000000000000	3 0043305570222831000000000000000000000000000000000	4 00050913679682519000000000000000000000000000000000000	5 00 1374 11190 6394137 315000000000000000000000000000000000000	6 00247121110674442379342000000000000000000000000000000000000					1 0056574000566544100000000000000000000000000000	12 00 284 1046 1160 466 1302 0000 0000 0000 0000 0000	13 00 356 126 646 1122 5320000000000000000000000000000000000	14 0087559443167700000000000000000000000000000000000	15 003801 4266 160020000000000000000000000000000000	16 007699263732100000000000000000000000000000000000	TOTAL  0 3 550 1872 1970 1414 1112 857 513 695 270 318 1699 777 34 110 0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
TOTAL	373	321	577	807	762	562	642	605	1379	1094	1001	508	474	378	307	202	

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 11

### LAKE CHARLES

# BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL S	SEASON	( SE	P TO	C VON								13237	7 HOUI	RLY O	BSERV	ATION	PAIRS
WIND SPEED							חזו	PECTI	ON BAI	ND							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	Ö	ŏ	ŏ	Ö	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ì
3	62	33	41	53	37	30	19	15	24	9	7	9	8	19	20	18	410
4	140	95	179	190	154	71	114	71	96	52	42	27	40	39	53	56	1427
5	130 94	100 95	172 138	221 191	182 167	115 76	120 120	86 66	111 102	62 50	49 37	33 17	34 16	27 12	52 37	63 45	1563 1269
6 7	82	91	133	132	129	88	80	56	77	50 50	38	8	19	9	24	38	1062
8	79	76	104	120	91	76	73	43	75	34	25	15	îí	13	18	30	890
9	52	51	71	64	68	64	59	37	43	25	14	6	5	4	8	25	602
10	78	94	112	88	64	71	87	40	68	24	32	6	8	6	16	36	838
11	44	41	42 65	43	25	27 40	37 43	20 30	21 28	15 20	9 5	6 2	1 0	1 6	10 13	21 21	370 449
12 13	51 37	51 29	30	36 26	29 14	15	27	21	27 27	20	6	3	3	2	15	27	309
14	27	22	20	15	- 6	14	19	19	31	10	6	4	3	3	-6	14	226
15	32	18	14	15	6	10	17	17	15	9	3	1	ī	2	6	10	182
16	18	9	3	12	6	. 6	12	6	13	3	0	0	1	1	5	9	112
17	16 13	10 9	7 6	6	2	12 6	3 4	6 5	12 10	3 6	0 1	0	0 2	0	1 2	8 9	90 87
18 19	5	0	Ö	Ö	2 5	3	ĭ	3	1	1	Ö	ŏ	í	ŏ	ő	3	29
źó	5	2	ĭ	4	ž	ŏ	ż	3	5	ż	ž	ŏ	ō	ŏ	ŏ	7	42
21	1	Ō	Ō	Ó	Ō	Ō	0	1	Ō	0	0	Ô	Ö	Ó	0	0	7
22	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	9
23 24	0	0	0	0	0	0	0	0	0	0	0 1	0	0	0	0	0	9 7
25	Ö	ő	ŏ	Ö	ŏ	ŏ	ŏ	ŏ	ŏ	ň	2	Ö	ŏ	ŏ	ŏ	ň	4
26	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ō	ŏ	ŏ	ŏ	ŏ	ŏ	ò
27	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
28	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
29 30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	U
31	ŏ	Ö	0	ŏ	Ď	Ö	ŏ	ŏ	ŏ	ň	ŏ	ň	ŏ	Ď	Ö	n	ň
32	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ
33	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
34	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35 36	0	Ü	0	0	0	0	0	0	0	0	0	0	0	0	0	U	U
36 37	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	0
38	Ŏ	Ŏ	Ō	Ō	Ŏ	Ō	Ō	Õ	Ö	Ŏ	Ō	Õ	Ō	ŏ	Ŏ	Ŏ	Ō
39	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Õ
40	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

TOTAL 976 837 1146 1233 1001 732 846 556 769 407 290 143 159 153 295 451

WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND)
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 12

### ATCHAFALAYA BAY

# BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASO	N ( D	EC TO	FEB	)							121	70 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIR	ECTIO	N RAN	מי							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-1 1-23 3-45 5-6-7 7-8 8-10 10-11 11-13 13-14-15 15-17 17-18 18-20 201-223 224-226 225-227 225-227 225-227 225-227 225-237 235-33 335-335-	10022696854351175201013000000000000000000000000000000000	2 609759286383635568248114400000000000000000000000000000000	8023617799851749660858402000000000000000000000000000000000	10049638455290667564351010000000000000000000000000000000000	1355835422590000000000000000000000000000000000	197 13603111145 11214 11214 1	, 15498861185554402724000000000000000000000000000000	149524037879668341000000000000000000000000000000000000	1 4 9 2 2 1 8 6 6 6 4 2 1 9 5 2 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 0 0 8 5 8 2 0 4 4 1 9 8 5 5 2 0 3 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 46654947891116253914100000000000000000000000000000000000	2 93690162239487433350300000000000000000000000000000000	1 1 8 9 6 8 3 1 1 8 9 6 7 6 9 3 5 5 5 3 2 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	14 832381633786966411141000000000000000000000000000000	1 4129598690052220333736119200000000000000000000000000000000000	16 4174267327005523001858874100000000000000000000000000000000000	1578 1783 1783 17914 10647 7754 10647 1064
TOTAL	92	989	980	886	1012	977	954	811	537	304	255	219	244	337	557	837	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 13

#### ATCHAFALAYA BAY

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEAS0	N ( M	AR TO	YAM C	)							113	44 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DI	RECTIO	N BAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-12 3-45 6-789 10-112 13-145 10-112 112-13 114-16 116-17 119-22 23-245 247-228 247-2289 257-2289 257-2289 257-2289 257-2289 257-2289 257-2289 257-2289 257-338 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-388 357-38	20010607657012310000000000000000000000000000000000	3487881476050348173855110000000000000000000000000000000000	5165624915251888861111663300000000000000000000000000	15481429961140221804761895000000000000000000000000000000000000	1699913121149947100000000000000000000000000000000	267055738321755727747614000000000000000000000000000000000	2130213149793311007774883379500000000000000000000000000000000000	100 106 711 166 1169 169 169 169 169 169 169 169	18 724 463 79 1118 5163 120 00 120 00 00 00 00 00 00 00	1880919976624052000000000000000000000000000000000	8888124 1291113225111022200000000000000000000000000	682028163842000000000000000000000000000000000000	72811459727240000000000000000000000000000000000	36470838453336342000000000000000000000000000000000	5469432189915409560040000000000000000000000000000000	1239463772114011489277300000000000000000000000000000000000	158 1633 1633 1633 1627 11621 1937 1649 1933 1933 1933 1933 1933 1933 1933 19
IUIAL	61	840	833	1091	1062	1408	1554	1298	590	281	171	159	111	111	210	223	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 14

#### ATCHAFALAYA BAY

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEASO	N ( J	UN TO	AUG	)							38	47 20	MIN	OBSER	VATIO	N PAIRS
WIND SPEED							DTR	FCTIO	N BAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-12 2-34 5-67 8-99 10-11 11-134 14-15 115-117 115-117 115-117 115-117 115-21 22-224 24-25 225-227 225-227 225-226 227-28 289-31 312-335 334-356 339-40 TOTAL	000000550000000000000000000000000000000	25055144886553002000500000000000000000000000000000	2501618830033202250220002000000000000000000000	11135688797280522222000000000000000000000000000000	1361465173335365555000002000000000000000000000000	11557885654 64	128826566665333885000000000000000000000000000	7545211994237200000000000000000000000000000000000	23 44 470 122 1688 1813 102 22 00 00 00 00 00 00 00 00 00 00 00 0	2227888202174468152200000000000000000000000000000000000	2123657441885522005000200000000000000000000000000	253327140564072255000000000000000000000000000000000	31318857622100000000000000000000000000000000000	52677789448875220005000500000000000000000000000000	723118955880002220000000000000000000000000000	222341018852200000000000000000000000000000000	207 257 5139 11361 11615 1166975 6975 1875 1875 1875 1875 1875 1875 1875 18
IOIAL	10	232	330	021	020	072	031	1002	1220	707	/30	071	031	465	280	236	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 15

#### ATCHAFALAYA BAY

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SE	ASON	( SEP	1 OT	( VOF								6371	20 M	IN OB	SERVA	TION	PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DIR 7	ECTIO 8	N BAN	D 10	11	12	13	14	15	16	TOTAL
WIND SPEED					5 907333478496501997341000000000000000000000000000000000000	6 001339574999 10339574999 1089039484410000000000000000000000000000000000					1			OB 1 4 4512293672555544991444630000000000000000000000000000000000			PAIRS  TOTAL  160 219 3869 687 781 864 1054 888 8456 610 522 5192 328 1558 84 87 41 1 1 30 1 00 00 00
36-37 37-38 38-39 39-40	0 0 0	0	0	0 0	0 0 0	0	0	0	0 0	0	0 0 0	0	0	0	0	0	0 0 0
TOTAL	42	916	·	1180	•		1015	918	554	265	189	238	307	365	524	579	ŭ

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 16

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASE	ON C	DEC T	O FEB	)							35	48 20	MIN	OBSE	RVATION	Y PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DIR 7	ECTION 8	I BANI 9	D 10	11	12	13	14	15	16	TOTAL
0-12-34-56-78-99-1112-12-14-56-78-99-1112-12-14-156-118-156-18-156-18-18-190-12-2234-256-227-28-289-3312-333-335-336-338-339-339-339-339-339-339-339-339-339	200055860152822118502028000000000000000000000000	140 140 140 844 1086 1457 1487 1487 1487 1487 1487 1487 1487 148	1425 8127561122018845711225000000000000000000000000000000000	16682396669209000000000000000000000000000000	14 28 116 165 226 198 339 251 15 51 80 00 00 00 00 00 00 00 00 00 00 00 00	1 1 5 2 1 1 1 1 2 1 1 1 2 5 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	8052822802212500520000000000000000000000	5254252020215000000000000000000000000000	520552522202825200000000000000000000000	58542220501022082005200000000000000000000000	080202802200500000520000000000000000000	2285151202521502000080000000000000000000	0582112228185511005252000000000000000000	221116653322254068642268992520000000000000000000000000000000000	112253137872 253137872 8411122 78411122367339223144222 5000000000000000000000000000000000	2 2 5 2 39 73 978 171 167 101 132 98 39 36 45 616 0 0 0 0 0 0 0	1 67 135 135 135 125 100 100 100 100 100 100 100 100 100 10
		-,,,,	_ , J_	1220	212	,0	, ,	02	"	04	72	70	100	4/7	1130	1057	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 17

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEASO	) NC	1AR TO	YAM C	)							20	85 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIRE	ECTION	I RANI	n							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-1 1-2 3-4 4-5 5-6 7-7 7-8 9-10 10-11 11-12-13 13-14 14-15-16 16-17 17-18-19-20 20-21-223-24 224-225-27 227-223 23-245-26 26-27 27-28-30 30-31 31-33-34 34-36-37 37-38-39-40	000009999049000000000000000000000000000	04934557501198042748859211668 110222448859211440000000000000000000000000000000000	040714999113311632183149000000000000000000000000000000000000	000003028746638268883333732800000000000000000000000000000	0004921751158333444940000000000000000000000000000000	00994489893788844000000000000000000000000000000000	000400499440000000000000000000000000000	000040004400000004000000000000000000000	000040400040000044000000000000000000000	000400004940049000400000000000000000000	000004444000090000000000000000000000000	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	000409009400000444400000000000000000000	0000992234494040404000000000000000000000	132623354 123 5752 123 123 123 123 123 123 123 123 123 12	000148899987789228932788440000000000000000000000000000000000	0 14835 14703 14835 14703 14835 14906 14835 14906 14835 14906 14835 14906 14835 14906 14835 14906 14835 14906 14835 14906 14835 14906 1490
TOTAL	71	2906	2306	1894	666	311	33	19	23	43	43	71	43	206	671	685	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 18

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

	SUMMER	SEASOI	N ( J	IUN TO	AUG	)							2	73 20	MIN	OBSER	NOITAV	PAIRS
W]	(ND SPEED (KNOTS)	1	2	3	4	5	6	DIRE	NOITS 8	BAND 9	10	11	12	13	14	15	16	TOTAL
	0-1-2-3-4-5-6-7-8-9-9-11-12-12-3-4-5-6-7-8-9-9-11-12-12-12-12-12-12-12-12-12-12-12-12-		0 0 0 0 73 1466 1466 109 109 109 0 0 0 0 0 0 0 0 0 0 0 0 0 0	009366336900000000000000000000000000000	003399666699360000000000000000000000000	733633633600000000000000000000000000000	0 7 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	000000000000000000000000000000000000000	000000000000000000000000000000000000000	060000000000000000000000000000000000000		00000000000000000000000000000000000000	0000038639600000000000000000000000000000	00000000000000000000000000000000000000	0 0 0 0 0 2199 183 366 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 36 73 1469 36 73 73 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 3 3 3 6 0 0 0 0 0 0 0 0 0 0 0 0 0 0	73 183 402 769 1062 1355 1648 915 1135 842 695 439 476 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
	TOTAL	U	726	203/	247U	007	13	U	U	v	U	13	ロング	1073	/07	200	73	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 19

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SE	ASON	( SEP	4 OT	( VOI								2149	20 M	IN OB	SERVA	TION	PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DIRI 7	101T3	BANI	10	11	12	13	14	15	16	TOTAL
0-1 1-22-34-56-77-8-9-10 10-11 11-12 12-13-14-15-16-17 115-16-17 115-17-18-19 19-221-223-24-25-224-25-224-25-224-25-225-231 23-331-334-335-336-336-335-336-336-338-39-40	000000030009404000000000000000000000000	4083399672445754727487000000000000000000000000000000000	1837493929844985713532211622116221162221162211622211622211622116222211622211622211622211622211622211622211622211622211622211622211622211622211622211622211621162	93125919483252667622400000000000000000000000000000000	01335546 18779997371026 11062313344 000000000000000000000000000000000	444444394000004300000000000000000000000	900000040009000000000000000000000000000	089004009900000000000000000000000000000	409440044000000000000000000000000000000	434400004000004400000000000000000000000	494440000000000000000000000000000000000	48379233000040000000000000000000000000000000	03253397344000440000000000000000000000000000	992613370133890000000000000000000000000000000000	43789216354 2327222007333830000000000000000000000000	40909231663616951890647030000000000000000000000000000000000	83 181 279 432 581 5395 670 558 649 795 333 1158 1443 603 000 000 000 000 000 000
TOTAL	32	1740	2382	2145	926	74	23	51	32	41	32	237	228	376	670	1005	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 20

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASO	) N ( D	EC TO	FEB	)							25	68 20	MIN	OBSER	RVATION	PAIRS
WIND SPEED							DIR	ECTIO	N BAN								
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-12 2-34-5 1-6-7-8 10-11 11-12 113-14 114-15 115-17 117-18 18-19 120-221 221-224 225-227 228-230 331-333 335-336 337-338 337-338 337-338 337-338	030 077 2738 1151 1533 1951 15333 0000 0000 0000 0000	70 70 70 50 120 120 120 120 120 120 120 120 120 12	110352669460587770693300000000000000000000000000000000	1107963455999779553112111111111111111111111111111111111	113944311755231175575119350030333000000000000000000000000000	3111533240511795811150300000000000000000000000000000000	0 115 232 157 277 1151 3254 150 000 000 000 000 000 000 000 000 000	193111783755593813772119300000000000000000000000000000000000	1933 1913 1313 1313 1313 1313 1313 1313	1917 3155288823111557050000000000000000000000000000000	11 11 11 11 11 11 11 11 11 11 11 11 11	2 0 3 5 9 1 1 3 1 1 2 2 2 2 1 1 5 5 7 3 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	112359257852212757913733000000000000000000000000000000000	7 7 7 7 31 11 19 23 27 27 46 60 31 15 50 8 23 7 7 7 27 7 27 19 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3315740760111020044835553303000000000000000000000000000	0 31 115 356 881 120 93 120 101 102 105 110 106 107 100 100 100 100 100 100 100 100 100	1635 10218 52385 77635 77319 5546669 6612 66612 1187 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 21

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEASO	) N ( M	1AR TO	YAM (	)							12	37 20	MIN	OBSER	VATION	PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DIR 7	ECTIO 8	N BAN	D 10	11	12	13	14	15	16	TOTAL
	1 880800044864688066680080000000000000000	2 1662431 12108979116488711164400 11211720400 11211720400 11211720 11211720 1121720 11211720 11211720 11211720 11211720 11211720 11211720 1121172	3	4 860842500000000000000000000000000000000000	5 04225 30587 1 8880080000000000000000000000000000000	6 888082808080800000000000 <b>f;@00</b> 0000					11 816 162 132 162 224 880 00 00 00 00 00 00 00 00 00 00 00	1	13 404200200800080000000000000000000000000	14 16 16 16 16 16 16 16 16 16 16 16 16 16	1 2 486087666402200000000000000000000000000000000	16 88 16 12 12 13 12 16 16 16 16 16 16 16 16 16 16 16 16 16	TOTAL 2910 2751377637763779387799875545799624032278885424600000000000000000000000000000000000
TOTAL	388	1697	1341	1018	913	509	404	396	323	242	242	242	379	476	646	776	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 22

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEAS0	и ( Ј	UN TO	AUG	)							6	104 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIR	ECTIO	N BAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-1	0	24	0	24	49	24	49	24	74	74	49	74	74	49	24	0	618
1- 2	0	24	24	49	49	0	24	0	74	24	24	24	74	0	Ó	24	420
2- 3	0	49	74	49	49	74	74	148	0	24	24	0	74	74	Ō	24	742
3- 4	0	0	99	123	74	0	99	99	24	49	99	24	24	24	123	Ö	866
4-5	0	74	74	49	173	99	49	49	74	24	Ö	24	99	74	24	123	1014
5- 6	0	99	99	49	74	74	49	24	49	0	198	49	49	Ò	49	Ō	866
6- 7	0	99	74	74	24	123	74	24	49	49	74	49	123	ŏ	99	24	965
7- 8	24	123	99	24	24	24	173	123	49	24	49	24	24	99	24	148	1064
8- 9	Ō	74	99	24	24	24	49	74	Ö	49	49	74	49	123	24	74	816
9-10	Ō	24	24	49	0	Ō	49	24	24	74	99	148	123	74	24	Ò	742
10-11	Ō	- 0	Ō	Ö	Ō	49	49	49	49	49	24	49	49	74	49	49	544
11-12	Ō	148	49	Ō	49	24	49	Ŏ	49	198	24	24	Ó	Ò	'n	Ó	618
12-13	Ō	0	0	49	0	24	0	Ō	Ö	49	24	Ō	Ŏ	ŏ	ŏ	ŏ	148
13-14	Ō	Ō	Ō	24	24	Ō	Ō	Ō	24	Ö	24	ŏ	Ŏ	24	ŏ	Ŏ	123
14-15	Ō	24	Ŏ	24	Ō	Ŏ	24	Ŏ	Ō	Ŏ	24	Ŏ	Ō	24	ŏ	ŏ	123
15-16	0	0	0	0	0	0	Ō	Ŏ	24	Õ	Ö	Ŏ	ō	Ō	ñ	ŏ	24
16-17	Ō	Ō	0	Ō	Ō	24	Ō	Ō	Ö	24	ō	ŏ	Ŏ	ŏ	24	ŏ	74
17-18	0	0	0	24	0	0	Ó	Ó	24	24	Ŏ	Õ	Ō	Ō	- i	ň	74
18-19	Ō	49	Ō	0	Ó	Ō	Ō	Ō	24	24	Ŏ	Ŏ	Ŏ	ŏ	ň	ň	99
19-20	0	0	0	Ō	0	0	0	0	0	Ō	Ō	Ō	Ō	Ō	ŏ	ñ	á
20-21	Ō	0	24	0	24	Ō	Ō	Ō	Ō	Ō	ā	Õ	Ŏ	ŏ	ň	ŏ	49
21-22	Ō	Ō	0	Ō	0	Ō	Ō	Ŏ	Ō	Ō	Ŏ	Ŏ	ŏ	ă	Õ	ň	'ó
22-23	Ó	Ó	G	Ö	Ō	Ō	Ō	Ō	Ō	Ō	Ŏ	Ŏ	Ŏ	ŏ	ŏ	Ŏ	Ğ
23-24	Ō	Ö	Ō	Ŏ	Ō	Ŏ	Ō	Ō	Ŏ	Ō	Ŏ	Ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ
24-25	Ō	Ō	Ö	Ō	Ō	Ö	Ō	Ō	Ō	Ō	ŏ	Ŏ	Ŏ	ŏ	ă	ŏ	ŏ
25-26	0	0	Ö	0	Ō	0	Ō	Ō	Ō	O.	Ō	Ō	Ō	Ö	ă	ň	Ŏ
26-27	Ö	Ô	Ö	Ö	Ö	Ö	Ó	Ō	Ŏ	Ö	Ŏ	Ŏ	Ö	õ	ŏ	ŏ	ň
27-28	0	0	0	0	0	0	Ó	Ō	Ŏ	Ō	Ŏ	Ō	Ō	Ŏ	ō	ŏ	Ŏ
28-29	Ö	0	0	0	0	0	Ó	Ö	Ŏ	Ō	Ŏ	Ŏ	Ŏ	Ŏ	ă	Ŏ	Ŏ
29-30	0	0	0	0	0	0	0	0	Ō	0	Ó	Ó	Ō	Ō	ă	Ŏ	Ŏ
30-31	0	0	0	0	0	0	0	0	0	0	Ó	Ó	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ
31-32	0	0	0	0	0	0	0	0	0	0	0	0	Ó	Ó	Ō	Ō	Ō
32-33	0	0	0	0	0	0	0	0	0	0	0	0	Ò	Ó	Ō	Õ	Ŏ
33-34	0	0	0	0	0	0	0	0	0	0	0	0	0	Ó	Ó	Ō	Ö
34-35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ō	Ō	Ŏ
35-36	0	0	0	0	0	0	0	0	0	0	0	Ó	0	Ó	Ó	Ō	Ö
36-37	0	0	0	0	0	0	0	0	0	0	0	0	0	Ó	0	0	Ö
37-38	0	0	0	0	0	0	0	0	0	0	Ō	Ö	Ŏ	Ŏ	ŏ	Õ	Ö
38-39	0	0	0	0	0	0	0	0	0	0	0	Ō	Ō	Ó	Ō	Ö	Ö
39-40	0	0	0	0	0	C	0	0	0	0	0	Ō	Ō	Ō	Ö	Ŏ	Ö
TOTAL	24	816	742	643	643	569	816	643	618	767	792	569	767	643	470	470	
												/					

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 23

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SE	ASON	( SEF	TO N	( VO								1273	20 M	IN OB	SERVA	TION	PAIRS
WIND SPEED							חזת	ECTIO	N BAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
WIND SPEED (KNOTS)  0-1 1-2 3-4 4-5 6-7 7-8 8-9 10-11 11-12 11-13 13-14 14-15-16 16-17 17-18 18-19 120-21 21-23 23-24 24-25 25-27 27-28 28-30	1 0707103505070007000000000000000000000000	2 702235829 105755758829 12035829 120300000000000000000000000000000000000	3 157221779711497 1157821397777000000700000700000700000700000070000	4 933450770422177000000770000007000	5 336477405595573010570000000000000000000000000000000	6 3477 4469 3177 3212 0570 000 000 000 000		EC 710 215718 357718 3577155007775000000000000000000000000000			11 623357 622157 377299357 2700000000000000000000000000000000000	1 2 143173553377777557007000000000000000000000	13 350268383758777570070000000000000000000000000	1 253763711212 253763711212	15 3907853387127151277050000000000000000000000000000	16 157 317 628 787 777 777 200 000 000	TOTAL  361 369 864 974 1107 989 1068 824 597 416 259 337 369 416 337 298 86 62 54 70 54 70 23 31
30-31 31-32 32-33 33-34 34-35 35-36 36-37 37-38 38-39	0 0 0 0 0 0	00000000	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	000000000000000000000000000000000000000	000000000000000000000000000000000000000	000000000000000000000000000000000000000	0 0 0 0 0 0 0 0	0 0 0 0 0 0 0	0 0 0 0 0 0 0 0	000000000000000000000000000000000000000	0 0 0 0 0 0	000000000000000000000000000000000000000	000000000000000000000000000000000000000	0 0 0 0 0 0 0	0 0 0 0 0 0 0	0 0 0 0 0 0	0 0 0 0 0 0
39-40 Total	164	1893	1146	809	549	0 518	322	329	0 455	306	0 424	361	463	0 487	934	0 832	0

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 24

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WIND SPEED	WINTER	SEASON	( DE	EC TO	FEB	)							172	1 20	MIN	OBSER	/ATION	PAIRS
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		1	2	3	4	5	6				10	11	12	13	14	15	16	TOTAL
TOTAL 0 69 302 2289 3759 2155 976 337 104 5 0 0 0 0 0	1- 2 2- 3 3- 4 4- 5 5- 6 6- 7 7- 8 8- 9 9-11 11-12 12-13 13-14 14-15 15-16 16-17 17-18 18-19 19-20 20-21 22-23 23-24 22-25 22-25 22-26 26-27 27-28 28-29 29-30 30-31 31-32 32-33 33-35 36-37 37-38 38-39 39-40	000000000000000000000000000000000000000	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	00072994350011511155430000000000000000000000000	500583225769891711191000000000000000000000000000000	5740099446691766691183994446331000000000000000000000000000000000	0116 12213 12734 12733 1263 12734 1273 1263 1273 1263 1273 1273 1273 1273 1273 1273 1273 127	0 114 662 1456 1456 1456 1569 1117 000 000 000 000 000 000 000 000 00	00 240 240 255 11 29 00 00 00 00 00 00 00 00 00 00 00 00 00	000601500000000000000000000000000000000	000050000000000000000000000000000000000			000000000000000000000000000000000000000				11 40 180 476 813 912 1417 1464 1301 987 604 395 290 220 319 209 139 52 87 58

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 25

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEASO	N ( M	AR T	YAM C	)							25	82 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							אומ	ECTIO	N RAN	n							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
(KNOTS)  0-1 2-34 5-67 8-99 11-12 12-13 13-14 15-16 16-18 18-19 19-21 22-22 23-22 23-22 25-26 26-28 29-31 31-32 33-34 35-36 36-38 39-40	1 7 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2 3770671330733030000000000000000000000000	3 90251142954673030000000000000000000000000000000000	4 11506 19705 1705 1705 1705 1705 1705 1705 1705 1	5 119918183355400013411100000000000000000000000000	6 033033411318887340703030000000000000000000000000000					11 27 19 30 00 00 00 00 00 00 00 00 00 00 00 00	12 755730000000000000000000000000000000000	13 11 75 73 11 30 00 00 00 00 00 00 00 00 00 00 00 00	14 371377000000000000000000000000000000000	15 11 33 00 00 00 00 00 00 00 00 00 00 00 00	16 3000030000000030000000000000000000000	TOTAL  247 23797 6228 8901 10848 6449 10848 2130 0300 0000 0000 0000 0000
TOTAL	19	251	770	1994	2072	2017	1243	546	441	309	108	69	81	42	19	11	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 26

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEASON	( J	UN TO	AUG	)							5	27 20	MIN	OBSER	VATION	PAIRS
WIND SPEED	1	2	3	4	5	6	DIR 7	ECTIO			11	12	13	14	15	16	TOTAL
NOTS)  0-1 2-3 3-4 5-6 6-7 8-9 9-10 10-11 112-13 13-15-16 16-17 17-18 19-20 20-21 222-23 23-25 26-28 29-30 30-31 32-32 33-33 335-38 36-38 36-38 39-40		2 0000000000000000000000000000000000000	3 0 0 0 0 0 188 37 5 6 75 75 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4 000049095433152168880000000000000000000000000000000000	5 053795099117329911732911533 11533 0000000000000000000000000000000000	6 6 671189514885120895120895120895120895120800000000000000000000000000000000000	7 0803186241987800000000000000000000000000000000000	8 0 0 0 0 18 75 189 754 1327 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	9 8007455800000000000000000000000000000000	10 00 188 180 00 00 00 00 00 00 00 00 00 00 00 00 0		12 00 18 00 00 00 00 00 00 00 00 00 00 00 00 00	13 0 0 0 0 18 0 0 0 0 0 0 0 0 0 0 0 0 0 0	14 0 0 18 18 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	15 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	16 00 00 18 00 00 00 00 00 00 00 00 00 00 00 00 00	TOTAL  75 132 303 7593 1726 11714 17596 3322 758 37 00 00 00 00 00 00 00 00 00 00 00 00 00
TOTAL	0	0	284	2220	2789	2087	1404	664	322	75	0	18	37	37	18	37	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 27

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WIND SPEED  (KNOTS)  1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 TOTAL  0 1 0 0 0 6 13 6 13 20 13 13 0 13 6 6 0 0 0 137  1 2 3 0 0 6 0 6 13 34 41 27 0 0 0 0 0 0 0 0 0 0 137  2 3 0 0 6 6 27 34 27 62 27 20 0 0 0 0 0 0 0 0 0 0 0 207  3 4 0 0 0 6 34 62 62 76 69 20 13 0 0 0 0 0 0 0 0 346  4 5 0 0 13 110 221 193 124 103 27 6 0 0 0 0 0 0 0 0 346  5 6 0 0 0 6 90 221 283 159 124 34 0 0 0 0 0 0 0 0 0 0 346  6 7 0 0 6 6 145 346 290 186 117 13 0 0 0 0 0 0 0 0 0 0 0 1188  8 9 0 0 13 263 491 353 271 66 110 6 0 0 0 0 0 0 0 0 0 1188  8 9 0 0 13 263 491 353 271 69 0 0 0 0 0 0 0 0 0 0 1388  8 9 0 0 16 6 20 373 498 367 76 0 0 0 0 0 0 0 0 0 0 0 1336  10-11 0 0 0 166 290 221 62 0 0 6 0 0 0 0 0 0 0 0 0 1336  10-11 0 0 0 166 290 221 62 0 0 6 0 0 0 0 0 0 0 0 0 0 1336  10-11 0 0 0 167 244 81 31 33 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	FALL SE	ASON (	SEP	TO 1	( VOF								1444	20 M	IN OBS	SERVA	TION P	AIRS
CKHÖTS    1   2   3   4   5   6   7   8   9   10   11   12   13   14   15   16   TOTAL	WIND SPEED							DIR	ECTIO	N RAN	n							
1-2 0 0 6 6 0 6 13 34 47 1 27 0 0 0 0 0 0 0 0 0 1 131		1	2	3	4	5	6					11	12	13	14	15	16	TOTAL
1-2 0 0 6 6 0 6 13 34 41 27 0 0 0 0 0 0 0 0 1 131	0- 1	n	n	n	6	13	6	13	20	13	13	Λ	13	4	6	n	0	117
2-3																		
\$\frac{4-5}{5-6}\$ 0 0 13 110 221 193 124 103 27 66 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2- 3	0	0	6	27	34	27	62	27	20	0	0	Ó	Ó	Ō	Ŏ	Ö	
5-6 0 0 6 790 221 283 159 124 34 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		0		6							13	0	0	0	0	0	0	346
6-7 0 0 0 6 145 346 290 186 117 13 0 0 0 0 0 0 0 0 1 108 8-9 8-9 0 0 13 180 533 373 166 110 6 0 0 0 0 0 0 0 0 0 0 1385 8-9 9 0 0 13 263 491 353 221 69 0 0 0 0 0 0 0 0 0 0 0 1412 9-10 0 0 20 373 498 367 76 0 0 0 0 0 0 0 0 0 0 0 0 1336 10-11 0 0 0 166 290 221 62 0 0 6 0 0 0 0 0 0 0 0 0 0 1336 10-11 0 0 0 166 290 221 62 0 0 6 0 0 0 0 0 0 0 0 0 0 747 11-12 0 0 0 0 207 242 159 488 66 0 0 0 0 0 0 0 0 0 0 0 0 747 11-13 0 0 0 0 90 124 48 13 13 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		•	-									•	0	0	0	0	0	803
7-8 0 0 13 180 533 373 166 110 6 0 0 0 0 0 0 0 0 1385 8-9 0 0 13 263 491 553 221 69 0 0 0 0 0 0 0 0 0 0 1412 9-10 0 0 0 20 373 498 367 76 69 0 0 0 0 0 0 0 0 0 0 0 1412 9-10 0 0 0 16 290 221 62 0 0 6 0 0 0 0 0 0 0 0 0 0 777 11-12 0 0 0 0 207 242 159 48 6 6 6 0 0 0 0 0 0 0 0 0 0 0 777 11-12 0 0 0 0 124 48 13 13 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		-	-	-							-		•		0	-	•	
8-9 0 0 13 263 491 355 221 69 0 0 0 0 0 0 0 0 0 1412 9-10 0 0 20 373 498 367 766 0 0 0 0 0 0 0 0 0 0 0 1336 10-11 0 0 0 166 290 221 62 0 0 6 0 0 0 0 0 0 0 0 777 11-12 0 0 0 0 207 242 159 48 6 6 0 0 0 0 0 0 0 0 0 0 0 777 11-12 0 0 0 0 90 124 48 13 13 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		-		-							-	-	•	•	•		•	
9-10		•									•	-	•	-	-	•	-	
10-11		•								-	-	-	•	-	0	•		
11-12		•		-:					-		Ų	-	•		Ü			
12-13		•		-							Ď	•	•		Ů			
13-14         0         0         0         117         131         62         0         6         0		•		-							•	-	•	-	V	-	-	
14-15		•		-						-	-	-	-	-	Ů	•	•	
15-16		n		-									•	-	ň	-	-	
16-17		ñ		-			-	-		-	_	•	•	-	ň	-	-	
17-18       0 <td></td> <td>ŏ</td> <td></td> <td></td> <td></td> <td></td> <td>-</td> <td>-</td> <td>•</td> <td></td> <td>•</td> <td>•</td> <td>•</td> <td>-</td> <td>ň</td> <td>•</td> <td>•</td> <td></td>		ŏ					-	-	•		•	•	•	-	ň	•	•	
18-19		Ŏ		-	-		-	-		-	_	-	-	-	ŏ	•	-	_
20-21		Ö	Ó	Ŏ	Ó	Ô	Ō	Ŏ	Ŏ	Ō	Ŏ	ŏ	ŏ		ŏ	ă		-
21-22	19-20	0	0	Ó	Ó	0	Ô	Ó	Ō	Ö	Ö	Ŏ	Ŏ	Ŏ	Ŏ		-	ă
22-23		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ó	Ŏ
23-24		•	•	•		0				•			•	0	0	0	0	0
24-25		•	-	•		0	•		•		-		-	-	0		-	0
25-26		-	-	•	_	0	•				•		•		Ō		-	0
26-27				•		0	•				-	•	-	-	0	-	-	Ō
27-28		•		_	_	U	-	-	•		-	•	•	•	0	-	-	0
28-29 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		•		•	-	U			•	-	•	•	•	•	0		-	0
29-30		•		•		Ü		•	•		•		•	-	•	-		Ų
30-31 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		•	•	•	•	•	•	-		-	•		•	-	•			V
31-32 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0				-		•			•	-	•		•	-	ň	-		Ň
32-33		ě	-	ŏ	-	-				-	•			-	ñ			ň
33-34	32-33	Ö	Ö	Ō	Ö	Ō	Ō				-	•	-	•	ŏ	•		ŏ
35-36		0	0	0	0	0	0	0	Ó	Ô	Ò	Ó	Ö	Ō	Ŏ	Ŏ	Ŏ	Ŏ
36-37 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		-		•		-		0	0	0	0	0	0	0	0	Ó	Ó	Ō
37-38 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		-	•	-	-	0			•	-	0		0	0	0	0	0	Ö
38-39 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0			-	•		0	-	-	•	-	•	-	-	0	0	0	0	0
39-40 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0			-	-	-	•			•	-	•		-	-	•	•	•	0
									-									0
TOTAL 0 0 96 1855 3372 2465 1253 713 173 41 0 13 4 4 0 0	39-40	U	U	U	0	0	U	Ü	0	0	0	0	0	0	0	0	0	0
	TOTAL	n	n	96	1855	3372	2465	1253	713	173	6.1	•	17			•	•	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 28

**GULF RETURN** 

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASON	( DE	с то	FEB	)							27	02 20	MIN	OBSER	VATION	PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DII 7	RECTI	DN BAN	ID 10	11	12	13	14	15	16	TOTAL
(KNOTS)  0- 1 1- 2 2- 3 3- 4 4- 5 6- 7 7- 8 8- 9 9-11 11-12 12-14 11-15 15-16 16-17 17-18 18-19 19-21 21-22 22-24 24-25 25-26 26-27 28-29 29-30 31-32 32-33 34-35 35-37 37-38 38-39 39-40	1 0000000000000000000000000000000000000	2 3000000000000000000000000000000000000	3 3000000000700000000000000000000000000	4 71807345227730000000000000000000000000000000000	5 4 0 3 0 9 9 2 2 3 3 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	6 07211777467369224770000000000000000000000000000000000	7 7 7 0 8 7 7 0 8 1 7 1 2 1 8 1 8 1 8 1 2 1 2 1 1 2 1 8 3 1 2 1 1 2 1 1 1 8 3 1 2 1 1 1 1 8 3 1 2 1 1 1 1 8 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1	8 43475545354479437000000000000000000000000000000000	9 13141666783367557000000000000000000000000000000000	10	11 7 0 0 0 7 1 1 1 5 7 0 8 2 2 2 2 8 0 0 0 0 0 0 0 0 0 0 0 0 0	12 110000025559230000000000000000000000000000	13 25 00 00 37 01 13 00 00 00 00 00 00 00 00 00 00 00 00 00	14 18 0 0 3 3 3 7 7 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	15 3033300330000000000000000000000000000	16 3003333000000000000000000000000000000	10 TAL 210 225 125 125 262 1384 1728 8647 381 128 8647 385 00 00 00 00 00 00 00 00 00 00 00 00 00
TOTAL	0	3	11	151	851	1835	2472	2235	1458	392	259	196	51	44	18	18	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 29

# GULF RETURN BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEASON	( MA	R TO	MAY	)							47	53 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIE	RECTIO	N BAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0- 1	0	0	0	6	21	21	31	6	14	8	2	6	2	2	0	0	122
1- 2	Ō	0	2	10	21	23	25	6	8	6	2	10	2	6	ž	ž	128
2- 3	0	0	2	18	35	37	56	27	18	21	2	12	4	0	0	2	239
3- 4	0	16	14	27	42	75	107	94	54	25	12	8	2	4	0	2	488
4~ 5	Ō	4	25	63	113	172	189	189	73	31	31	18	4	0	0	0	917
5- 6	0	0	4	71	105	220	262	340	132	37	31	18	_ 6	0	Ō	Ō	1232
6- 7	0	0	10	23	79	237	300	448	174	46	54	18	10	6	0	0	1411
7- 8	0	0	0	4	56	212	296	481	225	52	35	23	. 6	2	0	0	1397
8- 9 9-10	0	0	0	0	33 35	183 183	321 195	342 214	107 86	42 35	18	21	16	0	0	0	1087
18-11	n	ů	0	0	42	84	151	151	29	35 8	6 4	16 10	8 2	0	0	0	782
11-12	ň	Ö	ŏ	Ď	44	96	132	98	18	2	8	10	6	0	0	0	483 408
12-13	ň	ň	ŏ	ŏ	6	73	130	58	18	6	2	ŏ	Ö	Ö	0	Ŏ	296
13-14	ŏ	ŏ	ŏ	ŏ	6	56	111	31	-4	ŏ	ō	ŏ	ŏ	ŏ	ŏ	ŏ	210
14-15	Ŏ	Ö	Ŏ	ŏ	2	44	105	25	Ó	ž	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	178
15-16	Ŏ	Ō	Ŏ	Ō	0	25	73	25	Ŏ	ž	Õ	ŏ	Ŏ	ŏ	ŏ	ŏ	126
16-17	0	0	0	0	0	23	69	10	0	2	Ó	Ô	Ó	Ō	Ŏ	Ŏ	105
17-18	0	0	0	0	0	14	50	2	2	0	0	0	0	0	0	0	69
18-19	0	0	0	0	Ō	8	33	2	0	0	0	0	0	0	0	0	44
19-20	0	0	0	0	0	14	48	6	0	0	0	0	Ō	0	Ō	0	69
20-21	0	0	0	0	0	14	54	0	0	0	0	0	0	0	0	0	69
21-22	0	0	Ü	0	0	. 2	63	8	4	0	0	0	0	0	0	0	77
22-23 23-24	0	0	U	0	<b>0</b> 0	10	23 12	4 0	0	0	0	0	0	0	0	0	37
24-25	0 n	0	0	0	ů ů	2	12	0	0 0	Ů	0	0	0	0	0	0	14
25-26	0	Ö	Û	Ö	Ü	ů	0	Ö	Ö	ů	0	ű	0	0	0	0	0
26-27	ň	ő	ň	Ö	ň	ñ	ŏ	ŏ	ŏ	ñ	Ö	Ö	Ö	ň	Ö	Ö	Ů
27-28	Ď	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	Ö	ň
28-29	ŏ	ŏ	ŏ	Ŏ	Ŏ	ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ň	ň
29-30	0	0	0	0	0	0	0	0	0	0	Ö	Ó	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ
30-31	0	0	0	0	0	0	0	0	0	0	0	0	0	Ö	Ó	0	Ō
31-32	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32-33	Ō	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
33-34	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ō	0
34-35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35-36 36-37	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
36-37 37-38	0	0	0	0	0	0	0	0	U D	U 0	0	0	0	0	0	0	0
37-38 38-39	0	Ö	0	0	0	0	Ö	0	0	0	0	0	0	0	0	U	U
39-40	ő	Ö	ñ	Ö	Ď	Ů	Ô	Ŏ	Ö	ů	0	0	ů	0	0	0	0
3, 10	·	v	v	J	J	J	J	J	v	U	U	v	U	U	U	U	U
TOTAL	0	21	58	225	645	1838	2848	2577	974	330	212	166	71	21	2	6	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 30

#### **GULF RETURN**

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEASON	( JUI	OT P	AUG	)							15	20 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIE	PECTI	ON BAI	ND.							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0- 1	0	0	0	0	0	0	6	0	13	6	13	6	32	0	0	0	78
1- 2	Ö	Ö	ŏ	6	6	ŏ	13	52	46	19	13	13	6	19	Ö	Ğ	184
2- 3	ň	Ŏ	ŏ	19	19	39	59	59	59	26	19	13	6	Ťó	6	Ŏ	328
3- 4	Õ	Ď	ŏ	Ď	39	5 <b>9</b>	85	151	105	59	26	46	32	6	6	ŏ	618
<del>4</del> – 5	ŏ	Ŏ	Ŏ	Õ	32	85	125	276	210	72	59	19	6	19	ŏ	ŏ	907
5- 6	Ŏ	Ò	Ö	13	26	92	78	368	348	138	164	39	13	- 6	6	Ŏ	1296
6- 7	Ŏ	Ö	Ŏ	-6	19	52	46	289	342	138	243	59	46	ĕ	6	Õ	1256
7-8	Ò	0	0	0	6	46	125	184	394	131	125	78	32	Ō	Ō	Õ	1125
8- 9	Ó	0	0	0	13	46	131	217	217	157	125	59	19	Ō	13	Ò	1000
9-10	0	0	0	0	0	59	65	131	250	151	72	26	6	Ó	Ō	Ŏ	763
10-11	0	0	6	6	13	59	65	111	144	92	78	52	6	6	0	Ô	644
11-12	0	0	0	0	6	13	157	157	144	144	19	6	6	0	0	0	657
12-13	0	0	0	0	6	0	52	85	85	98	26	6	13	0	0	0	375
13-14	0	0	0	0	0	6	85	19	78	118	6	6	0	0	0	0	322
14-15	0	0	0	0	0	_ 0	65	6	6	72	32	Ō	0	0	0	0	184
15-16	Ō	0	0	0	0	13	46	0	0	78	6	0	0	0	0	Ō	144
16-17	0	0	0	0	0	0	13	0	0	59	6	0	0	0	0	0	78
17-18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
18-19	0	0	0	6	0	0	0	0	0	0	. 0	0	0	6	0	Ō	13
19-20	0	0	0	0	0	0	0	0	0	0	13	0	9	0	0	0	13
20-21	0	0	0	0	0	0	0	0	Õ	0	0	0	0	0	0	Ų	0
21-22 22-23	0	0	0	0	0 <b>0</b>	0	0	0	0	0	0	0	0	0	0	Ŭ	0
23-24	0	0	0	0	0	ň	ŏ	ů	0	Ů	6	Ü	0	0	. U	V	Ų
24-25	ñ	ň	Ö	ñ	ő	ň	ň	ñ	Ö	ň	ñ	ů	ň	ŏ	ň	Ŭ	6
25-26	ň	ŏ	ŏ	ő	ő	ň	ŏ	ň	ŏ	0	ň	ŏ	ŏ	Ö	ñ	ň	0
26-27	Õ	ň	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ñ	ŏ	Ď	ŏ	ŏ	ň	ň	ň
27-28	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ñ	ŏ
28-29	Ŏ	Ō	Ŏ	Ŏ	Ō	Ŏ	Ŏ	Ó	Ŏ	Ŏ	Ŏ	Ö	ŏ	ŏ	Ŏ	ŏ	Ď
29-30	0	0	Ó	Ó	0	0	Ó	Ó	Ō	Ō	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ
30-31	0	0	0	0	0	0	0	0	0	0	0	0	0	Ó	Ó	Ó	Ó
31-32	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
32-33	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
33-34	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
34-35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35-36	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
36-37	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
37-38	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ō
38-39	0	0	0	0	Õ	0	0	0	0	0	0	0	0	0	0	0	0
39-40	0	υ	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL	0	0	6	59	190	572	1223	2111	2447	1565	1046	434	230	72	39	0	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 31

#### **GULF RETURN**

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SE	ASON (	SEP	TO NOV	<i>(</i> )								1135	20 M	IN OBS	SERVA"	TION P	AIRS
WIND SPEED							חזת	FCTT	N BAN	n							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0- 1	•	0	0	0	0	0	17	8	0	17	8	0	0	0	0	0	52
1- 2	0	0	ŭ	Ö	ŏ	ŏ	35	8	8	8	8	ŏ	ŏ	ŏ	Ŏ	Ů	70
2- 3	ŏ	ŏ	ŏ	ŏ	17	52	44	88	35	17	8	ŏ	ŏ	ŏ	ŏ	ŏ	264
3- 4	ŏ	ň	ŭ	ň	17	61	79	149	88	26	ŏ	ň	ŏ	ă	ŏ	ŏ	422
4- 5	ŏ	ŏ	ŏ	ŏ	Ö	52	114	132	158	26	ŏ	ŏ	ă	ŏ	ă	ŏ	484
5-6	ň	Õ	ŏ	ŏ	ŏ	88	229	378	273	61	17	Ŏ	Ŏ	ŏ	ŏ	ŏ	1048
6- 7	ň	ŏ	Ŏ	Ŏ	17	114	325	387	176	44	35	8	Ŏ	ŏ	Ŏ	Ŏ	1110
7- 8	ŏ	Ŏ	Ŏ	Ŏ	Ö	79	502	651	422	132	26	Ō	Ò	Ŏ	Ō	Ŏ	1814
8- 9	Ŏ	Ō	Ö	Ō	Ó	35	440	616	308	17	0	8	0	Ó	0	Ó	1427
9-10	Ó	0	0	0	0	52	414	378	193	26	26	0	0	0	0	0	1092
10-11	0	0	0	0	0	52	387	264	8	17	0	0	0	0	0	0	731
11-12	0	0	0	0	0	70	132	123	105	17	0	0	0	0	0	0	449
12-13	0	0	0	0	0	17	158	96	88	17	0	0	8	0	0	0	387
13-14	0	0	0	0	0	0	185	35	70	17	0	8	0	8	0	0	325
14-15	0	0	0	0	0	0	44	0	52	26	0	Ō	Ō	0	8	0	132
15-16	0	0	Ō	0	0	0	17	8	. 8	44	8	0	0	0	0	0	88
16-17	0	0	0	0	0	0	17	0	44	8	0	0	0	0	8	0	79
17-18	0	0	0	0	0	0	0	0	8	0	0	0	0	0	0	0	8
18-19	0	0	0	0	0	0	Ŏ	0	0	8	0	0	0	0	0	0	8
19-20	0	0	0	0	0	0	0	0	0	0	0 0	ŭ	0	0	0	V	Ų
20-21	0	Ü	0	0	0	0	0	Ü	0	0	Ü	ŭ	ű	0	0	V	Ų
21-22 22-23	V	Ö	ŏ	Ô	0	V	Ö	ŭ	ů	Ŏ	Û	ŏ	ñ	Ö	ŏ	ů	0
23-24	Ď	ň	ů	ŏ	ň	ñ	ů	ŏ	Ů	ŏ	ä	ŏ	ă	ŏ	ŭ	Ů	Ů
24-25	Ď	ŏ	ŏ	Ö	ŏ	ň	ŏ	ň	ŏ	ň	ŏ	ň	ă	ŏ	ñ	n	ň
25-26	ň	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ň	ŏ	ŏ	ŏ	ň	õ	ŏ	ň	ň	ň
26-27	ň	ŏ	ŏ	ŏ	Ď	ŏ	ŏ	ň	ŏ	Ď	ň	ŏ	ă	ŏ	ă	ň	ŏ
27-28	ŏ	ŏ	Ď	ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	Ŏ	Ŏ	ŏ	ŏ	ŏ	Ď	ŏ
28-29	Ď	Ŏ	Ŏ	Ŏ	Ō	Ŏ	č	Ŏ	ŏ	Ŏ	Ŏ	Ö	Ŏ	ŏ	Ŏ	Ŏ	ŏ
29-30	Ŏ	Ŏ	Ŏ	Ŏ	0	Ó	Ö	Ŏ	Ŏ	Ó	Ō	Ö	Ó	Õ	Ö	Ŏ	Ŏ
30-31	Ò	0	0	Ó	0	0	G	0	Ô	0	0	0	0	0	0	Ô	0
31-32	0	0	0	0	0	0	G	G	0	0	0	0	0	0	0	0	0
32-33	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
33-34	0	Q	O	0	0	0	0	0	0	0	0	0	0	0	0	0	0
34-35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35-36	0	Ō	0	0	Ō	0	0	0	0	0	0	0	0	G	0	0	0
36-37	0	0	0	0	0	0	0	0	0	0	0	0	Ō	0	Ō	0	Ō
37-38	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ō
38-39	0	0	0	0	0	0	0	0	0	Q	0	0	0	0	0	0	0
39-40	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
70741		•	•				71/5							_		_	
TOTAL	0	ũ	0	0	52	6/8	3145	3330	2052	537	140	26	8	8	17	0	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 32

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEAS0	N ( D	EC TO	FEB	)							11	L48 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							ידמ	RECTIO	N RAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-1 1-2 2-3 4-5 5-6 7-7 8-9 10-11 11-13 11-13 11-15 15-17 17-18 18-19 120-21 21-223 224-225 225-226 227-228 229-30 30-31 31-33 33-34 34-35 35-36 36-37 37-38 38-39 39-40	000800000000000000000000000000000000000	00 88 60 173 00 00 00 00 00 00 00 00 00 00 00 00 00	80042243470000000000000000000000000000000	17826131657238880000000000000000000000000000000000	3477174206396870896870896870896870896870896870896870896870896877880000000000	6077 8348348951303444955664609126460912646090000000000000000000000000000000000	3482784745615668528288000000000000000000000000000000	604364 104846446 104888 10488 10488 10488 104888 10488 104888 104	51767846832002888800000000000000000000000000000	346714877408086000000000000000000000000000000	88 60 104 1137 108 000 000 000 000 000 000 000 000 000	8771370088000000000000000000000000000000	3176789622227000000000000000000000000000000000	886644426023352 5487 000000000000000000000000000000000000	0080271777000000000000000000000000000000	80000884800000000000000000000000000000	3189354357863858330302878000000000000000000000000000000

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 33

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SPRING	SEASO	N ( I	1AR TO	YAM C	)							4	95 20	MIN	OBSER	VATIO	N PAIRS
WIND SPEED							פדת	ECTIO	N RAN	ח							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0- 1	0	20	0	0	20	20	40	40	60	20	0	20	20	0	0	0	262
1- 2	0	0	40	40	60	40	80	60	0	0	20	20	0	0	20	0	383
2- 3	0	0	40	141	101	121	60	40	101	101	0	20	20	0	20	20	787
3- 4 4- 5	20 0	60 222	181 363	242 262	242 424	222 161	121 121	101 141	101 20	121 60	0	0	0	0	0	40	1454
5- 6	Ö	40	181	181	424	242	60	121	60	20	Ö	20	Ů	0	20 0	40 0	1838 1353
<b>6</b> - 7	ŏ	121	60	181	202	161	ő	161	60	ČŎ	ŏ	Õ	ŏ	ŏ	ů	ŏ	949
7- 8	Ó	60	20	101	222	80	Ō	80	60	20	ŏ	ŏ	ŏ	Ŏ	ŏ	ŏ	646
8- 9	0	40	60	141	141	121	0	80	40	0	0	0	Ô	0	0	Ō	626
9-10	0	20	40	141	121	101	0	40	e	0	20	0	0	40	20	0	545
10-11 11-12	0	40 20	20 0	80	60	80	20	0	0	0	0	0	0	0	0	40	343
12-13	0	20	ŏ	60 60	20 60	121 40	20 20	20 20	20 0	20 20	20 0	0	40 0	0 20	0	0	363
13-14	ŏ	_0	ŏ	0	40	70	20	40	0	20	Ö	ŏ	Ŏ	20	Ů	0	262 80
14-15	ŏ	20	ŏ	ŏ	20	ŏ	20	ŏ	ŏ	ŏ	ŏ	ň	ň	ő	ŭ	ñ	60
15-16	Ō	0	Ô	Ŏ	Ō	Ŏ	Ō	Ŏ	Ŏ	Ö	ŏ	20	ŏ	ŏ	ŏ	ŏ	20
16-17	0	Ō	0	0	0	0	0	20	0	0	0	0	0	Ó	Ō	Ŏ	20 20
17-18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
18-19 19-20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20-21	Û	Û	ň	Û	0	0	0	0	0	0	0	0	0	0	0	Ü	0
21-22	ŏ	ŏ	ŏ	Ö	ŏ	ŏ	ŏ	ů	ŏ	ŏ	ŏ	Ö	ŏ	ů	ŏ	0	U
22-23	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ň	ň
23-24	0	0	Ö	0	Ö	Ō	Ō	Õ	Ŏ	Ō	Ö	Ŏ	ō	ŏ	ŏ	ŏ	ŏ
24-25	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ó	Ō
25-26	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ō	0	0
26-27 27-28	0	0	0	0	0	0	0	0	0	Ŏ	0	0	0	0	0	0	0
28-29	ň	0	Ö	ŏ	ň	ŏ	ŏ	0	0	0 0	0	Ů	0	0	0	0	U O
29-30	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ů	ŏ	ŏ	ŏ	ŏ	ŏ	Ö	ŏ	ň
30-31	Ŏ	Ŏ	Ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ
31-32	0	0	0	0	0	0	0	0	0	0	0	0	Û	Ö	Ō	Ŏ	Ŏ
32-33	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
33-34 34-35	0	0	0	0	ō	0	0	0	0	0	0	0	0	0	0	0	Ō
35-36	Ů	0	0	0	0	0	0	0	0 <b>0</b>	0	0	0	0	0	0	0	0
36-37	ŏ	Ö	Ů	ů	ŏ	ů	0	0	ů	0	Ů	0	0	0	0	U	ď
37-38	ŏ	ŏ	ŏ	Ö	ŏ	ŏ	Ö	Ö	ŏ	Ů	ŏ	Û	Ö	0	Û	ň	ņ
38-39	Ŏ	Ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ņ
39-40	0	0	0	0	0	0	0	Ö	Ö	Ö	Ö	Ö	Ö	Ŏ	Ŏ	ŏ	ŏ
TOTAL	20	686	1010	1636	2161	1515	565	969	525	383	60	101	80	60		167	
IOINE	20	000	1010	1030	-101	1010	202	707	223	202	00	TOI	00	0.0	80	141	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 34

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

SUMMER	SEASON	( J	UN TO	AUG	)							:	249 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIR	FCTTC	N BAN	מו							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0- 1	0	0	40	0	0	40	0	80	80	160	0	80	40	0	40	40	602
ĭ- 2	Ŏ	Ŏ	Ö	Ŏ	80	80	Õ	Ŏ	Ŏ	120	40	40	40	40	40	Ö	481
2- 3	0	0	0	0	40	40	0	120	281	40	80	120	120	40	40	0	923
3- 4	0	0	0	0	0	0	0	120	200	240	80	40	80	40	120	120	1044
4-5	Ō	0	0	0	. 0	40	80	40	40	160	40	120	200	0	80	40	843
5- 6	0	0	0	40	120	40	40	40	0	200	200	160	80	40	40	40	1044
6- 7 7- 8	Ŏ	0 40	0	0	120	80	120 0	40	120 160	40 40	0	40 40	120 80	0 40	0	40 40	722 803
8- 9	0	0	Ö	Ö	40	160 40	200	200 120	120	40	ŏ	281	40	40	80	80	1044
9-10	Ů	ŏ	Õ	40	40	80	40	120	40	80	80	120	40	80	0	0	642
1Ó-11	Ô	ŏ	ŏ	Ö	Ŏ	80	ŏ	ň	ŏ	Õ	80	80	120	80	ŏ	Ď	441
11-12	ŏ	ŏ	Ŏ	Ŏ	40	Ŏ	8 Ŏ	ŏ	Ŏ	Ō	120	80	40	80	Ŏ	40	481
12-13	Ŏ	Ō	Ó	0	0	Ó	40	Ō	Ö	0	0	40	Ô	0	40	0	120
13-14	0	0	40	40	40	0	0	0	0	0	0	0	80	0	0	0	200
14-15	0	0	0	Ō	Ō	0	Ō	0	0	0	0	40	0	0	0	0	40
15-16	0	0	40	0	0	0	0	0	0	0	0	40	40	0	0	0	120
16-17	Ŏ	0	40	40	0	Ŏ	0	0	0	Ü	0	0	40	0	0	0	120
17-18	0	0	80 0	0	0	0	0	0	0	U	0	0	0	0 40	40 0	0	120 40
18-19 19-20	Ů	Ö	40	0	ŏ	ŏ	ŏ	Ů	Ö	ň	ŏ	0	ů	0	ů	ŏ	40
20-21	Õ	ŏ	0	ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	å	ŏ	ő	Ŏ	ő	ŏ	Ŏ	0
21-22	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	Ď	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ň	ů
22-23	ŏ	Ď	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	80	ŏ	ŏ	80
23-24	Ŏ	Ŏ	Õ	Ŏ	Ŏ	Ŏ	Õ	Õ	Ŏ	Ō	Ö	Ö	Ō	Ö	Õ	Ŏ	Ö
24-25	0	0	40	0	0	0	0	0	0	0	0	0	0	0	0	0	40
25-26	0	0	0	Ō	O	0	0	0	0	0	0	0	0	0	0	0	0
26-27	Ō	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Ō
27-28	0	0	0	0	0	0	0	0	0	0	0	0	0	Ŏ	0	0	0
28-29 29-30	0	0	0	0	0	0	0	0	0	0	Ů	0	0	0	0	U	Ů
30-31	Ô	ŏ	Ö	Ö	ŏ	0	ů	ŏ	Ö	Ö	ŏ	ŏ	ŏ	ŏ	Ů	ň	ň
31-32	ñ	ň	ŏ	ū	ŏ	ő	ñ	ŏ	ŏ	ŏ	ő	ŏ	ŏ	ŏ	ñ	ň	ň
32-33	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	Ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ	ŏ
33-34	Ŏ	Ö	Ö	Ö	Õ	Ö	Ö	Ö	Ö	Ō	Ö	Ö	Ŏ	Ō	Ō	Ö	Ŏ
34-35	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
35-36	Ō	0	0	0	Ō	Ō	0	0	0	0	0	0	0	0	0	0	O
36-37	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
37-38	0	0	Õ	0	Õ	Ō	Õ	Ŏ	0	0	Õ	0	0	0	0	0	0
38-39 39-40	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
37.40	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	0	U
TOTAL	0	40	321	160	522	682	602	763	1044	1124	722	1325	1164	562	<b>52</b> 2	441	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 35

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WIND SPEED (KNOTS)  1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16  TOTAL  0 - 1 0 0 0 0 0 83 83 0 0 41 125 41 41 83 41 41 41 0 0 583 1 - 2 0 0 0 0 0 83 83 125 0 166 41 41 41 41 41 41 41 41 41 41 61 833 3 - 4 0 0 0 0 0 1 83 83 83 125 0 166 41 41 125 41 0 0 41 41 41 61 833 3 - 4 0 0 0 0 0 41 83 83 83 125 0 166 41 41 125 41 0 0 41 41 41 61 833 4 - 5 0 0 0 0 0 0 83 83 83 125 0 166 41 41 125 41 0 0 41 41 19 1833 5 - 6 0 0 0 0 0 1 83 83 125 0 166 41 41 125 166 125 10 0 0 1 10 83 125 83 125 83 125 83 125 83 125 83 125 83 125 84 10 0 41 1983 5 - 6 0 0 0 0 0 5 11 125 208 0 0 125 166 125 10 0 0 1 1 1083 6 - 7 0 0 0 1 14 18 3 208 250 83 125 83 125 83 125 84 14 1 1 0 0 41 41 41 958 8 - 9 0 0 0 208 333 250 0 0 0 0 0 0 0 0 0 0 0 0 1 166 88 9 10 1166 8 - 9 0 0 0 208 333 250 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 1 16 83 125 11 10 11 10 0 0 0 1 11 166 88 9 10 11 10 10 0 0 1 11 10 0 0 0 1 11 10 0 0 0 1 11 1	FALL SE	ASON (	SEP	TO N	( VO								240	20 M	IN OB	SERVA	HOIT	PAIRS
CKNOT55	UZND CDEED							מזח	ECTIO	N DAN	ın.							
1-2	(KNOTS)	1	2	3	4	5	6					11	12	13	14	15	16	TOTAL
2-3		•		-				-										
3-4		•			•			•										
\$\frac{4}{-5}\$		-																
5-6         0         0         0         41         125         208         0         41         41         83         208         250         83         10         0         11         41         83         208         250         83         0 <td></td> <td>•</td> <td>•</td> <td>•</td> <td></td> <td>-</td> <td></td> <td></td> <td></td>		•	•	•											-			
6-7 0 0 41 41 83 208 250 83 0 0 41 41 41 83 0 0 916 8-9 0 0 0 541 166 291 125 0 0 0 0 0 0 0 41 41 0 1166 8-9 0 0 0 0 298 333 250 0 0 0 0 0 0 0 41 41 41 41 81 958 9-10 0 0 0 291 125 375 83 0 0 0 0 0 0 0 41 41 41 41 41 81 958 10-11 0 0 0 0 41 83 83 83 125 41 0 0 0 0 0 0 0 0 0 0 0 0 316 10-11 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 0 0 33 0 208 12-13 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 0 0 33 3 13-14 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 291 125 14-15 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 291 125 14-15 0 0 0 0 0 125 14 41 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 291 125 14-15 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		ŏ			ŏ													
8-9 0 0 0 0 208 333 250 0 0 0 0 0 0 41 41 41 41 41 41 31 61 958 9-10 0 0 0 0 291 125 375 83 0 0 0 0 0 0 0 41 0 0 0 916 10-11 0 0 0 0 41 83 83 125 41 0 0 0 0 0 0 0 83 0 208 12-13 0 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 83 0 208 12-13 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 333 13-14 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 125 125 83 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		Ö	Ó	41	41	83			83	0	0	41	41	41	83	Ö		
9-10		0		-					-		-	•	•					
10-11		0							•	•	•	•	-					
11-12		0									•	•	•	•		•	•	
12-13         0         0         0         125         125         83         0		0	•	•							•	•		-			-	
13-14         0         0         0         41         41         0         0         0         0         0         0         0         0         41         125           14-15         0         0         0         0         125         41         41         0         0         41         0		V						•			•	ŭ		•	•		•	
14-15		ň		•	-						•	ň	•	•	•		•	
15-16         0 <td></td> <td>ň</td> <td>•</td> <td>-</td> <td></td> <td></td> <td></td> <td></td> <td>ň</td> <td>•</td> <td></td> <td></td> <td></td> <td></td> <td>•</td> <td>-</td> <td></td> <td></td>		ň	•	-					ň	•					•	-		
17-18       0 <td></td> <td>ŏ</td> <td>Ŏ</td> <td>Ŏ</td> <td>Ŏ</td> <td></td> <td></td> <td></td> <td>Ŏ</td> <td>Ŏ</td> <td></td> <td>Ŏ</td> <td>Ŏ</td> <td></td> <td>Ŏ</td> <td>Ŏ</td> <td>ŏ</td> <td></td>		ŏ	Ŏ	Ŏ	Ŏ				Ŏ	Ŏ		Ŏ	Ŏ		Ŏ	Ŏ	ŏ	
18-19	16-17	Ŏ	Ó	0		0	0	0	0		0	0		0	0	Ō	Ó	
19-20		0					•		0		•	0			-	•	Q	0
20-21		0	•			-	-	-	0		•	0		-	-	-	0	0
21-22         0 <td></td> <td>0</td> <td>•</td> <td></td> <td>-</td> <td>•</td> <td>•</td> <td></td> <td>0</td> <td></td> <td>•</td> <td>•</td> <td></td> <td>0</td> <td></td> <td>•</td> <td>0</td> <td>0</td>		0	•		-	•	•		0		•	•		0		•	0	0
22-23		Ü	•			•	•	•	U		-	•	•	U	-	•	Ų	U
23-24		ň	•				•	•	ň		•	•		•	•	-	V	Ů
24-25		ň	•			ñ	•	•	ň		•	-		•	•	•	ň	ň
25-26		ŏ	ŏ	•	•	ŏ	ŏ	•	ŏ	•	•	•	•	•	•	•	ŏ	ň
27-28       0 <td></td> <td>Ŏ</td> <td></td> <td>Ŏ</td> <td>Ŏ</td> <td>Ŏ</td> <td>ŏ</td> <td>Ŏ</td>		Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ	Ŏ		Ŏ	Ŏ	Ŏ	ŏ	Ŏ
28-29 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		0				•	•		•		0	•			0	0	0	0
29-30		0				•	•	•	•		•	•	•	•		•	0	0
30-31 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		0			•		•		0		•	•		•	•	•	0	0
31-32 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		Ü					•	•	Ų		•	•		•	•	•	Ų	Ŭ
32-35 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		ň							ň		•	•		•	•	•	Ŭ	V
33-34		ŏ				•	•	•	ŏ		•	•		•	•	•	ň	ň
35-36	33-34	Ŏ			Ŏ	Ŏ	Ŏ	Ŏ	Ŏ		ŏ	Ŏ		•	-	ŏ	ŏ	ŏ
36-37		0			•		•		0		•	0		0	0	Ō	Ó	Ö
37-38 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		Ų							0		•	•		0	•	•	0	0
38-39 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		U			•		-	•	Ō	•	•	•		•	•	•	0	0
39-40 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		n			•				Ų			•		•	•	•	Ü	0
												-	•	•				Ü
TOTAL 0. 0 41 1291 1500 2166 1083 500 500 333 791 500 291 333 333 333		•	•	•	·	·	•	•	•	•	•	J	•	•	•	,	U	U
	TOTAL	0.	0	41	1291	1500	2166	1083	500	500	333	791	500	291	333	333	333	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 36

**GULF HIGH** 

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WINTER	SEASO	N ( D	EC TO	FEB	)							5	04 20	MIN	OBSER	VATION	PAIRS
WIND SPEED							DIR	ECTIO	N BAN	מו							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
	1 00999880000000000000000000000000000000	2 0999990000000000000000000000000000000	3 000000000000000000000000000000000000	4 0 39 0 159 10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	5 0 0 19 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	6 0 0 9 3 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0					11 00 39 158 415 77 158 77 00 00 00 00 00 00 00 00 00 00 00 00	12 199 198 1798 1778 1780 00 00 00 00 00 00 00 00 00 00 00 00 0	13 190 199 199 100 000 000 000 000 000 000	14 39 00 99 198 59 00 00 00 00 00 00 00 00 00 00 00 00 00	15 0091789917891119911111111111111111111111	16 9099999999000000000000000000000000000	TOTAL  138 277 615 1250 1349 1607 19249 674 416 218 218 00 00 00 00 00 00 00 00 00 00 00 00 00
TOTAL	317	178	138	138	39	59	436	1011	615	1349	1567	1309	575	714	1031	515	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 37

**GULF HIGH** 

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

WIND SPEED DIRECTION BAND (KNOTS) 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15  U-1 0 0 0 0 0 0 51 0 154 154 51 0 51 0 0 1-2 0 0 51 0 51 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 51 1 1 0 1; 4 51 1;	OTAL 463 567 237 855
(KNOTS)     1     2     3     4     5     6     7     8     9     10     11     12     13     14     15       0-1     0     0     0     0     0     51     0     154     154     51     0     51     0     0       1-2     0     0     51     0     0     0     0     51     309     51     0     0     0	0 0 0 0 51 1 1 0 1; 4 51 1;	463 567 237 855
1-2 0 0 51 0 51 0 0 0 51 309 51 0 0 0	0 51 1 1 0 1; 4 51 1; 3 0 1;	567 237 855
2-3	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	54425716492664600000000000000000000000000000000

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 38

GULF HIGH
BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION
( IN PERCENT OCCURRENCE X100 )

SUMMER	SEASO	и ( J	ин то	AUG	)							;	876 20	MIN	OBSER	VATION	PAIRS
WIND SPEED (KNOTS)	1	2	3	4	5	6	DIR 7	ECTIO 8	N BAN	D 10	11	12	13	14	15	16	TOTAL
0-1 1-2 2-3 3-4 4-5 5-6 6-7 7-8 8-9 9-11 11-12 12-13 13-14 14-15 15-16 16-17 17-18 18-19 19-21 21-22 22-24 24-25 25-26 24-25 25-27 27-28 28-29 230-31 31-33 33-34 34-35 35-36 37-38 38-39 39-40	00000001120000000000000000000000000000	0 1120 341 991 691 110000000000000000000000000000	0 112343340 110000000000000000000000000000000	57 1227 227 220 1111 000 000 000 000 000 000 000 00	11 04 22 11 11 00 00 00 00 00 00 00 00 00 00 00	22 0 341 111 111 111 0 0 0 0 0 0 0 0 0 0 0 0 0	22 34 30 111 110 00 00 00 00 00 00 00 00 00 00	0 22 79 11 22 21 10 00 00 00 00 00 00 00 00 00 00 00 00	11923488544000000000000000000000000000000000	21122117 1757470000000000000000000000000000000000	8555465172272000000000000000000000000000000000	457 796 2179 3655 223388 1157 0 110 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	34586285022544120000000000000000000000000000000	0 1184 1194 1194 1205 1536 1137 1137 1137 1137 1137 1137 1137 11	1109749971110000000000000000000000000000	00079981299001000000000000000000000000000000	308 3429 17692 14761 14761 1470 114 114 114 110 110 110 110 110 110 11

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 39

GULF HIGH

### BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SEA	ASON	( SEP	TO N	( VO								341	L 20 M	IN OB	SERVA	TION	PAIRS
WIND SPEED							DIP	ECTIO	N BAL	מנ							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
	1 0000000000000000000000000000000000000	2 000000000000000000000000000000000000	3 52 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	4 9000000000000000000000000000000000000	5 0000000000000000000000000000000000000	6 0000000000000000000000000000000000000	7 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	8 000000000000000000000000000000000000	9 877099 1178090 0000000000000000000000000000000000	1 05735187777888552 000000000000000000000000000000000	11 558759768899000000000000000000000000000000000	12 29714181607179567 1148700000000000000000000000000000000000	13	1	15 0586997611000000000000000000000000000000000	16 800770008779000000000000000000000000000	TOTAL 615 938 997 1524 703 8595 586 4985 388 992 00 00 00 00 00 00 00 00 00 00 00 00 00
TOTAL	0	351	322	117	87	146	117	117	703	1612	879	1466	1906	909	791	469	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 40

#### **GULF TROPICAL DEPRESSION**

## BIVARIATE DISTRIBUTION OF WIND SPEED AND DIRECTION ( IN PERCENT OCCURRENCE X100 )

FALL SEA	ASON (	SEP	TO NO	<i>i</i> )								267	20 M	IN OBS	ERVA	TION P	AIRS
WIND SPEED							DIR	FCTI	N BAN	D							
(KNOTS)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	TOTAL
0-1 1-23 4-56 5-78 8-10 10-11 11-13 11-13 11-13 11-13 11-13 11-13 12-14 14-16 11-13 11-18 18-19 120-21 221-223 221-224 221-223 221-223 221-223 221-223 221-223 221-223 221-23 221-23 231-336 3	000000000000000000000000000000000000000	000000000000000000000000000000000000000	000000000000000000000000000000000000000			00000000000000000000000000000000000000	0000000422231299944472231129931122142233129944472337	0007779444447997422184374997777700000000000000000000000000000	0000 3771 1744 1877 1877 1877 1877 1877 1124 1124 1124 1124 1124 1124 1124 11	0 0 0 0 0 37 112 37 0 0 0 37 0 0 0 0 37 0 0 0 0 37 0 0 0 0	000000004777077000077333333333333333333	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		000000000000000000000000000000000000000	000000000000000000000000000000000000000	000000000000000000000000000000000000000	0 0 0 74 74 262 149 636 674 749 861 1086 1123 786 486 337 262 187 74 37 149 37 37 0 0 0 0
IUIAL	Ų	U	U	U	3/	524	209/	39/0	2546	411	299	112	0	0	0	0	

THE WIND DIRECTION IS DIVIDED INTO 16 BANDS(22.5 DEGREES IN EACH BAND).
BAND 1 IS CENTERED ABOUT 0.0 DEGREES AND INDICATES WINDS BLOWING FROM THE NORTH.
INCREASING BAND NUMBERS REFLECT INCREASING WIND ANGLES(MEASURED CLOCKWISE).

Table 41

Maximum Winter Wind Speeds, knots,

at New Orleans and Lake Charles

	Direction Band															
<u>Year</u>	1	_2	_3	_4	_5	_6	_7	_8	_9	10	11	12	13	14	<u>15</u>	16
						New	Orle	ans								
1961	18	16	17	19	16	19	13	18	20	25	17	14	20	22	18	19
1962	23	20	16	18	18	13	16	18	22	25 25	18	18	20	13	19	20
1963	24	22	20	24	17	12	14	16	19	20	17	11	12	18	17	24
1964	19	17	22	17	18	13	16	20	20	20	25	26	22	16	17	17
1965	21	15	17	15	17	13	23	19	24	15	14	20	23	16	14	18
1966	16	17	19	17	16	18	24	19	22	30	10	16	19	14	23	16 22
1967 1968	17 19	15 17	12 18	16 15	13 15	12 14	13 16	14 22	18 18	18 18	17 20	17 17	12 20	13 20	16 21	22 19
1969	20	16	16	24	25	24	17	21	17	16	16	18	19	16	20	18
1970	18	14	18	16	15	15	17	16	18	14	14	17	15	14	18	20
1971	17	17	20	19	16	12	17	30	24	19	16	10	16	14	20	16
1972	21	15	16	11	13	14	20	16	16	21	14	5	7	9	18	19
1973 1974	20 18	17 15	16 11	14 14	15 13	17 16	20 21	29 24	17 17	15 22	14 16	11 15	18 20	27 21	18 17	18 16
1975	19	16	14	13	15	17	15	17	20	16	19	15	18	17	16	22
					r	ako	Char	100								
						arc	Ollar	ICS								
10/0	0.0	0.7	0.0	0.0	- (		<b>.</b> 0	0.0	0.0	0.0	10	1.0	1.5	10	1.5	10
1963 1964	28 20	27 20	22 21	23 22	16 17	17 16	18 18	22 16	20 19	22 20	19 22	16 18	15 19	12 18	15 18	18 21
1965	17	17	16	17	16	16	18	18	24	16	14	17	15	14	21	18
1966	22	16	16	16	20	17	22	24	24	13	10	11	8	16	18	20
1967	20	18	18	13	11	14	16	15	16	17	14	12	15	12	14	17
1968	20	20	16	17	14	16	16	18	20	14	13	8	10	15	17	22
1969 1970	18 22	15 15	15 16	15 17	19 18	16 20	15 19	19 13	17 19	17 17	13 20	15 26	15 12	17 14	19 17	19 26
							_	_	-							22
1971 1972	20 20	17 21	19 20	21 22	19 11	26 16	18 18	20 16	23 19	20 17	18 10	14 12	16 10	19 9	20 18	22 18
1973	20	22	17	16	14	21	22	24	25	15	19	20	16	16	21	22
1974	20	16	14	14	20	20	19	26	17	17	16	15	27	30	14	18
1975	24	18	18	16	24	19	28	22	27	15	13	16	16	17	21	22

Table 42

Maximum Spring Wind Speeds, knots,

at New Orleans and Lake Charles

		Direction Band														
Year	_1	_2	_3	_4	_5	_6	_7	_8	_9	<u>10</u>	11	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>
						New	Orle	eans								
1961	20	17	19	18	18	17	15	25	18	17	15	18	19	22	21	17
1962 1963	15 18	15 18	17 16	15 15	18 15	15 17	18 16	22 18	18 22	18 25	14 18	20 15	14 18	18 16	19 18	18 16
1964	16	16	14	14	19	20	18	22	25	25	26	17	19	15	18	16
1965	15	16	15	14	13	15	16	17	18	20	12	14	20	14	16	15
1966	18	14	20	17	20	15	17	20	18	21	15	16	22	15	17	16
1967	16	16	16	15	19	16	24	20	20	16	13	11	22	17	19	17
1968 1969	18 17	14 17	14 18	15 20	16 18	18 18	22 26	20 29	23 32	17 14	14 17	19 19	22 18	17 15	17 17	17 17
1970	14	16	16	15	15	17	20	17	28	15	18	12	15	12	15	17
1971	18	19	12	16	12	14	16	20	24	14	14	17	14	14	16	21
1972	20	18	16	15	15	18	22	16	20	14	17	14	14	11	12	21
1973 1974	16	15	13	12	17	27	27	28	23	18	16	17	18	16	25	17
1974	15 19	19 15	15 15	14 16	18 19	19 20	18 16	20 18	22 23	16 17	18 16	15 17	17 24	16 15	20 16	16 18
		_	-							-,						
					Ī	ake	Char	les								
1963	18	20	18	18	14	14	20	22	24	22	18	15	11	12	24	20
1964 1965	18 18	20 18	18 20	18 13	19 18	16 16	18 17	20 19	26 22	25 15	24 15	16 10	19 10	20 10	23 18	20 18
1966	18	17	18	18	16	15	18	19	20	19	20	15	9	21	21	16
1967	18	25	18	17	15	16	19	23	20	16	11	9	7	7	16	16
1968	18	15	14	15	13	12	18	18	23	16	13	13	15	20	21	14
1969	18	15	18	18	17	26	20	18	28	16	14	17	13	13	20	17
1970	15	17	15	19	12	20	15	21	24	16	13	18	15	14	18	15
1971	22	20	18	17	15	15	18	18	22	17	19	9	16	22	21	23
1972 1973	17 20	13 20	23 24	23 25	22 22	15 24	18 32	18 28	20 27	19 18	15 18	10 18	12 16	18 17	16 20	15 27
1974	17	18	14	19	16	16	23	28	25	18	15	12	15	5	16	20
1975	17	13	15	19	16	26	25	26	26	14	14	12	16	20	17	22

Table 43

Maximum Summer Wind Speeds, knots,

at New Orleans and Lake Charles


15 16
•
10 12
10 13 10 11
12 10
11 13
13 11
12 9
10 13 7 12
12 30
14 9
12 21
12 10
13 11
10 11 12 10
12 10
11 16
11 11 13 12
10 10 9 7
5 10
11 10
12 8
12 8
14 11
14 12 16 11
17 23

Table 44

Maximum Fall Wind Speeds, knots,

at New Orleans and Lake Charles

	Direction Band 1 2 3 4 5 6 7 8 0 10 11 12 12 14 15 16															
Year	_1	_2	_3	_4	_5	_6	_7	_8	_9	<u>10</u>	11	12	13	14	<u>15</u>	16
						New	0rle	ans								
1961 1962	15 15	18 16	18 15	20	16	15	23	22	17	10	9	12	13	11	17	16
1963	16	24	20	15 16	16 20	10 17	12 15	15 22	19 14	14 12	12 10	12 14	15 17	15 13	15 17	17 18
1964	20	14	14	17	17	12	13	17	15	15	10	8	10	13	14	13
1965	15	15	14	14	13	10	6	6	16	17	15	11	12	10	13	13
1966	15	16	13	13	15	15	10	11	12	14	8	9	12	14	22	21
1967	15	14	15	15	15	16	18	16	14	15	17	14	20	14	14	16
1968 1969	23 18	18 19	20 15	17 15	18 16	15 13	14 18	18 15	21 14	15 13	16 10	25 8	19	15	17	17 18
1970	18	15	15	12	17	15	16	13 17	20	9	12	12	10 16	15 13	17 13	18
1971	16	16	14	15	13	13	12	13	9	16	9	5	8	11	11	13
1972	17	15	15	21	15	17	16	24	24	7	13	13	15	14	15	14
1973	15	17	15	17	17	15	17	18	24	18	14	10	18	14	13	17
1974 1975	16 21	14 23	16 15	15 12	16	13 15	15	14	16 17	8 13	11	8	11	15	17	14
1975	21	23	13	12	13	13	12	19	17	13	16	6	10	13	17	17
					$\bar{\Gamma}$	ake	Char	<u>les</u>								
1963 1964	18	25	18	23	24	18	17	21	18	20	14	11	8	11	13	20
1965	19 13	18 14	20 13	18 11	15 12	12 13	15 14	15 13	20 11	18 15	13 15	10 14	14 9	8 5	15 13	13 16
1966			10			13		13		13	13	17		J	15	10
1967	17	16	12	18	16	16	13	15	18	14	14	16	14	16	16	14
1968	16	14	17	17	8	17	20	17	21	14	12	13	18	15	18	18
1969	17	13	15	13	16	15	16	17	17	10	14	13	7	13	11	18
1970	23	22	18	17	20	19	19	21	23	15	16	10	9	12	17	18
1971	19	20	17	16	16	14	14	15	16	18	15	6	9	8	10	16
1972 1973	16 14	14 17	14 18	17 20	16	18	15 16	15	18	13	11	8	10	12	14	15
1973	14 18	17 17	18 14	20 14	25 12	17 19	16 16	18 19	20 17	13 10	13 13	8 6	15 14	16 10	16 15	20 18
1975	20	18	16	18	12	14	18	22	14	11	9	5	7	12	16	23






Figure 1. Location of weather stations (WS) in the Atchafalaya Bay




Figure 2. Prototype data collected in the Atchafalaya Bay




Figure 3. Isobaric contour map of FOR circulation




Figure 4. Prototype wind measurements during FOR circulation




Figure 5. Isobaric contour map of FGR circulation




Figure 6. Prototype wind measurements during FGR circulation




Figure 7. Isobaric contour map of CH circulation




Figure 8. Prototype wind measurements during CH circulation




Figure 9. Isobaric contour map of CR circulation




Figure 10. Prototype wind measurements during CR circulation




Figure 11. Isobaric contour map of GR circulation




Figure 12. Prototype wind measurements during GR circulation




Figure 13. Isobaric contour map of GH circulation




Figure 14. Prototype wind measurements during GH circulation




Figure 15. Comparison of wind speed at stations WS-E and WS-B




Figure 16. Comparison of wind direction at stations WS-E and WS-B  $\,$ 




Figure 17. Comparison of wind speed at stations WS-E and WS-D




Figure 18. Comparison of wind direction at stations WS-E and WS-D




Figure 19. Frequency of occurrence of wind speed during winter (all weather types)




Figure 20. Frequency of occurrence of wind direction during winter (all weather types)




Figure 21. Frequency of occurrence of wind speed during spring (all weather types)

## WIND DIRECTION (SPRING SEASON, MAR-MAY)




Figure 22. Frequency of occurrence of wind direction during spring (all weather types)




Figure 23. Frequency of occurrence of wind speed during summer (all weather types)




Figure 24. Frequency of occurrence of wind direction during summer (all weather types)




Figure 25. Frequency of occurrence of wind speed during fall (all weather types)




Figure 26. Frequency of occurrence of wind direction during fall (all weather types)




Figure 27. Cumulative yearly frequency of occurrence of wind speed (all weather types)




Figure 28. Cumulative yearly frequency of occurrence of wind direction (all weather types)




Figure 29. Duration of CH winds during winter

## CONTINENTAL HIGH (MAR-MAY)




Figure 30. Duration of CH winds during spring




Figure 31. Duration of CH winds during summer



Figure 32. Duration of CH winds during fall




Figure 33. Duration of FOR winds during winter




Figure 34. Duration of FOR winds during spring




Figure 35. Duration of FOR winds during summer




Figure 36. Duration of FOR winds during fall




Figure 37. Duration of CR winds during winter




Figure 38. Duration of CR winds during spring




Figure 39. Duration of CR winds during summer




Figure 40. Duration of CR winds during fall




Figure 41. Duration of GR winds during winter




Figure 42. Duration of GR winds during spring




Figure 43. Duration of GR winds during summer




Figure 44. Duration of GR winds during fall




Figure 45. Duration of FGR winds during winter




Figure 46. Duration of FGR winds during spring




Figure 47. Duration of FGR winds during summer




Figure 48. Duration of FGR winds during fall




Figure 49. Duration of GH winds during winter




Figure 50. Duration of GH winds during spring




Figure 51. Duration of GH winds during summer




Figure 52. Duration of GH winds during fall




Figure 53. Duration of GTD winds during fall




Figure 54. Duration of winds during winter (all weather types)




Figure 55. Duration of winds during spring (all weather types)




Figure 56. Duration of winds during summer (all weather types)




Figure 57. Duration of winds during fall (all weather types)