

TA7
E8
no. ERDC/EL
SR-00-15

LIBRARY
USE ONLY

ERDC/EL

Environmental Laboratory



**US Army Corps
of Engineers®**

Engineer Research and
Development Center

Field Survey of Contaminant Concentrations in Existing Wetlands in the San Francisco Bay Area

C. R. Lee, D. L. Brandon, J. W. Simmers, H. E. Tatem
R. A. Price, and S. P. Miner

November 2000

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.



PRINTED ON RECYCLED PAPER

Field Survey of Contaminant Concentrations in Existing Wetlands in the San Francisco Bay Area

by C. R. Lee, D. L. Brandon, J. W. Simmers,
H. E. Tatem, R. A. Price, and S. P. Miner

Environmental Laboratory
U.S. Army Engineer Research and Development Center
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Final report

Approved for public release; distribution is unlimited

SUMMARY

There is an increased public awareness of the importance of wetlands and a heightened interest in restoration and creation of wetlands using dredged material. Dredged material is being tested for potential use in wetland creation and restoration projects. In order to evaluate the acceptability of wetland creation and restoration with dredged material, establishment of some form of reference wetland baseline from which to make informed evaluations is often necessary. Test data must be interpreted in relationship to realistic circumstances. The reference baseline is usually chosen from the particular location where wetlands will be created or restored.

The objective of this study was to determine the concentrations of contaminants in sediments, plants and animals in existing wetlands near proposed wetland creation sites and to establish a reference wetland baseline for the San Francisco Bay area. The data collected would become an initial wetland baseline that can be used to interpret and put perspective on results of wetland testing of dredged material from the San Francisco Bay area.

Thirteen naturally occurring wetlands were sampled in marine, estuarine and freshwater locations along San Francisco and Suisun Bays and in the Sacramento River Basin. Wetland sediment, plant and animal samples were collected and transported to the U.S. Army Engineer Waterways Experiment Station (WES) for processing and analysis. Samples were analyzed for metals, butyltins, petroleum aromatic hydrocarbons, pesticides and polychlorinated biphenyls.

The naturally-occurring wetlands in the San Francisco Bay area and the adjacent estuarine and freshwater areas contained relatively low levels of most metal, PCBs, PAHs, butyltin, and pesticide contaminants in soil/sediment, plants, and animals. Metals such as lead, chromium and arsenic appeared to have elevated concentrations in some plants and animals. There was a very depauperate faunal component in all the naturally-occurring wetlands surveyed, that may be the result of a more subtle impact. This survey was conducted toward the end of a five year drought in the region. This climatic condition no doubt influenced the existing fauna available for sampling.

CONTENTS

	Page
SUMMARY.....	1
LIST OF FIGURES.....	3
LIST OF TABLES.....	5
PREFACE.....	6
I. INTRODUCTION.....	7
Background.....	7
Purpose and Scope.....	7
Objectives.....	7
II. FIELD SURVEY.....	8
Approach.....	8
Methods and Materials.....	8
Site Selection.....	8
Plant and Animal Identification.....	9
Field Collection Technique.....	9
Laboratory Procedures.....	10
Results and Discussion.....	22
Chemical Analysis.....	22
III. CONCLUSIONS AND RECOMMENDATIONS.....	81
REFERENCES.....	82
APPENDIX A: FIELD SURVEY/PLANT AND ANIMAL TISSUE CONCENTRATIONS.....	83

LIST OF FIGURES

	Page
Figure II-1 Sampling was Accomplished by Helicopter.....	12
Figure II-2 Field Survey Map for Sites 1-7, 9, and 14.....	13
Figure II-3 Field Survey Map for Site 8.....	14
Figure II-4 Field Survey Map for Sites 10-13.....	15
Figure II-5 Field Sampling Site 1 Hamilton Air Force Base (Reference).....	16
Figure II-6 Field Sampling Site 2 Sears Point Road/ Cullinan Ranch	16
Figure II-7 Field Sampling Site 3 Dutchman Slough/ Cullinan Ranch	17
Figure II-8 Field Sampling Site 4 Lower Tubbs Island Wetland	17
Figure II-9 Field Sampling Site 5 Petaluma Marsh.....	18
Figure II-10 Field Sampling Site 7 Sonoma Baylands	18
Figure II-11 Field Sampling Site 8 Deepwater Slough	19
Figure II-12 Field Sampling Site 9 Roe Island, NWS Concord	19
Figure II-13 Field Sampling Site 10 Browns Island	20
Figure II-14 Field Sampling Site 11 Near Franks Tract	20
Figure II-15 Field Sampling Site 13 Staton Island, South Fork	21
Figure II-16 Field Sampling Site 14 Suisun Slough (Reference)	21
Figure II-17 Mean Zinc Concentrations in Soil from Sites 1 through 14	49
Figure II-18 Mean Copper Concentrations in Soil from Sites 1 through 14 ...	50
Figure II-19 Mean Cadmium Concentrations in Soil from Sites 1 through 14	51
Figure II-20 Mean Arsenic Concentrations in Soil from Sites 1 through 14	52
Figure II-21 Mean Selenium Concentrations in Soil from Sites 1 through 14	53
Figure II-22 Mean Mercury Concentrations in Soil from Sites 1 through 14 ..	54
Figure II-23 Mean Lead Concentrations in Soil from Sites 1 through 14	55
Figure II-24 Mean Nickel Concentrations in Soil from Sites 1 through 14 ...	56
Figure II-25 Mean Chromium Concentrations in Soil from Sites 1 through 14	57
Figure II-26 Mean Zinc Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	58

Figure II-27 Mean Copper Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	58
Figure II-28 Mean Cadmium Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	58
Figure II-29 Mean Arsenic Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	59
Figure II-30 Mean Selenium Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	59
Figure II-31 Mean Mercury Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	59
Figure II-32 Mean Lead Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	60
Figure II-33 Mean Nickel Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	60
Figure II-34 Mean Chromium Concentrations Grouped by Site of Plants <i>Spartina</i> (SF), <i>Salicornia</i> (SO), <i>Scirpus</i> (SS), and <i>Typha</i> (TL)	60
Figure II-35 Mean Zinc Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	61
Figure II-36 Mean Copper Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	61
Figure II-37 Mean Chromium Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	61
Figure II-38 Mean Arsenic Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	62
Figure II-39 Mean Selenium Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	62
Figure II-40 Mean Mercury Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	62
Figure II-41 Mean Lead Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	63
Figure II-42 Mean Nickel Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	63
Figure II-43 Mean Chromium Concentrations Grouped by site of Organisms <i>Corbicula</i> (CO), <i>Modiolus</i> (MD), <i>Nassarius</i> (SN)	63

LIST OF TABLES

	Page
Table II-1 Wetland Field Survey List.....	25
Table II-2 Butyltin Concentration in Naturally-occurring Wetland Plants and Soils.....	26
Table II-3 Heavy Metal Concentration in Naturally-occurring Wetland Plants and Soils.....	28
Table II-4 PCB Concentration in Naturally-occurring Wetland Plants and Soils.....	31
Table II-5 PAH Concentration in Naturally-occurring Wetland Plants and Soils.....	34
Table II-6 Pesticide Concentration in Naturally-occurring Wetland Plants and Soils.....	43
Table II-7 Summary of Concentrations of Contaminants in Soils Under Field Conditions.....	64
Table II-8 Summary of Concentrations of Contaminants in Plants Under Field Conditions.....	67
Table II-9 Summary of Concentrations of Contaminants in Animals Under Field Conditions.....	75

PREFACE

This report presents the results of a field survey of existing wetlands in the San Francisco Bay area performed for Messrs. Brian Walls, Duke Roberts, Mark Dettle and Tom Kendall, project managers at the San Francisco District of the US Army Corps of Engineers. The study was conducted by the US Army Engineer Waterways Experiment Station (WES) during the period July 1990 through September 1991.

Work was performed by Dr. Charles R. Lee, Soil Scientist; Dr. Henry E. Tatem, Zoologist; Dr. John W. Simmers, Research Biologist; Mr. Richard A. Price, Research Agronomist; Mr. Dennis L. Brandon, Statistician; Contaminant Mobility and Regulatory Criteria Group (CMRCG), Environmental Processes and Effects Division (EPED), Environmental Laboratory (EL); and Mr. Scott P. Miner, Ecologist, San Francisco District, U.S. Army Corps of Engineers (SPN).

Animal bioassessment acknowledges Mr. Lawrence Bird (ASCI Corporation), and Ms. Heather Holifield, Mr. Michael Pendavis, and Mr. Johnny McGuffie (University Contract Students) for conducting the laboratory portion of this study. Plant bioassessment acknowledges Ms. Erika Seals and Ms. Elizabeth Tominey (University Contract Students) for laboratory processing and analysis of sediment and plant tissue. Heavy metals analyses of samples from the plant bioassay were provided by the Analytical Laboratory Group, Environmental Engineering Division, USAE-WES, Vicksburg, Mississippi. All other chemical analyses of sediment, water, and tissues were performed by Dr. Eric Crecelius, Battelle/Marine Sciences Laboratory, Sequim, WA.

At the time of the study, work was conducted under the supervision of Dr. Bobby L. Folsom, Jr., Chief, CMRCG; Mr. Donald L. Robey, Chief, EPED; Dr. John Harrison, Chief, EL, and Mr. Roderick A. Chisholm II, Chief, Environmental Branch, SPN.

At the time of the study, COL Larry Fulton, EN, was Commander and Director during the preparation of this report. Technical Director was Dr. Robert W. Whalin.

This report should be cited as follows:

Lee, C.R., Brandon, D.L., Simmers, J.W., Tatem, H.E., Price, R.A., and Miner, S.P. (2000). "Field Survey of Contaminant Concentrations in Existing Wetlands in the San Francisco Bay Area," ERDC/EL SR-00-15, U.S. Army Engineer Research and Development Center, Vicksburg, MS.

I. INTRODUCTION

Background

Each year the Corps of Engineers dredges sediment from harbors and channels throughout the San Francisco Bay Area to maintain navigation and commerce. Productive use of dredged material to restore and create wetlands has gained more interest in recent years. Suitable dredged material has been used productively in over 120 locations across the U. S. (US Army EM-1110-2-5026). The importance of wetlands to the productivity of estuaries has been realized even more recently in the San Francisco Bay Area. A heightened public interest has emerged to restore wetland acreage that has dwindled away over the past 50 years. Consequently, there has been increased public desire to create and restore wetlands in the San Francisco Bay area in recent years. Dredged material was thought to be of potential value in wetland creation or restoration.

Purpose and Scope

The purpose of this report is to describe the results of a field survey of existing wetland sites in the San Francisco Bay Area and to establish a wetland baseline data set.

Objectives

The objectives of the survey were:

- 1) to identify relatively undisturbed wetlands typical of the San Francisco Bay area;
- 2) to collect samples of the dominant plants, animals (where present) and wetland soil from selected marine and estuarine wetlands in the vicinity of San Francisco Bay;
- 3) to analyze each plant tissue, animal tissue, and soil sample for the presence of contaminants, including toxic heavy metals, polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAHs), and butyltin compounds such as Tributyltin (TBT);

4) to document the location and appearance of each of the sampling sites for future reference by map location and through aerial photographs.

II. FIELD SURVEY

Approach

The interpretation of the results of biological and chemical testing of a sediment to evaluate its potential use in wetland creation requires a yardstick (i.e. reference database) for comparison. For this reason, naturally-occurring wetlands in the San Francisco Bay area were identified and the soil/sediment and the indigenous plant and animal communities were sampled. In coordination with personnel of the USACE San Francisco District, sites were selected that are considered to be typical undisturbed wetlands by the District and the Federal and State resource agencies. Unfortunately, since settlement, the San Francisco Bay Area has been the source of anthropomorphic disturbance that has resulted in both modification of the pre-settlement landscape and the introduction of numerous plant and animal species. As a result, it is not always possible to locate the desired species or a sufficient biomass of the desired species for analysis. During the summer of 1990, when the field survey was conducted, animal species, live populations of bivalve mollusks in particular, were not present in either the marine or estuarine wetlands. The paucity of animals has certainly limited the comparative value of the following survey, however, the plant and sediment/soil collections do provide a suitable data base for the establishment of a baseline for wetlands in the San Francisco Bay Area comparison with the species employed in the mesocosm test procedures.

Methods and Materials

Site Selection. The initial selection of the wetlands to be considered was provided by the USACE San Francisco District personnel and consisted of wetlands selected within known wetland refuges and locations generally thought to have been little affected by anthropomorphic activities during recent years, or as in the case of Site 8, the disturbance was well documented and the site was of

interest to the District.

The potential sites were surveyed from the air and if there were no obvious reasons to reject the site, such as proximity to industrial activity, a location within the site was selected for the field sampling (Figure II-1). On several occasions if the helicopter employed by the field collection personnel was not able to land, if the field crew was not able to reach a suitable plant community due to dense vegetation once landed, or if the appropriate plant species were not present, the collection site was relocated as required.

Plant and Animal Identification. Plants and animals collected were identified using appropriate resource materials and reports such as Fernald (1950), Josselyn (1983), and Gosner (1979). Where appropriate, local botanists were consulted to confirm the plant identifications in the field.

Field Collection Technique. Locations of field collections, water salinity, and plant and animal species collected are given in Table II-1. In the marine wetland areas Spartina foliosa and Salicornia subterminalis were the predominate species. Spartina was collected from the low marsh (nearest the water) and Salicornia was collected from the high marsh (the zone inland from the Spartina). In general, two samples of Spartina labeled A and B were collected from the intertidal, low marsh and two samples of Salicornia labeled C and D were collected from the more upland, high marsh. In the estuarine and fresh water areas of the survey, the dominant low and high marsh plants were collected as before, labels A and B designated low marsh and C and D designated high marsh. Due to the variability of the less marine habitats, plant species varied between Typha, Scirpus, and Salicornia, depending on the wetland area. Each sample collected consisted of the amount of plant material that could be encompassed by a 28.7-cm square made from a folding carpenter's ruler, or 823.7 cm². The plants were clipped 5 cm above the ground. Plant material from each sample was placed in a Ziploc bag or a trash can liner, depending on the amount of vegetation, and placed on ice for shipment to the WES.

After the plants were collected a soil sample of the surface material was collected from each of the sampling locations, A-D. Soil samples were placed in Ziploc bags and placed on ice for shipment to WES. A refractometer was used to measure the salinity of the water.

Any animals suitable as sentinel species were collected at each field collection site. Animal collections represent a composite sample rather than two discrete points within the field site. When a single species was found in sufficient numbers to provide appropriate biomass for chemical analysis, the animals were collected, placed in Ziploc bags and placed on ice for shipment to WES.

At each site the location was plotted on a map (Figures II-2, II-3, and II-4) and an aerial photograph was made of the site, looking north at 30- to 45-m altitude (Figure II-5 - II-16).

Laboratory Procedures. Plant, animal, and soil samples were shipped and stored at 4°C until processed. The plant leaf samples were rinsed three times in reverse osmosis (RO) purified water blotted with paper towels, and weighed. Animal sentinel species (mollusks) were rinsed in RO water and the soft tissues removed from the shells. Only the soft tissues were submitted for chemical analysis. Soil samples were composited to form one sample from each field site. Plant tissue, animal tissue, and soils were placed in acid-washed, hexane-rinsed glassware and shipped at 4°C to Battelle Pacific Northwest Laboratory for chemical analysis. Freeze dried and ground sediment samples were analyzed by energy dispersive X-ray fluorescence for As, Cr, Cu, Ni, Pb and Zn (Nielson and Sanders 1983). The other metals were analyzed by atomic absorption spectrometry (AA) after the sediment was totally dissolved in a mixture of nitric, perchloric and hydrofluoric acids at an elevated temperature (130 degrees C) in a sealed Teflon container. Mercury was quantified by cold vapor atomic absorption spectrometry and the other metals (Ag, Cd and Sb) were quantified by Zeeman graphite furnace AA with matrix modifiers. Sediment and tissue samples were extracted with a mixture of methylene chloride, tropolone and sodium sulfate for

the Tributyltin (TBT) analyses. The extract was derivatized and analyzed by gas chromatography with a flame photometric detector (GC-FPD) similar to the method of Ungery et al. (1986). Sediments were analyzed for base-neutral acids using US EPA Method 625, which indicates solvent extraction, column cleanup and the quantification by GC/MS. The PCBs and DDT were analyzed by US EPA Method 8080 which quantified by GC-ECD. Volatiles were analyzed by US EPA Method 624 using GC/MS. All samples for tributyltin analyses were placed in hexane rinsed and oven-dried amber glass containers and frozen prior to shipping.



Figure II-1. Sampling was Accomplished by Helicopter.




Figure II-2. Field Survey Map for Sites 1-7, 9, and 14.




Figure II-3. Field Survey Map for Site 8.




Figure II-4. Field Survey Map for Sites 10-13.



Figure II-5. Field Sampling Site 1 Hamilton Air Force Base (Reference)



Figure II-6. Field Sampling Site 2 Sears Point Road/ Cullinan Ranch



Figure II-7. Field Sampling Site 3 Dutchman Slough/ Cullinan Ranch




Figure II-8. Field Sampling Site 4 Lower Tubbs Island Wetland



Figure II-9. Field Sampling Site 5 Petaluma Marsh



Figure II-10. Field Sampling Site 7 Sonoma Baylands



Figure II-11. Field Sampling Site 8 Deepwater Slough



Figure II-12. Field Sampling Site 9 Roe Island, NWS Concord



Figure II-13. Field Sampling Site 10 Browns Island



Figure II-14. Field Sampling Site 11 Near Franks Tract



Figure II-15. Field Sampling Site 13 Staton Island, South Fork



Figure II-16. Field Sampling Site 14 Suisun Slough (Reference)

Results and Discussion

Chemical Analysis. Results of the analyses for metal, butyltin, PAH, PCB, and pesticide contaminants are shown in Tables II-2 through II-6. Metal concentrations (Table II-3) represent naturally-occurring background levels for the enormous San Francisco Bay area. These levels of metals result from the presence of heavy metals in the earth's crust, water borne metals, metals in tide water and any atmospheric fallout. These data represent areas thought to be relatively undisturbed, and uncontaminated by agricultural or industrial activities (Site 8 however, was the only disturbed site that had been created from dredged material). While the sediment arsenic concentrations are relatively low and range up to 23.7 mg/kg for site 1 and plant tissue contents are at or below detection limits, those few animals collected had tissue arsenic concentrations approaching or at a FDA-type tissue arsenic content of 10 ug/g (dry weight basis) for mollusks and crustacea used by Australia (Lee et al. 1991). While the few animals sampled in this field survey did not show elevated levels of chromium, zinc or lead, snails collected from the Tiburon area and used in wetland bioassays of dredged material from Oakland Harbor and J. F. Baldwin Ship Channel showed levels of chromium up to 74.9 ug/g, zinc up to 797 ug/g and lead up to 31.6 ug/g (Lee et al. 1994). These bioassays were conducted at the same time of the Field Survey and indicate that chromium, zinc and lead concentrations in certain native wetland animals may be elevated in the San Francisco Bay area. Of particular concern is the lead levels that appear to approach and exceed the 25 ug/g lead concentration established in Australia for human consumption of mollusc. Both Spartina and Salicornia plant species collected during the Field Survey showed tissue lead concentrations up to 4.9 and 5.4 ug/g, respectively. These values approach and exceed the 5.0 ug/g concentration established by the Dutch for mixed animal feed (van Driel et al. 1985). These data suggest that lead contents of some wetland plants and certain wetland animals in existing wetlands may be of concern to the foodwebs associated with these sites in the San Francisco Bay area. The presence of copper at what may appear to be an elevated level in the animals collected at the field sites is likely related to the copper-containing respiratory pigment

characteristic of the Mollusca as a group. The butyltin levels are generally near or below detection limits with the exception of tributyltin in bivalve mollusks (Table II-2). Butyltin values in boldface print are above detection limits. Modiolus collected at Site 1 contained 34.9-38.3 ug/kg tributyltin and Corbicula collected at Site 13 contained 40.7 ug/kg. These levels are the highest determined in any survey animals and probably reflect trace amounts accumulated from the water filtered by these mollusks. Since butyltins do not exist in nature, the levels reported are assumed to be the result of contamination from marine antifouling coatings. PCBs were not found above the detection limits with the exception of some trace amounts of Aroclor 1254 in the wetland soils collected at Sites 1-4 (Table II-4). As noted for butyltin compounds, PCBs are not found naturally in the environment and their presence above detection limits indicates some anthropomorphic contamination. The presence of some PAHs at levels greater than detection in the wetland soils at some sites may also be indicative of anthropomorphic influences (Table II-5). Those PAHs indicated in boldface print are above the detection limits but at the same time they are still relatively low and generally do not exceed 50-100 ug/kg. Pesticides were notably below detection limits with only a few exceptions (boldfaced in Table II-6).

The naturally-occurring wetlands in the San Francisco region that were selected for this survey appeared to be relatively uncontaminated by post-settlement agriculture and industrialization. Even Site 8 constructed on dredged material contained only low levels of the contaminants evaluated. Arsenic tissue contents observed in the few animals collected appeared to be close or at the action level established in Australia for mollusks and crustacea. Further study of arsenic in wetland foodwebs in the San Francisco Bay area appears to be warranted. Likewise, some wetland plants and animals were observed to contain elevated levels of chromium, zinc and/or lead. Lead particularly was observed to approach and exceed tissue lead contents established for plant feed mixes by the Dutch and lead concentrations in mollusks established by the Australian. Further evaluation of chromium, zinc and lead in existing wetlands of San Francisco Bay appears to be warranted.

Although the levels of anthropomorphic contaminants appear to be low, all the selected sites were characterized by a lack of animals, particularly those that could have been used as sentinel species. All the marine sites were characterized by the dead remains of what must have only recently been extensive beds of ribbed mussels. Although the plant communities have survived, there is a need to at least develop a plausible explanation for the lack of living mussels. The introduction and proliferation of a tiny exotic clam from Asia, Potamocorbula amurensis may be a contributing factor. This species out-competes and is a more efficient feeder than existing species. In the brackish and freshwater sites, the clam Corbicula was represented also by many shells and only a few live animals. The invasion of Potamocorbula amurensis also includes brackish waters such as in Suisun Bay. Snails were equally scarce on all sites but Site 8. This lack of animals is quite peculiar since the snails, and mussels are invasive species from the U. S. East Coast, and the clams are an equally opportunistic species from Asia. While it is likely that the introduction of the exotic species (Nassarius, Modiolus, and Corbicula) accompanied some disturbance of the California wetlands, these are very hardy species and would have been expected to survive subsequent disturbances. However, Potamocorbula amurensis could even be out-competing these species. It is realized that the entire San Francisco Bay area has suffered from an extensive drought over the past five years and could have contributed to the observation of few live animal species in the wetlands sampled. Likewise, the faunal component of San Francisco Bay wetlands is not well documented and perhaps the fauna may not be particularly diverse or abundant in the West Coast wetlands.

Table II-1. Wetland Field Survey Site List of Samples Collected.

Site	Location	Samples
1	Hamilton Air Force Base/Antenna Field Natural saltmarsh, 26 ppt. Selected as reference marsh for laboratory tests	<i>Spartina foliosa</i> (2) <i>Salicornia</i> sp. (2) Mussels (1) Soil (4)
2	Sears Point Road, adjacent to Cullinan Ranch, recent accreted sediment salt marsh, 30 ppt	<i>Spartina foliosa</i> (2) <i>Salicornia</i> sp. (2) Soil (4)
3	Dutchman Slough, adjacent to Cullinan Ranch, natural salt/brackish marsh, 22 ppt	<i>Spartina foliosa</i> (2) <i>Salicornia</i> sp. (2) Soil (4)
4	Lower Tubbs Island, natural salt marsh 29 ppt	<i>Spartina foliosa</i> (2) <i>Salicornia</i> sp. (2) Soil (4)
5	Petaluma Marsh, natural brackish marsh 27 ppt	<i>Spartina foliosa</i> (2) <i>Salicornia</i> sp. (2) Soil (4)
6	Castro Cove, natural salt marsh,	no permission
7	Sonoma Baylands, natural salt marsh, adjacent to potential restoration site 32 ppt	<i>Spartina foliosa</i> (2) <i>Salicornia</i> sp. (2) Soil (4)
8	Deepwater Slough, salt marsh on dredged material, some contamination, 45 ppt	<i>Salicornia</i> sp. (4) Snails (1) Soil (4)
9	Roe Island, NWS Concord, natural brackish marsh, 8 ppt	<i>Scirpus</i> sp. (4) Soil (4)
10	Browns Island, natural brackish marsh, 4 ppt	<i>Typha</i> sp. (4) Soil (4)
11	Near Franks Tract, natural freshwater marsh, a potential restoration site, <2 ppt	<i>Scirpus</i> sp. (4) Soil (4)
12	San Joaquin River, natural freshwater marsh, between Rindge & McDonald tracts	omitted
13	Staton Island, on South Fork, below Brack tract, freshwater marsh, 0 ppt	<i>Typha</i> sp. (4) <i>Corbicula</i> sp. (1) Soil (4)
14	Suisun Slough, natural brackish marsh, 10-12 ppt, selected as reference marsh for laboratory tests	<i>Scirpus</i> sp. (2) <i>Salicornia</i> sp. (2) Soil (4)

+ number of samples

Table II-2. Butyltin Concentration in Naturally-occurring Wetland Plants and Soils (Concentration in ug/kg dry-weight)

Site		Tetrabutyl Tin	Tributyl Tin	Dibutyl Tin	Monobutyl Tin
1	Soil Plants	<1.3	2.3	<1.4	<1.3
	<i>Spartina a</i>	<4.7	9.2	<4.3	19.8
	<i>Spartina b</i>	<3.3	<3.7	<3.1	<3.1
	<i>Salicornia c</i>	<1.6	<1.8	<1.5	<1.5
	<i>Salicornia d</i>	<3.2	7.4	<2.9	21.1
	Animals				
	<i>Modiolus R1</i>	<3.9	34.9	9.3	7.8
	<i>Modiolus R2</i>	<5.0	38.3	<5.0	<4.6
2	Soil Plants	0.5	2.6	3.6	17.0
	<i>Spartina a</i>	<2.3	<2.5	<2.2	<2.2
	<i>Spartina b</i>	<3.1	<3.4	<2.9	<2.9
	<i>Salicornia c</i>	9.7	6.5	<3.5	7.1
	<i>Salicornia d</i>	<3.2	7.4*	<2.9	12.5*
3	Soil Plants	3.0	2.6	<1.4	2.9
	<i>Spartina a</i>	<2.1	2.9	<2.1	<1.9
	<i>Spartina b</i>	<3.6	8.3	3.7	5.1
	<i>Salicornia c</i>	2.2	3.1	6.6	15.6
	<i>Salicornia d</i>	3.3	4.8	4.4	7.1
4	Soil Plants	<1.4	3.1	2.0	2.3
	<i>Spartina a</i>	2.7	5.2	2.5	NA
	<i>Spartina b</i>	<4.2	<4.6	<3.9	<3.9
	<i>Salicornia c</i>	3.2	6.0	19.0	64.3
	<i>Salicornia d</i>	<3.2	7.0*	<2.9	17.6
5	Soil Plants	<1.2	3.1	1.7	<1.2
	<i>Spartina a</i>	<2.2	5.2	<2.2	<2.0
	<i>Salicornia c</i>	<2.9	6.0*	<2.7	18.1
	<i>Salicornia d</i>	54.7	35.8	2.3	5.3
7	Soil Plants	2.9	2.0	9.6	2.1
	<i>Spartina a</i>	<4.1	<4.4	<3.8	<3.8
	<i>Spartina b</i>	<3.3	<3.6	<3.1	<3.1
	<i>Salicornia c</i>	8.2	9.6*	5.6	6.1
	<i>Salicornia d</i>	<5.8	12.6*	13.2	<5.3
8	Soil	2.0	2.3	<1.4	<1.3
	<i>Salicornia a</i>	2.4	4.5	2.2	53.5
	<i>Salicornia b</i>	<3.1	5.3	<3.1	<2.9
	<i>Salicornia c</i>	2.0	3.5	11.1	24.6
	<i>Salicornia d</i>	<2.3	4.0	2.8	2.1
	Animals				
	<i>Cerithidea?</i>	<1.4	3.5	4.2	1.7
	<i>Cerithidea?</i>	<0.6	1.4	0.9	1.6

Table II-2 Concluded. Butyltin Concentration in Naturally-occurring Wetland Plants and Soils (Concentration in ug/kg dry-weight)

Site		Tetrabutyl Tin	Tributyl Tin	Dibutyl Tin	Monobutyl Tin
9	Soil Plants	<1.9	3.2	9.6	2.1
	<i>Scirpus</i> a	6.1	8.3*	4.6	4.3
	<i>Scirpus</i> b	<3.2	6.5*	<2.9	<2.9
	<i>Scirpus</i> c	<3.8	8.4*	3.6	5.0
	<i>Scirpus</i> d	<5.1	14.7*	6.7	<4.6
10	Soil Plants	<1.5	3.6	<1.6	4.7
	<i>Typha</i> a	11.4*	4.7*	2.5*	9.5*
	<i>Typha</i> b	6.1*	5.7*	3.0*	4.1*
	<i>Typha</i> c	11.0*	3.9*	2.8*	<2.2
	<i>Typha</i> d	6.3*	2.2*	3.7*	14.0*
11	Soil Plants	<0.9	33.4	<0.9	<0.9
	<i>Scirpus</i> a	<4.1	5.6	5.6	<3.7
	<i>Scirpus</i> b	5.5	5.2	2.6	9.5
	<i>Scirpus</i> c	<2.2	4.1	<2.1	4.4
13	Soil Plants	<0.9	1.8	<0.9	<0.9
	<i>Typha</i> a	13.1*	8.4*	4.4*	7.0*
	<i>Typha</i> b	14.7*	6.8*	4.1*	5.5*
	<i>Typha</i> c	<3.2	<3.6	<3.0	<3.0
	<i>Typha</i> d	18.3*	4.3*	2.3*	3.3*
	Animals				
	<i>Corbicula</i>	14.6	40.7	30.1	11.8
14	Soil Plants	<1.3	3.5	1.8	2.4
	<i>Scirpus</i> a	1.2	2.2	1.1	NA
	<i>Salicornia</i> c	2.4	4.8	2.2	35.1
	<i>Salicornia</i> d	<3.1	4.4	<3.0	5.6

* indicates analyte detected in the blank

**Table II-3. Heavy Metal Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in mg/kg dry-weight)**

Site		As	Cr	Cu	Ni	Pb	Se	Zn	Cd	Hg
1	Soil	23.7	174.0	71.6	102.0	36.3	0.33	137.2	0.33	0.515
	Plants									
	<i>Spartina a</i>	<0.96	<4.7	4.63	1.96	4.1	<0.74	27.6	0.055	0.006
	<i>Spartina b</i>	<0.86	7.1	4.35	4.34	2.2	<0.64	21.2	0.032	0.015
	<i>Salicornia c</i>	<1.0	4.2	7.92	3.07	4.1	<0.77	18.0	0.051	0.01
	<i>Salicornia d</i>	0.94	<3.7	10.45	2.71	<2.4	<0.70	16.6	0.069	0.012
	Animals									
	<i>Modiolus R1</i>	8.76	4.0	23.1	7.74	1.71	4.19	71.7	3.53	0.398
	<i>Modiolus R2</i>	8.93	3.3	20.5	5.33	1.39	3.52	71.1	3.45	0.304
2	Soil	18.5	219.0	90.6	125.4	36.8	0.33	158.9	0.32	0.469
	Plants									
	<i>Spartina a</i>	<1.2	8.9	6.44	4.61	3.0	0.85	30.5	0.063	0.02
	<i>Spartina b</i>	<1.1	<6.3	7.2	4.11	4.7	<0.76	34.8	0.066	0.02
	<i>Salicornia c</i>	<0.91	1.8	10.8	2.47	0.61	<2.20	40.0	0.16	0.019
	<i>Salicornia d</i>	<1.1	10.6	13.9	6.07	3.9	<0.78	31.5	0.089	0.022
3	Soil	18.2	179.0	70.1	145.2	33.0	0.42	166.1	0.41	0.166
	Plants									
	<i>Spartina a</i>	1.27	7.2	13.7	8.76	1.39	<0.63	98.0	0.06	0.022
	<i>Spartina b</i>	1.04	7.7	13.9	9.29	1.84	<0.64	84.9	0.12	0.025
	<i>Salicornia c</i>	1.0	1.8	8.0	3.31	0.66	<0.65	26.6	0.05	0.016
	<i>Salicornia d</i>	<0.86	2.6	12.0	5.27	0.93	<0.64	25.8	0.08	0.021
4	Soil	13.4	214.0	72.6	135.5	35.7	0.17	160.1	0.31	0.439
	Plants									
	<i>Spartina a</i>	1.82	2.5	8.9	2.05	0.60	<0.68	60.9	0.07	0.014
	<i>Spartina b</i>	<1.2	<6.9	6.43	3.24	4.9	<0.85	25.9	0.043	0.012
	<i>Salicornia c</i>	<1.0	5.9	19.1	6.29	1.42	<0.79	45.7	0.29	0.038
	<i>Salicornia d</i>	<0.76	4.6	6.52	1.66	<2.10	<0.63	12.04	0.094	0.014
5	Soil	14.4	179.0	67.6	125.9	34.1	0.25	158.4	0.26	0.419
	Plants									
	<i>Spartina a</i>	0.99	8.5	11.4	9.1	2.04	<0.65	65.5	0.08	0.027
	<i>Spartina b</i>	<1.1	<5.1	8.86	3.29	<2.7	<0.81	44.9	0.22	0.008
	<i>Salicornia c</i>	<0.62	6.8	8.68	5.66	2.80	<0.65	15.7	0.039	0.012
	<i>Salicornia d</i>	<0.88	4.1	11.5	4.49	4.49	<0.66	44.3	0.06	0.018

**Table II-3. Continued. Heavy Metal Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in mg/kg dry-weight)**

Site		As	Cr	Cu	Ni	Pb	Se	Zn	Cd	Hg
7	Soil	10.6	195.0	67.5	119.8	33.8	0.33	157.5	0.33	0.469
	Plants									
	<i>Spartina a</i>	<1.1	<6.0	4.64	4.29	3.6	<0.79	28.5	0.043	0.009
	<i>Spartina b</i>	<0.99	8.9	6.1	7.40	2.7	<0.72	25.5	0.064	0.017
	<i>Salicornia c</i>	1.14	6.6	8.79	5.37	<2.30	<0.69	19.5	0.067	0.011
	<i>Salicornia d</i>	2.20	25.4	17.7	19.20	5.40	<0.73	37.5	0.10	0.059
8	Soil	5.29	224.0	35.9	72.2	20.9	<0.14	88.5	0.14	0.074
	Plants									
	<i>Salicornia a</i>	<0.003	0.4	9.7	<1.7	0.23	<1.10	27.3	0.13	0.024
	<i>Salicornia b</i>	<0.99	0.5	8.8	1.47	0.49	<0.77	57.4	0.10	0.018
	<i>Salicornia c</i>	<0.85	0.4	8.7	1.48	0.92	<0.66	36.0	0.21	0.030
	<i>Salicornia d</i>	<0.83	0.4	8.9	<0.93	0.38	<0.65	36.3	0.15	0.025
	Animals									
	<i>Cerithidea?</i> 1	11.62	2.2	93.6	10.2	1.15	1.33	401.0	1.03	0.180
	<i>Cerithidea?</i> 1	9.22	2.1	74.3	8.5	1.43	1.04	309.0	1.03	0.172
	<i>Cerithidea?</i> 2	2.5	1.2	23.5	4.5	0.82	1.47	131.4	0.34	0.055
9	Soil	19.3	183.0	68.5	107.7	85.6	0.41	142.2	0.28	0.383
	Soil (dup)	20.7	168.0	72.4	106.6	84.6	0.39	145.5	0.28	0.394
	Plants									
	<i>Scirpus a</i>	<0.71	6.4	6.83	4.26	<2.0	<0.60	41.7	0.20	0.020
	<i>Scirpus b</i>	<0.82	3.9	6.64	7.92	2.50	<0.62	43.5	0.37	0.026
	<i>Scirpus c</i>	<4.2	<4.2	10.13	2.03	2.50	<0.65	39.7	0.35	0.012
	<i>Scirpus d</i>	<0.79	3.9	5.52	1.97	<2.00	<0.58	27.2	0.19	0.024
10	Soil	17.2	126.0	67.9	93.3	47.8	0.91	135.0	0.56	0.321
	Plants									
	<i>Typha a</i>	<0.79	<3.4	4.06	2.28	<2.0	<0.63	19.0	0.035	0.016
	<i>Typha b</i>	<0.77	<3.6	4.95	2.16	<1.9	<0.63	17.8	0.067	0.026
	<i>Typha c</i>	<0.79	<3.5	5.36	2.64	2.19	<0.63	18.6	0.055	0.022
	<i>Typha d</i>	<0.87	<4.1	10.18	2.54	<2.1	<0.69	21.3	0.100	0.012

**Table II-3. Concluded. Heavy Metal Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in mg/kg dry-weight)**

Site		As	Cr	Cu	Ni	Pb	Se	Zn	Cd	Hg
11	Soil	15.3	181.0	50.3	83.3	13.7	0.16	89.8	0.22	0.283
	Plants									
	<i>Scirpus a</i>	0.87	2.7	31.1	6.70	0.87	<0.62	89.9	0.17	0.050
	<i>Scirpus b</i>	<0.89	4.0	17.4	9.39	1.03	<0.65	133.0	0.24	0.044
	<i>Scirpus c</i>	<0.79	0.7	15.3	4.47	0.49	<0.56	88.7	0.16	0.018
	<i>Scirpus d</i>	<0.84	1.9	13.6	5.81	0.76	<0.61	59.3	0.13	0.028
13	Soil	5.36	110.0	24.2	32.2	14.0	<0.14	161.7	0.55	0.059
	Plants									
	<i>Typha a</i>	<0.91	<7.1	9.41	7.40	4.0	<0.66	61.0	0.13	0.014
	<i>Typha b</i>	<0.9	8.0	7.59	9.40	2.3	<0.63	93.6	0.14	0.015
	<i>Typha c</i>	<0.87	<4.2	5.12	4.27	2.8	<0.62	34.3	0.07	0.016
	<i>Typha d</i>	<0.83	<4.0	4.0	8.31	<2.1	<0.62	98.8	0.09	0.010
	Animals									
	<i>Corbicula</i>	10.79	4.3	164.1	5.78	1.89	3.98	273.0	3.34	0.469
14	Soil	16.9	193.0	77.3	122.1	32.5	0.25	164.7	0.36	0.362
	Plants									
	<i>Scirpus a</i>	<0.79	3.3	7.7	3.47	1.18	<0.58	48.4	0.08	0.038
	<i>Salicornia c</i>	<0.92	3.6	10.1	3.78	0.99	<0.70	30.8	0.17	0.034
	<i>Salicornia d</i>	<0.95	1.7	11.4	1.85	0.71	<0.71	29.8	0.07	0.019

Table II-4. PCB Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Aroclor 1016	Aroclor 1221	Aroclor 1232	Aroclor 1242	Aroclor 1248	Aroclor 1254	Aroclor 1260
1	Soil	<30	<30	<30	<30	<30	150	<30
	Plants							
	<i>Spartina</i> a	<100	<100	<100	<100	<100	<100	<100
	<i>Spartina</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> c	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> d	<100	<100	<100	<100	<100	<100	<100
	Animals							
	<i>Modiolus</i> R1	<100	<100	<100	<100	<100	<100	<100
	<i>Modiolus</i> R2	<100	<100	<100	<100	<100	<100	<100
2	Soil	<30	<30	<30	<30	<30	83	<30
	Plants							
	<i>Spartina</i> a	<100	<100	<100	<100	<100	<100	<100
	<i>Spartina</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> c	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> d	<100	<100	<100	<100	<100	<100	<100
3	Soil	<30	<30	<30	<30	<30	210	<30
	Plants							
	<i>Spartina</i> a	<20	<20	<20	<20	<20	<20	<20
	<i>Spartina</i> b	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> c	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> d	<20	<20	<20	<20	<20	<20	<20
4	Soil	<30	<30	<30	<30	<30	120	<30
	Plants							
	<i>Spartina</i> a	<20	<20	<20	<20	<20	<20	<20
	<i>Spartina</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> c	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> d	<100	<100	<100	<100	<100	<100	<100
5	Soil	<30	<30	<30	<30	<30	<30	<30
	Plants							
	<i>Spartina</i> a	<20	<20	<20	<20	<20	<20	<20
	<i>Spartina</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> c	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> d	<20	<20	<20	<20	<00	<100	<100

Table II-4 Continued. PCB Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)

Site		Aroclor 1016	Aroclor 1221	Aroclor 1232	Aroclor 1242	Aroclor 1248	Aroclor 1254	Aroclor 1260
7	Soil Plants	<30	<30	<30	<30	<30	<30	<30
	<i>Spartina</i> a	<100	<100	<100	<100	<100	<100	<100
	<i>Spartina</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> c	<100	<100	<100	<100	<100	<100	<100
	<i>Salicornia</i> d	<100	<100	<100	<100	<100	<100	<100
8	Soil Plants	<30	<30	<30	<30	<30	<30	<30
	<i>Salicornia</i> a	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> b	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> c	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia</i> d	<20	<20	<20	<20	<20	<20	<20
	Animals							
	<i>Cerithidea?</i> 1	<100	<100	<100	<100	<100	<100	<100
	<i>Cerithidea?</i> 2	<100	<100	<100	<100	<100	<100	<100
9	Soil Plants	<30	<30	<30	<30	<30	<30	<30
	<i>Scirpus</i> a	<100	<100	<100	<100	<100	<100	<100
	<i>Scirpus</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Scirpus</i> c	<100	<100	<100	<100	<100	<100	<100
	<i>Scirpus</i> d	<100	<100	<100	<100	<100	<100	<100
10	Soil Plants	<50	<50	<50	<50	<50	<50	<50
	<i>Typha</i> a	<100	<100	<100	<100	<100	<100	<100
	<i>Typha</i> b	<100	<100	<100	<100	<100	<100	<100
	<i>Typha</i> c	<100	<100	<100	<100	<100	<100	<100
	<i>Typha</i> d	<100	<100	<100	<100	<100	<100	<100
11	Soil Plants	<30	<30	<30	<30	<30	<30	<30
	<i>Scirpus</i> a	<20	<20	<20	<20	<20	<20	<20
	<i>Scirpus</i> b	<20	<20	<20	<20	<20	<20	<20
	<i>Scirpus</i> c	<20	<20	<20	<20	<20	<20	<20
	<i>Scirpus</i> d	<20	<20	<20	<20	<20	<20	<20

**Table II-4 Concluded. PCB Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)**

Site		Aroclor 1016	Aroclor 1221	Aroclor 1232	Aroclor 1242	Aroclor 1248	Aroclor 1254	Aroclor 1260
13	Soil	<30	<30	<30	<30	<30	<30	<30
	Plants							
	<i>Typha a</i>	<100	<100	<100	<100	<100	<100	<100
	<i>Typha b</i>	<100	<100	<100	<100	<100	<100	<100
	<i>Typha c</i>	<100	<100	<100	<100	<100	<100	<100
	<i>Typha d</i>	<100	<100	<100	<100	<100	<100	<100
	Animals							
	<i>Corbicula</i>	<100	<100	<100	<100	<100	<100	<100
14	Soil	<30	<30	<30	<30	<30	<30	<30
	Plants							
	<i>Scirpus a</i>	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<20	<20
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<20	<20

Table II-5. PAH Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Acenaph- thene	Acenaph- thyrene	Anthr- acene	Benzo[a] Anthracene	Benzo[b] Fluoranthene	Benzo[k] Fluoranthene	Benzo[a] Pyrene
1	Soil	12	15	38	100	96	82	130
	Plants							
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<10
	Animals							
	<i>Modiolus R1</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Modiolus R2</i>	<10	<10	<10	<10	<10	<10	<10
2	Soil	<10	<10	15	41	58	44	63
	Plants							
	<i>Spartina a</i>	<10	<10	26	<10	<10	<10	<10
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<10
3	Soil	<10	<10	<10	22	40	26	39
	Plants							
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<10
4	Soil	<10	<10	15	47	67	50	80
	Plants							
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<10

Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)

Site		Acenaph- thene	Acenaph- thylene	Anthr- cene	Benzo[a] Anthracene	Benzo[b] Fluoranthene	Benzo[k] Fluoranthene	Benzo[a] Pyrene
5	Soil Plants	<10	<10	<10	<10	<10	<10	<10
	<i>Spartina</i> a	<10	<10	<10	<10	<10	<10	<10
	<i>Spartina</i> b	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> c	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> d	<10	<10	<10	<10	<10	<10	<10
7	Soil Plants	<10	<10	16	67	82	72	86
	<i>Spartina</i> a	<10	<10	<10	<10	<10	<10	<10
	<i>Spartina</i> b	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> c	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> d	<10	<10	<10	<10	<10	<10	<10
8	Soil Plants	<10	<10	<10	<10	15	11	11
	<i>Salicornia</i> a	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> b	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> c	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia</i> d	<10	<10	<10	<10	<10	<10	<10
	Animals							
	<i>Cerithidea?</i> 1	<10	<10	<10	<10	<10	<10	<10
	<i>Cerithidea?</i> 2	<10	<10	<10	<10	<10	<10	<10
9	Soil Plants	<10	<10	17	56	83	67	62
	<i>Scirpus</i> a	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus</i> b	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus</i> c	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus</i> d	<10	<10	<10	<10	<10	<10	<10
10	Soil Plants	19	120	97	150	211	150	130
	<i>Typha</i> a	<10	<10	<10	<10	<10	<10	<10
	<i>Typha</i> b	<10	<10	<10	<10	<10	<10	<10
	<i>Typha</i> c	<10	<10	<10	<10	<10	<10	<10
	<i>Typha</i> d	<10	<10	<10	<10	<10	<10	<10

Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Acenaph- thene	Acenaph- thyrene	Anthr- cene	Benzo[a] Anthracene	Benzo[b] Fluoranthene	Benzo[k] Fluoranthene	Benzo[a] Pyrene
11	Soil Plants	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus a</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus b</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Scirpus d</i>	<10	<10	<10	<10	<10	<10	<10
13	Soil Plants	<10	<10	<10	29	18	20	22
	<i>Typha a</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Typha b</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Typha c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Typha d</i>	<10	<10	<10	<10	<10	<10	<10
	Animals							
	<i>Corbicula</i>	<10	<10	<10	<10	<10	<10	<10
14	Soil Plants	<10	<10	<10	11	18	13	16
	<i>Scirpus a</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<10
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<10

Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)

Site		Benzo [g,h,i] perylene	Chrysene	Dibenzo [a,h] anthracene	Fluor- anthene	Fluorene	Iproto- pyrene	2-Methyl- Naph- thalene	Naph- thalene
1	Soil	110	100	19	190	<10	99	30	61
	Plants								
	<i>Spartina a</i>	<10	<10	<10	<10	15	<10	29	63
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	<20	42
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	20	50
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<20	<50
	Animals								
	<i>Modiolus R1</i>	<10	<10	<10	<10	<10	<10	45	120
	<i>Modiolus R2</i>	<10	<10	<10	<10	<10	<10	<30	61
2	Soil	68	51	10	94	<10	59	27	53
	Plants								
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	21	<50
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	32	<50
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	24	61
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	37	98
3	Soil	53	27	<10	54	<10	43	35	64
	Plants								
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	24	68
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	29	88
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<20	<50
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	28	83
4	Soil	88	53	11	110	<10	77	25	50
	Plants								
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	<20	<50
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	30	63
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	25	73
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	<20	41
5	Soil	<10	<10	<10	<10	<10	<10	15	34
	Plants								
	<i>Spartina a</i>	<10	<10	<10	<10	<10	<10	25	68
	<i>Spartina b</i>	<10	<10	<10	<10	<10	<10	<20	<50
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	<20	30
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	24	59

**Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)**

Site		Benzo [g,h,i] perylene	Chrysene	Dibenzo [a,h] anthracene	Fluor- anthene	Fluorene	Indeno- 1,2,3- pyrene	2-Methyl- Naph- thalene	Naph- thalene
7	Soil Plants	100	71	15	120	<10	87	12	26
	<i>Spartina</i> a	<10	<10	<10	<10	<10	<10	28	54
	<i>Spartina</i> b	<10	<10	<10	<10	<10	<10	<20	28
	<i>Salicornia</i> c	<10	<10	<10	<10	<10	<10	<20	30
	<i>Salicornia</i> d	<10	<10	<10	11	<10	<10	<20	30
8	Soil Plants	15	15	<10	18	<10	11	<10	56
	<i>Salicornia</i> a	<10	<10	<10	<10	<10	<10	30	16
	<i>Salicornia</i> b	<10	<10	<10	<10	<10	<10	28	89
	<i>Salicornia</i> c	<10	<10	<10	<10	<10	<10	25	68
	<i>Salicornia</i> d	<10	<10	<10	<10	<10	<10	28	89
	Animals								
	<i>Cerithidea?</i> 1	<10	<10	<10	<10	<10	<10	<30	<60
	<i>Cerithidea?</i> 2	<10	11	<10	<10	<10	<10	<30	<60
9	Soil Plants	<10	<10	17	56	83	67	NA	62
	<i>Scirpus</i> a	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> b	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> c	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> d	<10	<10	<10	<10	<10	<10	NA	<10
10	Soil Plants	19	120	97	150	211	150	NA	130
	<i>Typha</i> a	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Typha</i> b	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Typha</i> c	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Typha</i> d	<10	<10	<10	<10	<10	<10	NA	<10
11	Soil Plants	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> a	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> b	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> c	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Scirpus</i> d	<10	<10	<10	<10	<10	<10	NA	<10

Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)

Site		Benzo [g,h,i] Perylene	Chrysene	Dibenzo [a,h] anthracene	Fluor- anthene	Fluorene	Ide- no- 1,2,3- Pyrene	2-Methyl- Naph- thalene	Naph- thalene
13	Soil	<10	<10	<10	29	18	20	NA	22
	Plants								
	<i>Typha a</i>	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Typha b</i>	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Typha c</i>	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Typha d</i>	<10	<10	<10	<10	<10	<10	NA	<10
	Animals								
	<i>Corbicula</i>	<10	<10	<10	<10	<10	<10	NA	<10
14	Soil	<10	<10	<10	11	18	13	NA	16
	Plants								
	<i>Scirpus a</i>	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Salicornia c</i>	<10	<10	<10	<10	<10	<10	NA	<10
	<i>Salicornia d</i>	<10	<10	<10	<10	<10	<10	NA	<10

NA - not available

Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Phenanthrene	Pyrene
1	Soil	94	240
	Plants		
	<i>Spartina a</i>	31	<10
	<i>Spartina b</i>	13	<10
	<i>Salicornia c</i>	10	<10
	<i>Salicornia d</i>	28	<10
	Animals		
	<i>Modiolus R1</i>	37	26
	<i>Modiolus R2</i>	14	<10
2	Soil	36	120
	Plants		
	<i>Spartina a</i>	30	<10
	<i>Spartina b</i>	20	<10
	<i>Salicornia c</i>	22	<10
	<i>Salicornia d</i>	<10	<10
3	Soil	25	72
	Plants		
	<i>Spartina a</i>	14	<10
	<i>Spartina b</i>	14	<10
	<i>Salicornia c</i>	<10	<10
	<i>Salicornia d</i>	13	<10
4	Soil	42	140
	Plants		
	<i>Spartina a</i>	<10	<10
	<i>Spartina b</i>	38	<10
	<i>Salicornia c</i>	13	<10
	<i>Salicornia d</i>	<10	<10
5	Soil	<10	<10
	Plants		
	<i>Spartina a</i>	17	<10
	<i>Spartina b</i>	<10	<10
	<i>Salicornia c</i>	<10	<10
	<i>Salicornia d</i>	<10	<10

Table II-5 Continued. PAH Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Phenanthrene	Pyrene
7	Soil	45	160
	Plants		
	<i>Spartina</i> a	37	<10
	<i>Spartina</i> b	12	12
	<i>Salicornia</i> c	<10	<10
	<i>Salicornia</i> d	<10	12
8	Soil	<10	20
	Plants		
	<i>Salicornia</i> a	16	<10
	<i>Salicornia</i> b	12	<10
	<i>Salicornia</i> c	20	<10
	<i>Salicornia</i> d	15	<10
	Animals		
	<i>Cerithidea?</i> 1	<10	<10
	<i>Cerithidea?</i> 2	<10	10
9	Soil	20	89
	Plants		
	<i>Scirpus</i> a	17	19
	<i>Scirpus</i> b	18	11
	<i>Scirpus</i> c	18	19
	<i>Scirpus</i> d	11	<10
10	Soil	76	240
	Plants		
	<i>Typha</i> a	<10	<10
	<i>Typha</i> b	<10	<10
	<i>Typha</i> c	11	<10
	<i>Typha</i> d	20	<10
11	Soil	<10	<10
	Plants		
	<i>Scirpus</i> a	18	<10
	<i>Scirpus</i> b	18	<10
	<i>Scirpus</i> c	10	<10
	<i>Scirpus</i> d	14	<10

**Table II-5 Concluded. PAH Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)**

Site		Phenanthrene	Pyrene
13	Soil	20	46
	Plants		
	<i>Typha a</i>	12	10
	<i>Typha b</i>	<10	<10
	<i>Typha c</i>	18	<10
	<i>Typha d</i>	<10	<10
	Animals		
	<i>Corbicula</i>	<10	<10
14	Soil	13	33
	Plants		
	<i>Scirpus a</i>	<10	<10
	<i>Salicornia c</i>	<10	<10
	<i>Salicornia d</i>	<10	<10

Table II-6 Pesticide Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Aldrin	a-BHC	b-BHC	d-BHC	g-BHC	Chlor-dane	4,4-DDD	4,4-DDE	4,4-DDT
1	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Spartina a</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	Animals									
	<i>Modiolus R1</i>	<10	<10	<10	<10	<10	<10	<10	<10	<10
	<i>Modiolus R2</i>	<10	<10	<10	<10	<10	<10	<10	<10	<10
2	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Spartina a</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
3	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Spartina a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Spartina b</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia d</i>	<2.0	2.3	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
4	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Spartina a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
5	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Spartina a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia d</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0

Table II-6 Pesticide Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Aldrin	a-BHC	b-BHC	d-BHC	g-BHC	Chlor-dane	4,4-DDD	4,4-DDE	4,4-DDT
7	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	3.6	<3.0
	Plants									
	<i>Spartina</i> a	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Spartina</i> b	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia</i> c	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia</i> d	<20	<20	<20	<20	<20	<30	<20	<20	<20
8	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Salicornia</i> a	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia</i> b	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia</i> c	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia</i> d	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	Animals									
	<i>Cerithidea?</i> 1	<10	<10	<10	<10	<10	<10	<10	<10	<10
	<i>Cerithidea?</i> 2	<10	<10	<10	<10	<10	<10	<10	<10	<10
9	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Scirpus</i> a	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Scirpus</i> b	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Scirpus</i> c	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Scirpus</i> d	<20	<20	<20	<20	<20	<30	<20	<20	<20
10	Soil	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0
	Plants									
	<i>Typha</i> a	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Typha</i> b	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Typha</i> c	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Typha</i> d	<20	<20	<20	<20	<20	<30	<20	<20	<20
11	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Scirpus</i> a	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Scirpus</i> b	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Scirpus</i> c	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Scirpus</i> d	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0

Table II-6 Pesticide Concentration in Naturally-occurring Wetland Plants and Soils
 (Concentration in ug/kg wet-weight)

Site		Aldrin	a-BHC	b-BHC	d-BHC	g-BHC	Chlor-dane	4,4-DDD	4,4-DDE	4,4-DDT
13	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Typha a</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Typha b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Typha c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Typha d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	Animals									
	<i>Corbicula</i>	<10	<10	<12	<24	<10	<10	<10	<115	<30
14	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants									
	<i>Scirpus a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia d</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0

Table III-6 Continued. Pesticide Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)

Site		Dieldrin	Endo-sulfan I	Endo-II	Endo-Sulfate	Endrin Aldehyde	Hepta-chlor	Hepta-chlor	Meth-oxy-chlor	Toxa-phene
1	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<200
	Plants									
	<i>Spartina a</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	Animals									
	<i>Modiolus R1</i>	<10	<10	<10	<10	<10	<10	<10	<10	<500
	<i>Modiolus R2</i>	<10	<10	<10	<10	<10	<10	<10	<10	<500
2	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<200
	Plants									
	<i>Spartina a</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<20	<20	<20	<200
3	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<200
	Plants									
	<i>Spartina a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<200
	<i>Spartina b</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<200
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<200
	<i>Salicornia d</i>	<2.0	2.3	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<200
4	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<200
	Plants									
	<i>Spartina a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<200
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20

Table II-6 Continued. Pesticide Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg wet-weight)

Site		Dieldrin	Endo-sulfan I	Endo-sulfan II	Endo-sulfan Sulfate	Endrin	Endrin Aldehyde	Hepta-chlor	Hepta-chlor	Methoxychlor	Toxaphene
								chlor	Epoxide	chlor	
5	Soil Plants	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	<i>Spartina a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Salicornia d</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
7	Soil Plants	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	3.6	<3.0	<3.0
	<i>Spartina a</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Spartina b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Salicornia c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Salicornia d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
8	Soil Plants	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	<i>Salicornia a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia b</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia d</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	Animals										
	<i>Cerithidea?</i> 1	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10
	<i>Cerithidea?</i> 2	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10
9	Soil Plants	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	<i>Scirpus a</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Scirpus b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Scirpus c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Scirpus d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20

Table II-6 Concluded. Pesticide Concentration in Naturally-occurring Wetland Plants and Soils
(Concentration in ug/kg Wet Weight)

Site		Dieldrin	Endo-sulfan I	Endo-sulfan II	Sulfate	Endrin	Endrin Aldehyde	Hepta-chlor	Hepta-chlor	Met-oxy-chlor	Toxa-phene
			I	II				Epoxide			
10	Soil	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0	<5.0
	Plants										
	<i>Typha a</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Typha b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Typha c</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Typha d</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
11	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants										
	<i>Scirpus a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Scirpus b</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Scirpus c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Scirpus d</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
13	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants										
	<i>Typha a</i>										
	<i>Typha b</i>	<20	<20	<20	<20	<20	<30	<20	<20	<20	<20
	<i>Typha c</i>										
	<i>Typha d</i>	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20
	Animals										
	<i>Corbicula</i>	16	<10	<10	<10	18	<10	42	<10	<10	
14	Soil	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0	<3.0
	Plants										
	<i>Scirpus a</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia c</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
	<i>Salicornia d</i>	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0

TYPE=Soil




Figure II-17. Mean Zinc Concentrations in Soil from Sites 1 through 14.

TYPE=Soil




Figure II-18. Mean Copper Concentrations in Soil from Sites 1 through 14.

TYPE=Soil




Figure II-19. Mean Cadmium Concentrations in Soil from Sites 1 through 14.




Figure II-20. Mean Arsenic Concentrations in Soil from Sites 1 through 14.

TYPE=Soil




Figure II-21. Mean Selenium Concentrations in Soil from Sites 1 through 14.

TYPE=Soil




Figure II-22. Mean Mercury Concentrations in Soil from Sites 1 through 14.

TYPE=Soil




Figure II-23. Mean Lead Concentrations in Soil from Sites 1 through 14.

TYPE=Soil




Figure II-24. Mean Nickel Concentrations in Soil from Sites 1 through 14.




Figure II-25. Mean Chromium Concentrations in Soil from Sites 1 through 14.




Figure II-26. Mean Zinc Concentrations of Plants, *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.




Figure II-27. Mean Copper Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.




Figure II-28. Mean Cadmium Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.

MEAN OF AS BY TYPE GROUPED BY SITE

AS MEAN




Figure II-29. Mean Arsenic Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.

MEAN OF SE BY TYPE GROUPED BY SITE

SE MEAN




Figure II-30. Mean Selenium Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.

MEAN OF HG BY TYPE GROUPED BY SITE

HG MEAN




Figure II-31. Mean Mercury Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.




Figure II-32. Mean Lead Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.




Figure II-33. Mean Nickel Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.




Figure II-34. Mean Chromium Concentrations of Plants *Spartina* (SF), *Salicornia* (SO), *Scirpus* (SS), and *Typha* (TL) Grouped by Site.

MEAN OF ZN BY TYPE GROUPED BY SITE

ZN MEAN




Figure II-35. Mean Zinc Concentrations of Organisms *Corbicula* (CO), *Modiolus* (MD), *Nassarius* (SN) Grouped by site.

MEAN OF CU BY TYPE GROUPED BY SITE

CU MEAN




Figure II-36. Mean Copper Concentrations of Organisms *Corbicula* (CO), *Modiolus* (MD), *Nassarius* (SN) Grouped by site.

MEAN OF CD BY TYPE GROUPED BY SITE

CD MEAN




Figure II-37. Mean Chromium Concentrations of Organisms *Corbicula* (CO), *Modiolus* (MD), *Nassarius* (SN) Grouped by site.

MEAN OF AS BY TYPE GROUPED BY SITE




Figure II-38. Mean Arsenic Concentrations of Organisms Corbicula (CO), Modiolus (MD), *Nassarius* (SN) Grouped by site.

MEAN OF SE BY TYPE GROUPED BY SITE




Figure II-39. Mean Selenium Concentrations of Organisms Corbicula (CO), Modiolus (MD), *Nassarius* (SN) Grouped by site.

MEAN OF HG BY TYPE GROUPED BY SITE




Figure II-40. Mean Mercury Concentrations of Organisms Corbicula (CO), Modiolus (MD), *Nassarius* (SN) Grouped by site.

MEAN OF PB BY TYPE GROUPED BY SITE




Figure II-41. Mean Lead Concentrations of Organisms *Corbicula* (CO), *Modiolus* (MD), *Nassarius* (SN) Grouped by site.

MEAN OF NI BY TYPE GROUPED BY SITE




Figure II-42. Mean Nickel Concentrations of Organisms *Corbicula* (CO), *Modiolus* (MD), *Nassarius* (SN) Grouped by site.

MEAN OF CR BY TYPE GROUPED BY SITE




Figure II-43. Mean Chromium Concentrations of Organisms *Corbicula* (CO), *Modiolus* (MD), *Nassarius* (SN) Grouped by site.

Table II-7 Summary of Concentrations of Contaminants in Soils Under Field Conditions
(Concentrations in mg/kg, dry-weight for metals, and ug/kg, wet-weight all others)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
<u>Metals</u>	Mean	Range	Mean	Range	Mean	Range
As	14.87	5.29 - 23.7	18.5	16.9 - 19.3	10.3	5.3 - 15.3
Cr	197.7	174.0 - 224.0	167.5	126.0 - 193.0	145.5	110.0 - 181.0
Cu	68.6	35.9 - 90.6	71.5	67.9 - 77.3	37.3	24.2 - 50.3
Ni	120.8	72.2 - 145.2	107.4	93.3 - 122.1	57.8	32.2 - 83.3
Pb	32.9	20.9 - 36.8	62.6	32.5 - 85.6	13.9	13.7 - 14.0
Se	0.28*	<0.14 - 0.42	0.49	0.25 - 0.91	0.15*	<0.14 - 0.16
Zn	146.67	88.5 - 166.1	146.9	135.0 - 164.7	125.8	89.8 - 161.7
Cd	0.30	0.33 - 0.41	0.37	0.28 - 0.56	0.39	0.22 - 0.55
Hg	0.364	0.074 - 0.515	0.365	0.321 - 0.394	0.171	0.059 - 0.283
<u>Butyltins</u>						
Tetrabutyltin	1.76*	<1.2 - 2.9	1.6*	<1.3 - <1.9	<0.9 ^s	<0.9 [@]
Tributyltin	2.57	2.0 - 3.1	3.4	3.2 - 3.6	17.6	1.8 - 33.4
Dibutyltin	3.01*	<1.4 - 89.6	4.3*	<1.6 - 9.6	<0.9 ^s	<0.9 [@]
Monobutyltin	3.83*	<1.3 - 17.0	3.1	2.1 - 4.7	<0.9 ^s	<0.9 [@]
<u>PCBs</u>						
Aroclor 1016	<30 ^s	<30 [@]	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]
Aroclor 1221	<30 ^s	<30 [@]	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]
Aroclor 1232	<30 ^s	<30 [@]	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]
Aroclor 1242	<30 ^s	<30 [@]	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]
Aroclor 1248	<30 ^s	<30 [@]	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]
Aroclor 1254	93.3 ^s	<30 - 210	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]
Aroclor 1260	<30 ^s	<30 [@]	<36.7 ^s	<30 - <50	<30 ^s	<30 [@]

* : This mean contains at least one less than value.

@ : Every variable in this set was this same value.

^s : All values were less than detection limits.

Table II-7 Continued. Summary of Concentrations of Contaminants in Soils Under Field Conditions (Concentrations in ug/kg, wet-weight basis)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
<u>PAHs</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
Acenaph-thene	10.3 [*]	<10 - 12	13.0 [*]	<10 - 19	<10 ^s	<10 ^e
Acenaph-thylene	10.7 [*]	<10 - 15	46.7 [*]	<10 - 120	<10 ^s	<10 ^e
Anthracene	16.3 [*]	<10 - 38	41.3 [*]	<10 - 97	<10 ^s	<10 ^e
Benzo [a] Anthracene	42.4 [*]	<10 - 100	72.3	11 - 150	19.5 [*]	<10 - 29
Benzo [b] Fluoranthene	52.6 [*]	<10 - 96	104.0	18 - 211	14 [*]	<10 - 18
Benzo [k] Fluoranthene	42.1 [*]	<10 - 82	76.7	13 - 150	15 [*]	<10 - 20
Benzo [a] Pyrene	59.9 [*]	130 - <10	69.3	16 - 130	16 [*]	<10 - 22
Benzo [g,h,i] perylene	63.4 [*]	<10 - 110	13.0 [*]	<10 - 19	<10 ^s	<10 ^e
Chrysene	46.7 [*]	<10 - 100	46.7 [*]	<10 - 120	<10 ^s	<10 ^e
Dibenzo [a,h] anthracene	12.1 [*]	<10 - 19	41.3 [*]	<10 - 97	<10 ^s	<10 ^e
Fluor-anthene	85.1 [*]	<10 - 190	72.3	11 - 150	20	11 - 29
Fluorene	<10 ^s	<10 ^e	104.0	18 - 211	18	18 [*]
Ideeno-1,2,3-pyrene	55.1 [*]	<10 - 99	76.7	13 - 150	16.5	13 - 20
2-Methyl-Naphthalene	22.0 [*]	<10 - 30	NA	NA	NA	NA
Naphthalene	49.1	26 - 64	69.3	16 - 130	19	16 - 22
Phenan-threne	37.4 [*]	<10 - 94	15.0 [*]	<10 - 20	16.5	13 - 20
Pyrene	108.9 [*]	<10 - 240	28.0 [*]	<10 - 46	39.5	33 - 46

* : This mean contains at least one less than value.

^e : Every variable in this set was this same value.

^s : All values were less than detection limits.

NA: Not available.

Table II-7 Concluded. Summary of Concentrations of Contaminants in Soils Under Field Conditions (Concentrations in ug/kg, wet-weight basis)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
<u>Pesticides</u>						
Aldrin	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
a-BHC	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
b-BHC	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
d-BHC	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
g-BHC	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Chlordane	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
4,4-DDD	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
4,4-DDE	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
4,4-DDT	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Dieldrin	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Endosulfan I	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Endosulfan II	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Endosulfan sulfate	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Endrin	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Endrin Aldehyde	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Heptachlor	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Heptachlor Epoxide	3.09 ^o	3.6 - <3.0	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Methoxychlor	<3.0 ^s	<3.0 ^o	<3.7 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o
Toxaphene	<115.6 ^s	<200 - <3.0	<4.0 ^s	<3.0 - <5.0	<3.0 ^s	<3.0 ^o

* : This mean contains at least one less than value.

o : Every variable in this set was this same value.

s : All values were less than detection limits.

Table II-8 Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, dry-weight basis)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
Metals						
As: <i>Spartina</i>	1.14*	<0.86 - 1.82	NA	NA	NA	NA
<i>Salicornia</i>	0.91*	<0.003 - 2.20	<0.94 ^s	<0.92 - <0.95	NA	NA
<i>Scirpus</i>	NA	NA	<1.46 ^s	<0.71 - <4.2	0.85*	0.79 - <0.87
<i>Typha</i>	NA	NA	<0.81 ^s	<0.77 - <0.87	<0.88 ^s	<0.83 - <0.91
Cr: <i>Spartina</i>	6.65*	2.5 - 8.9	NA	NA	NA	NA
<i>Salicornia</i>	4.99*	0.4 - 25.4	2.65	1.7 - 3.6	NA	NA
<i>Scirpus</i>	NA	NA	4.34*	3.3 - 6.4	2.33	0.7 - 4.0
<i>Typha</i>	NA	NA	<3.65 ^s	<3.4 - <4.1	5.83*	<4.0 - 8.0
Cu: <i>Spartina</i>	8.05	4.35 - 13.9	NA	NA	NA	. NA
<i>Salicornia</i>	10.7	6.52 - 19.1	10.75	10.1 - 11.4	NA	NA
<i>Scirpus</i>	NA	NA	7.36	5.52 - 10.13	19.4	13.6 - 31.1
<i>Typha</i>	NA	NA	6.14	4.06 - 10.18	6.53	4.0 - 9.41
Ni: <i>Spartina</i>	5.20	1.96 - 9.29	NA	NA	NA	NA
<i>Salicornia</i>	4.45*	<0.93 - 19.20	2.82	1.85 - 3.78	NA	NA
<i>Scirpus</i>	NA	NA	3.93	1.97 - 4.26	6.59	4.47 - 9.39
<i>Typha</i>	NA	NA	2.41	2.16 - 2.64	5.35	4.27 - 9.40
Pb: <i>Spartina</i>	2.81*	0.60 - 4.90	NA	NA	NA	NA
<i>Salicornia</i>	2.07*	0.23 - 5.40	0.85	0.71 - 0.99	NA	NA
<i>Scirpus</i>	NA	NA	2.04*	1.18 - 2.50	0.79	0.49 - 1.03
<i>Typha</i>	NA	NA	2.05*	<1.9 - 2.19	2.8*	<2.1 - 4.0
Se: <i>Spartina</i>	0.73*	<0.63 - 0.85	NA	NA	NA	NA
<i>Salicornia</i>	<0.82 ^s	<0.63 - <2.20 ^s	<0.70 - <0.71		NA	NA
<i>Scirpus</i>	NA	NA	<0.61 ^s	<0.58 - <0.65	<0.61 ^s	<0.56 - <0.6
<i>Typha</i>	NA	NA	<0.65 ^s	<0.63 - <0.69	<0.63 ^s	<0.62 - <0.6
Zn: <i>Spartina</i>	45.7	21.2 - 98.0	NA	NA	NA	NA
<i>Salicornia</i>	30.6	12.04 - 57.4	30.3	29.8 - 30.8	NA	NA
<i>Scirpus</i>	NA	NA	40.1	27.2 - 48.4	92.7	59.3 - 133.
<i>Typha</i>	NA	NA	19.2	17.8 - 19.0	71.9	34.3 - 98.8
Cd: <i>Spartina</i>	0.076	0.032 - 0.22	NA	NA	NA	NA
<i>Salicornia</i>	0.109	0.05 - 0.29	0.12	0.07 - 0.17	NA	NA
<i>Scirpus</i>	NA	NA	0.24	0.08 - 0.37	0.18	0.13 - 0.24
<i>Typha</i>	NA	NA	0.064	0.035 - 0.100	0.11	0.07 - 0.14
Hg: <i>Spartina</i>	0.016	0.008 - 0.027	NA	NA	NA	NA
<i>Salicornia</i>	0.022	0.01 - 0.038	0.027	0.019 - 0.034	NA	NA
<i>Scirpus</i>	NA	NA	0.024	0.012 - 0.038	0.035	0.018 - 0.0
<i>Typha</i>	NA	NA	0.019	0.012 - 0.026	0.014	0.010 - 0.0

* : This mean contains at least one less than value.

** : Every variable in this set was this same value.

^s : All values were less than detection limits.

NA: Not applicable. No plants of this species at this site.

Table II-8 Continued. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
<u>Butyltins</u>						
Tetrabutyltin:						
<i>Spartina</i>	3.24 [*]	<2.1 - 2.7	NA	NA	NA	NA
<i>Salicornia</i>	6.65 [*]	<1.6 - 54.7	2.75 [*]	<3.1 - 2.4	NA	NA
<i>Scirpus</i>	NA	NA	3.88 [*]	1.2 - 6.1	3.93 [*]	1.2 - 5.5
<i>Typha</i>	NA	NA	8.7 [*]	2.2 - 11.4	12.33 [*]	<3.2 - 18.3
Tributyltin:						
<i>Spartina</i>	4.82 [*]	<2.5 - 9.2	NA	NA	NA	NA
<i>Salicornia</i>	7.51 [*]	<1.8 - 35.8	4.6	4.4 - 4.8	NA	NA
<i>Scirpus</i>	NA	NA	8.02 [*]	2.2 - 14.7	4.97	2.2 - 5.6
<i>Typha</i>	NA	NA	4.13 [*]	2.2 - 5.7	5.78 [*]	<3.6 - 8.4
Dibutyltin:						
<i>Spartina</i>	3.07 [*]	<2.1 - 3.7	NA	NA	NA	NA
<i>Salicornia</i>	5.18 [*]	<1.4 - 13.2	2.6 [*]	<3.0 - 2.2	NA	NA
<i>Scirpus</i>	NA	NA	3.78 [*]	<2.9 - 6.7	3.43 [*]	1.1 - 5.6
<i>Typha</i>	NA	NA	3.0 [*]	2.5 - 3.7	3.45 [*]	2.3 - 4.4
Monobutyltin:						
<i>Spartina</i>	4.78 [*]	<1.9 - 19.8	NA	NA	NA	NA
<i>Salicornia</i>	15.6 [*]	<1.3 - 64.3	20.35	5.6 - 35.1	NA	NA
<i>Scirpus</i>	NA	NA	4.2 [*]	<2.9 - 5.0	5.87 [*]	<3.7 - 9.5
<i>Typha</i>	NA	NA	7.45 [*]	<2.2 - 14.0	4.7 [*]	<3.0 - 7.0
<u>PCBs</u>						
Aroclor 1016						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<55.0 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e
Aroclor 1221						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<55.0 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e
Aroclor 1232						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<55.0 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e

* : This mean contains at least one less than value.

e : Every variable in this set was this same value.

s : All values were less than detection limits.

^ : Indicates analyte detected in the blank.

NA: Not applicable. No plants of this species at these sites.

Table II-8 Continued. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
<u>PCBs</u>						
Aroclor 1242						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<55.0 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e
Aroclor 1248						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<60 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e
Aroclor 1254						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<60 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e
Aroclor 1260						
<i>Spartina</i>	<73.3 ^s	<20 - <100	NA	NA	NA	NA
<i>Salicornia</i>	<60 ^s	<20 - <100	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<100 ^s	<100 ^e	<20 ^s	<20 ^e
<i>Typha</i>	NA	NA	<100 ^s	<100 ^e	<100 ^s	<100 ^e
<u>PAHs</u>						
Acenaphthene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Acenaph-thylene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Anthracene						
<i>Spartina</i>	11.3 ^s	<10 - 26	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e

^s : This mean contains at least one less than value.

^e : Every variable in this set was this same value.

[:] : All values were less than detection limits.

NA: Not applicable. No plants of this species at these sites.

Table II-8 Continued. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
PAHs						
Benzo [a]						
Anthracene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Benzo [b]						
Fluoranthene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Benzo [b]						
Fluoranthene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Benzo [k]						
Fluoranthene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Benzo [a]						
Pyrene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Benzo [g,h,i]						
perylene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Chrysene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e

* : This mean contains at least one less than value.

e : Every variable in this set was this same value.

: All values were less than detection limits.

NA: Not applicable/Not available. No plants of this species at these sites.

Table II-8 Continued. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
Dibenzo [a,h] anthracene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Fluoranthene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	10.06 [*]	<10 - 11	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Fluorene						
<i>Spartina</i>	10.42 [*]	<10 - 15	NA	NA	NA	NA
<i>Salicornia</i>	10.06 [*]	<10 - 11	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Indeno-1,2,3- pyrene						
<i>Spartina</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Salicornia</i>	<10 ^s	<10 ^e	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
2-Methyl- Naphthalene						
<i>Spartina</i>	24.83 [*]	<20 - 32	NA	NA	NA	NA
<i>Salicornia</i>	24.31 [*]	<20 - 37	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	NA	NA	NA	NA
<i>Typha</i>	NA	NA	NA	NA	NA	NA
Naphthalene						
<i>Spartina</i>	56.17 [*]	28 - 88	NA	NA	NA	NA
<i>Salicornia</i>	57.31 [*]	16 - 98	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
Phenanthrene						
<i>Spartina</i>	20.5 [*]	<10 - 31	NA	NA	NA	NA
<i>Salicornia</i>	13.69 [*]	<10 - 37	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	14.8 [*]	<10 - 18	15	10 - 18
<i>Typha</i>	NA	NA	12.75 [*]	<10 - 20	12.5 [*]	<10 - 18
Pyrene						
<i>Spartina</i>	10.17 [*]	<10 - 12	NA	NA	NA	NA
<i>Salicornia</i>	10.13 [*]	<10 - 12	<10 ^s	<10 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e
<i>Typha</i>	NA	NA	<10 ^s	<10 ^e	<10 ^s	<10 ^e

* : This mean contains at least one less than value.

• : Every variable in this set was this same value.

° : All values were less than detection limits. NA: Not applicable.

Table II-8 Continued. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
<u>Pesticides</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
Aldrin:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
a-BHC:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	11.02 ^e	<2.0 ^e - 2.3	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
b-BHC:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
d-BHC:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
g-BHC:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Chlordane:						
<i>Spartina</i>	<20.7 ^s	<2.0 - <30	NA	NA	NA	NA
<i>Salicornia</i>	<16 ^s	<2.0 - <30	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
4,4-DDD:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
4,4-DDE:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e

^s : This mean contains at least one less than value.

^e : Every variable in this set was this same value.

^o : All values were less than detection limits.

^o : There was a less than value much higher than this highest actual number.

NA: Not applicable/Not available. No plants of this species in these sites.

Table II-8 Continued. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
Pesticides						
4,4-DDT:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Dieldrin:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Endosulfan I:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	11.02 ^s	<2.0 ^s - 2.3	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Endosulfan II:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Endosulfan sulfate:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Endrin:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Endrin						
Aldehyde:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e

^e : Every variable in this set was this same value.

^s : All values were less than detection limits.

NA: Not applicable/Not available. No plants of this species at these sites.

Table II-8 Concluded. Summary of Concentrations of Contaminants in Plants Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
Pesticides						
Heptachlor:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Heptachlor Epoxide:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Methoxychlor:						
<i>Spartina</i>	<14 ^s	<2.0 - <20	NA	NA	NA	NA
<i>Salicornia</i>	<11 ^s	<2.0 - <20	<2.0 ^s	<2.0 ^e	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e
Toxaphene:						
<i>Spartina</i>	<123.5 ^s	<2.0 - <200	NA	NA	NA	NA
<i>Salicornia</i>	<80.75 ^s	<2.0 - <200	NA	NA	NA	NA
<i>Scirpus</i>	NA	NA	<2.0 ^s	<2.0 ^e	<2.0 ^s	<2.0 ^e
<i>Typha</i>	NA	NA	<20	<20	<20 ^s	<20 ^e

^s : This mean contains at least one less than value.

^e : Every variable in this set was this same value.

^o : All values were less than detection limits.

^h : In this range there was a less than value much higher than this highest actual value.

NA: Not applicable/Not available. No plants of this species at these sites.

Table II-9 Summary of Concentrations of Contaminants in Animals Under Field Conditions (Concentrations in mg/kg metals and ug/kg butyltins)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
<u>Metals</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
As: <i>Modiolus</i>	8.85	8.76 - 8.93	NA	NA	NA	NA
<i>Cerithidea</i>	7.78	2.5 - 11.62	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	10.79	10.79 ^o
Cr: <i>Modiolus</i>	3.65	3.3 - 4.0	NA	NA	NA	NA
<i>Cerithidea</i>	1.83	1.2 - 2.2	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	4.3	4.3 ^o
Cu: <i>Modiolus</i>	21.85	20.5 - 23.1	NA	NA	NA	NA
<i>Cerithidea</i>	63.8	23.5 - 93.6	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	164.1	164.1 ^o
Ni: <i>Modiolus</i>	6.54	5.33 - 7.74	NA	NA	NA	NA
<i>Cerithidea</i>	7.73	4.5 - 10.2	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	5.78	5.78 ^o
Pb: <i>Modiolus</i>	1.55	1.39 - 1.71	NA	NA	NA	NA
<i>Cerithidea</i>	1.22	0.82 - 1.43	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	1.89	1.89 ^o
Se: <i>Modiolus</i>	3.86	3.52 - 4.19	NA	NA	NA	NA
<i>Cerithidea</i>	1.28	1.04 - 1.47	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	3.98	3.98 ^o
Zn: <i>Modiolus</i>	71.4	71.1 - 71.7	NA	NA	NA	NA
<i>Cerithidea</i>	280.5	131.4 - 309	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	273.0	273.0 ^o
Cd: <i>Modiolus</i>	3.49	3.45 - 3.53	NA	NA	NA	NA
<i>Cerithidea</i>	0.80	0.34 - 1.03	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	3.34	3.34 ^o
Hg: <i>Modiolus</i>	0.351	0.304 - 0.398	NA	NA	NA	NA
<i>Cerithidea</i>	0.136	0.055 - 0.180	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	0.469	0.469 ^o
<u>Butyltins</u>						
Tetrabutyltin						
<i>Modiolus</i>	<4.45 ^s	<3.9 - <5.0	NA	NA	NA	NA
<i>Cerithidea</i>	<1.00 ^s	<0.6 - <1.4	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	14.6	14.6 ^o
Tributyltin						
<i>Modiolus</i>	36.6	34.9 - 38.3	NA	NA	NA	NA
<i>Cerithidea</i>	2.2	1.4 - 3.5	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	40.7	40.7 ^o

= : Dry-weight basis for metals; wet-weight for butyltins.

o : Every variable in this set was this same value.

^s : All values were less than detection limits.

NA : Not applicable/Not available. No animals of this species at this site.

Note : There were no animals analyzed from the estuarine sites.

Table II-9 Continued. Summary of Concentrations of Contaminants in Animals Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
<u>Butyltins</u>						
<u>Dibutyltin</u>						
Modiolus	7.15*	<5.0 - 9.3	NA	NA	NA	NA
Cerithidea	2.55	0.9 - 4.2	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	30.1	30.1*
<u>Monobutyltin</u>						
Modiolus	6.2*	<4.6 - 7.8	NA	NA	NA	NA
Cerithidea	1.65	1.6 - 1.7	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	11.8	11.8*
<u>PCBs</u>						
<u>Aroclor 1016</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e
<u>Aroclor 1221</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e
<u>Aroclor 1232</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e
<u>Aroclor 1242</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e
<u>Aroclor 1248</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e
<u>Aroclor 1254</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e
<u>Aroclor 1260</u>						
Modiolus	<100 ^s	<100 ^e	NA	NA	NA	NA
Cerithidea	<100 ^s	<100 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<100 ^s	<100 ^e

* : Note - there were no animals analyzed from the estuarine sites.

* : This mean contains at least one less than value.

e : Every variable in this set was this same value.

s : All values were less than detection limits.

NA : Not applicable/Not available. No animals of this species at this site.

Table II-9 Continued. Summary of Concentrations of Contaminants in Animals Under Field Conditions (Concentrations in ug/kg, wet-weight)⁻

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
<u>PAHs</u>						
Acenaphthene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Acenaphthylene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Anthracene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Benzo [a]						
 Anthracene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Benzo [b]						
 Fluoranthene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Benzo [k]						
 Fluoranthene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Benzo [a]						
 Pyrene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Benzo [g,h,i]						
 Perylene						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e

= : Note; there were no animals analyzed from the estuarine sites.

- : This mean contains at least one less than value.

• : Every variable in this set was this same value.

* : All values were less than detection limits.

NA : Not applicable/Not available. No animals of this species at this site.

Table II-9 Continued. Summary of Concentrations of Contaminants in Animals Under Field Conditions (Concentrations in ug/kg, wet-weight)

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
<u>PAHs</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
Chrysene						
Modiolus	<10 ^s	<10 ^e	NA	NA	NA	NA
Cerithidea	10.5 ["]	<10 - 11	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Dibenzo [a,h]						
Anthracene						
Modiolus	<10 ^s	<10 ^e	NA	NA	NA	NA
Cerithidea	<10 ^s	<10 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Fluoranthene						
Modiolus	<10 ^s	<10 ^e	NA	NA	NA	NA
Cerithidea	<10 ^s	<10 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Fluorene						
Modiolus	<10 ^s	<10 ^e	NA	NA	NA	NA
Cerithidea	<10 ^s	<10 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Indeno-1,2,3-pyrene						
Modiolus	<10 ^s	<10 ^e	NA	NA	NA	NA
Cerithidea	<10 ^s	<10 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
2-Methyl-Naphthalene						
Modiolus	37.5 ["]	<30 - 45	NA	NA	NA	NA
Cerithidea	<30 ^s	<30 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	NA	NA
Naphthalene						
Modiolus	90.5 ["]	61 - 120	NA	NA	NA	NA
Cerithidea	<60 ^s	<60 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Phenanthrene						
Modiolus	25.5	14 - 37	NA	NA	NA	NA
Cerithidea	<10 ^s	<10 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Pyrene						
Modiolus	18 ["]	<10 - 26	NA	NA	NA	NA
Cerithidea	<10 ^s	<10 ^e	NA	NA	NA	NA
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e
Corbicula	NA	NA	NA	NA	<10 ^s	<10 ^e

- : Note; there were no animals analyzed from the estuarine sites.

" : This mean contains at least one less than value.

e : Every variable in this set was this same value.

^s : All values were less than detection limits.

NA : Not applicable/Not available. No animals of this species at this site.

Table II-9 Continued. Summary of Concentrations of Contaminants in Animals Under Field Conditions (Concentrations in ug/kg, wet-weight)^a

<u>Pesticides</u>	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>	<u>Mean</u>	<u>Range</u>
Aldrin						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
a-BHC						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
b-BHC						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<12 ^s	<12 ^e
d-BHC						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<24 ^s	<24 ^e
g-BHC						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Chlordane						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
4,4-DDD						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
4,4-DDE						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<115 ^s	<115 ^e
4,4-DDT						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<30 ^s	<30 ^e
Dieldrin						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<16 ^s	<16 ^e

= : Note; there were no animals analyzed from the estuarine sites.

* : Every variable in this set was this same value.

: All values were less than detection limits.

NA : Not applicable/Not available. No animals of this species at this site.

Table II-9 Concluded. Summary of Concentrations of Contaminants in Animals Under Field Conditions (Concentrations in ug/kg wet-weight)⁻

	Marine Sites: 1 - 8		Estuarine Sites: 9, 10, and 14		Freshwater Sites: 11 - 13	
	Mean	Range	Mean	Range	Mean	Range
<u>Pesticides</u>						
Endosulfan I						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Endosulfan II						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Endosulfan Sulfate						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Endrin						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	18 ^s	18 ^e
Endrin Aldehyde						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Heptachlor						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	42 ^s	42 ^e
Heptachlor Epoxide						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Methoxychlor						
<i>Modiolus</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	<10 ^s	<10 ^e
Toxaphene						
<i>Modiolus</i>	<500 ^s	<500 ^e	NA	NA	NA	NA
<i>Cerithidea</i>	<10 ^s	<10 ^e	NA	NA	NA	NA
<i>Corbicula</i>	NA	NA	NA	NA	NA	NA

⁻ : Note; there were no animals analyzed from the estuarine sites.

^e : Every variable in this set was this same value.

^s : All values were less than detection limits.

NA : Not applicable/Not available. No animals of this species at this site.

III. CONCLUSIONS AND RECOMMENDATIONS

The naturally-occurring wetlands in the San Francisco Bay area and the adjacent estuarine and fresh water areas appear to contain relatively low levels of most metal, PCB, PAH, butyltin, and pesticide contaminants in soil/sediment, plants, and animals. Metals such as lead, chromium and arsenic appeared to have elevated concentrations in some plants and animals. There is, however, a very depauperate faunal component in all the naturally occurring wetlands surveyed, that may be the result of a more subtle impact. The introduction and proliferation of a tiny exotic clam from Asia, Potamocorbula amurensis may be a contributing factor. This species out-competes and is a more efficient feeder than existing species. In the brackish and freshwater sites, the clam Corbicula was represented also by many shells and only a few live animals. The invasion of Potamocorbula amurensis also includes brackish waters such as in Suisun Bay. Snails were equally scarce on all sites but Site 8. This lack of animals is quite peculiar since the snails, and mussels are invasive species from the U. S. East Coast, and the clams are an equally opportunistic species from Asia. While it is likely that the introduction of the exotic species (Nassarius, Modiolus, and Corbicula) accompanied some disturbance of the California wetlands, these are very hardy species and would have been expected to survive subsequent disturbances. However, Potamocorbula amurensis could even be out-competing these species. This survey was conducted toward the end of a five year drought experienced in the region. This climatic condition no doubt influenced the existing fauna available for sampling. Further documentation of the fauna of the San Francisco Bay area wetlands appears to be warranted. In addition, further evaluation of the status of arsenic, lead and chromium in wetland foodwebs in the San Francisco Bay area.

The data presented in this report establishes an initial baseline for wetlands in the San Francisco Bay Area and can be used to interpret wetland test results for wetland creation or restoration projects. As more information becomes available, this baseline should be updated to include all ongoing and future data collection activities.

REFERENCES

- Barr, A. J., Goodnight, J. H., Sall, J. P., and Helwig, J. T., American Public Health Association. 1976. "A Users Guide to SAS 76," SAS Institute, Inc., Raleigh, NC.
- Fernald, M. L. 1950. "Gray's Manual of Botany," 8th ed. Corrected Printing 1970. Van Nostrand Co., N.Y.
- Gosner, Kenneth L. 1979. "A Field Guide to the Atlantic Seashore," Houghton Mifflin, Boston, MA.
- Josselyn, Michael. 1983. "The Ecology of San Francisco Bay Tidal Marshes: A Community Profile". Report FWS/OBS-83/23. U.S. Fish and Wildlife Service, Slidell, LA.
- Krone, C.A., Brown, D.W., Burrows, D.G., Bogar, R.G., Chan, S.L. and Varanasi, U. "A Method for Analysis of Butyltin Species and Measurement of Butyltins in Sediment and English Sole Livers from Puget Sound", Marine Environmental Research. 27. pp 1-18.
- Lee, C. R., et al. 1991. General Decision Making Framework for Management of Dredged Material, Example Application to Commencement Bay, Washington, Miscellaneous Paper D-91-1, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
- Nielson, K.K., and Sanders, R.W. 1983. "Multielement Analysis of Unweighed Biological and Geological Samples Using Backscatter and Fundamental Parameters." Adv. X-ray Anal. 26, pp 385-390.
- Rice, C.D., Espourteille, F.A., Huggett, R.J. 1987. "Analysis of Tributyltin in Estuarine Sediments and Oyster Tissue, Crassostrea virginica. Applied Organometal Chemistry., 1, pp 541-544.
- Unger, M.A., MacIntyre, W.G., Greaves, J., and Huggett, R.J. 1986. "GC Determination of Butyltins in Natural Waters by Flame Photometric Detection of Hexyl Derivatives with Mass Spectrometric Confirmation." Chemosphere 15(4), pp 461-470.
- USEPA. 1984. United States Environmental Protection Agency, "EPA Par VIII 40 CFR, Part 136, Methods 608 and 625".
- USEPA. 1986. United States Environmental Protection Agency Office of Solid Waste and Emergency Response, "Test Methods for Evaluating Solid Wastes, SW-846 Methods 8080 and 8270". 3rd Ed.

APPENDIX A

Field Survey

Plant and Animal Tissue Concentrations

a. Plant Codes

SPA Spartina Foliosa
SCI Scirpus olynei
SAL Salicornia subterminalis
TYP Typha latifolia

b. Animal Codes

SN Cerithidea ?
CB Corbicula fluminea
MO Modiolus demissus

PLANT METAL RESULTS

(Concentrations In mg/kg Dry Weight, ppm)

Ballistic Code	Sponsor Code	As	Cr	Cu	Ni	Pb	Se	Zn	Ag	Cd	Hg
245-1	R5B-SPA	1.1 U	5.1 U	8.86	3.29	2.7 U	0.81 U	44.9	0.2	0.22	0.008
245-2	R4B-SPA	1.2 U	6.9 U	6.43	3.24	4.9	0.85 U	25.9	0.13	0.043	0.012
245-3	R7A-SPA	1.1 U	6 U	4.64	4.29	3.6	0.79 U	28.5	0.07	0.043	0.009
245-4,5,6,	R2A-SPA-1,2,3	1.2 U	8.9	6.44	4.61	3.0	0.85 U	30.5	0.14	0.063	0.02
245-7	R1A-SPA	0.96 U	4.7 U	4.63	1.96	4.0	0.74 U	27.6	0.11	0.055	0.006
245-8,9	R2B-SPA-1,2	1.1 U	6.3 U	7.2	4.11	4.7	0.76 U	34.8	0.23	0.066	0.02
245-10	RID-SAL-1	0.94	3.7 U	10.45	2.71	2.4 U	0.70 U	16.6	0.01	0.069	0.012
245-11	RID-SAL-2	0.96 U	3.7 U	11.46	2.68	2.5 U	0.72 U	20.0	0.02	0.093	0.016
245-12,14	RIC-SAL-1,3	1.0 U	4.2	7.92	3.07	4.1	0.77 U	18.0	0.01	0.051	0.01
245-13,15	RIC-SAL-2,4	1.2 U	4.8 U	8.93	2.47	2.9 U	0.87 U	18.5	0.02	0.082	0.017
245-16,18	R2D-SAL-1,3	1.1 U	10.6	13.9	6.07	3.9	0.78 U	31.5	0.03	0.089	0.022
245-17,19	R2D-SAL-2,4	0.96 U	5.2 U	11.7	3.03	4.6	0.74 U	23.2	0.01 U	0.1	0.014
245-20,21	R13C-TYP-1,2	0.87 U	4.2 U	5.12	4.27	2.8	0.62 U	34.3	0.01 U	0.07	0.016
245-22,23	R13B-TYP-1,2	0.9 U	8	7.59	9.40	2.3	0.63 U	93.6	0.01 U	0.14	0.015
245-24,25	R13D-TYP-1,2	0.83 U	4 U	4	8.31	2.1 U	0.62 U	98.8	0.01 U	0.09	0.01
245-26	R13A-TYP	0.91 U	7.1 U	9.41	7.40	4.0	0.66 U	61.0	0.01 U	0.13	0.014
245-27	R10D-TYP	0.87 U	4.1 U	10.18	2.54	2.1 U	0.69 U	21.3	0.03	0.1	0.012
245-28	R10A-TYP	0.79 U	3.4 U	4.06	2.28	2.0 U	0.63 U	19.0	0.01 U	0.035	0.016
245-29	R10C-TYP	0.79 U	3.5 U	5.36	2.64	2.19	0.63 U	18.6	0.02	0.055	0.022
245-30,31	R10B-TYP-1,2	0.77 U	3.6 U	4.95	2.16	1.9 U	0.63 U	17.8	0.02	0.067	0.026
245-32	R7B-SPA	0.99 U	8.9	6.1	7.40	2.7	0.72 U	25.5	0.06	0.064	0.017
245-33	R1B-SPA	0.86 U	7.1	4.35	4.34	2.2	0.64 U	21.2	0.12	0.032	0.015
245-34,35,36 REP 1	R9B-SCI-1,2,3 REP 1	0.82 U	3.9	6.64	7.92	2.50	0.62 U	43.5	0.06	0.37	0.026
245-34,35,36 REP 2	R9B-SCI-1,2,3 REP 2	0.82 U	3.6 U	7.72	9.95	2.1 U	0.63 U	49.2	0.05	0.38	0.024
245-37	R9C-SCI	0.03 U	4.2 U	10.13	2.03	2.50	0.65 U	39.7	0.09	0.35	0.012
245-38,39	R9D-SCI-1,2	0.79 U	3.9	5.52	1.97	2.00 U	0.58 U	27.2	0.04	0.19	0.024
245-40,41,42 REP 1	R9A-SCI-1,2,3 REP 1	0.71 U	6.4	6.83	4.26	2.00 U	0.60 U	41.7	0.05	0.2	0.02
245-40,41,42 REP 2	R9A-SCI-1,2,3 REP 2	0.75 U	4.4	8.4	5.65	2.0 U	0.60 U	41.3	0.05	0.19	0.021
245-43	R7C-SAL-1	1.14	6.6	8.79	5.37	2.30 U	0.69 U	19.5	0.02	0.067	0.011
245-44	R7-SAL-2	0.93 U	5.7 U	8.88	4.04	2.40 U	0.79 U	22.4	0.02	0.14	0.017
245-45	R5C-SAL-1	0.62 U	6.8	8.68	5.66	2.80	0.65 U	15.7	0.02	0.039	0.012
245-46	R5C-SAL-2	1.18	7.9 U	10.5	5.40	2.60 U	0.90 U	45.2	0.01 U	0.083	0.013
245-47	R4D-SAL-1	0.76 U	4.6	6.52	1.66	2.10 U	0.63 U	12.04	0.01	0.094	0.014
245-48	R4D-SAL-2	0.91 U	8	11.09	2.14	2.40 U	0.79 U	30.7	0.02	0.16	0.019
245-49	R7D-SAL-1	2.20	25.4	17.7	19.20	5.40	0.73 U	37.5	0.05	0.1	0.059
245-50	R7D-SAL-2	1.00 U	7.7	8.94	2.96	2.60 U	0.86 U	22.6	0.01 U	0.1	0.016
Procedural Blank		N/A	N/A	N/A	N/A	N/A	N/A	N/A	0.01	0.01 U	0.004 U

U indicates analyte not detected at detection limit shown

N/A indicates not applicable. Note: procedural blanks are not appropriate for XRF analyses.

STANDARD REFERENCE MATERIAL

(Concentrations in mg/kg Dry Weight, ppm)

Battelle Code	Sponsor Code	As	Cr	Cu	Ni	Pb *	Se	Zn	Ag	Cd	Hg
SRM 1571 ORCHARD LEAVES, REP 1		11.2	4.3 U	12.7	1.07	45.7	0.77 U	27.8	0.01 U	0.15	0.129
SRM 1571 ORCHARD LEAVES, REP 1A		8.72	4.1 U	12.4	1.57	48.5	0.70 U	26.2	0.01 U	0.16	
SRM 1571 ORCHARD LEAVES, REP 2		10.54	3.6 U	11.68	1.49	43.5	0.60 U	25.1			
SRM 1571 ORCHARD LEAVES, REP 2A		9.97	3.5 U	11.23	1.20	43.3	0.60 U	23.9			
SRM 1571 ORCHARD LEAVES, REP 3		10.1	5	10.75	1.05	43.5	0.59 U	25.8			
Certified Value:		14 ±2	NC	12 ± 1	1.3 ±0.2	45 ±3	0.08 ±0.01	25 ±3	NC	0.11 ±0.02	0.155 ±0.015
SRM 1566A OYSTER TISSUE (RICHLAND), REP 1		14.43	3.3 U	65.7	2.20	2.9	2.03	872.0	1.28	4.09	0.059
SRM 1566A OYSTER TISSUE (MSL), REP 1		13.59	3.5	68.4	2.05	3.6	2.04	892.0	1.46	4.15	
SRM 1566A OYSTER TISSUE (RICHLAND), REP 2		14.98	4.6	64.7	2.20	2.0 U	2.36	817.0			
SRM 1566A OYSTER TISSUE (MSL), REP 2		14.95	3.1 U	64.7	2.18	2.5	2.02	837.0			
SRM 1566A OYSTER TISSUE (RICHLAND), REP 3		13.35	3.1 U	62.7	2.14	2.1	2.17	845.0			
SRM 1566A OYSTER TISSUE (MSL), REP 3		14.93	3.3 U	64.6	1.85	1.9 U	2.27	884.0			
Certified Value:		14 ±1.2	1.43 ±0.46	66.6 ± 4.3	2.25 ±0.44	71 ±0.014*	2.21 ±0.24	830 ±57	1.68 ±0.15	4.15 ±0.38	0.0642 ±0.0067

* Lead determined by ICP-MS, not XRF.

PAH RESULTS FOR WES PLANT SAMPLES

cl# 245

(Concentrations in ug/Kg Dry Weight, ppb)

Battelle Code	Sponsor Code	Date Ext'd	Date Anal'd	% Moist.	Acenaph-thene	Acenaph-thylene	Anthra-cene	Benzo[a]Anthra-cene	Benzo[b]Fluor-anthene	Benzo[k]Fluor-anthene	Benzo[a]pyrene	Benzo-(g,h,i)-perylene
BLANK 1	BLANK 1	3/18/91	4/11/91	NA	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-1	R5B-SPA	3/18/91	4/11/91	88	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-2	R4B-SPA	3/18/91	4/11/91	85	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-3	R7A-SPA	3/18/91	4/11/91	81	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-4,5,6,	R2A-SPA-1,2,3	3/18/91	4/11/91	80	10 U	10 U	26	10 U	10 U	10 U	10 U	10 U
245-7	R1A-SPA	3/18/91	4/11/91	82	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-8,9	R2B-SPA-1,2	3/18/91	4/11/91	85	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-10,	RID-SAL-1	3/18/91	4/11/91	73	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-11,	RID-SAL-2	3/18/91	4/11/91	91	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-12,14	RIC-SAL-1,3	3/18/91	4/11/91	72	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-13,15	RIC-SAL-2,4	3/18/91	4/11/91	89	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-16,18	R2D-SAL-1,3	3/18/91	4/11/91	73	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-17,19	R2D-SAL-2,4	3/18/91	4/11/91	85	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
BLANK 2	BLANK 2	11/29/90	12/12/90	NA	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-20,21	R13C-TYP-1,2	11/29/90	12/12/90	86	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-22,23	R13B-TYP-1,2	11/29/90	12/12/90	88	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-24,25	R13D-TYP-1,2	11/29/90	12/12/90	81	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-26	R13A-TYP	12/13/90	12/14/90	90	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-27	R10D-TYP	11/29/90	12/12/90	88	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-28	R10A-TYP	11/29/90	12/12/90	85	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-29	R10C-TYP	11/29/90	12/12/90	81	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-30,31	R10B-TYP-1,2	11/29/90	12/12/90	82	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-32	R7B-SPA	11/29/90	12/12/90	80	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-33	R1B-SPA	12/13/90	12/14/90	86	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-33, DUP	R1B-SPA, DUP	12/13/90	12/14/90	NA	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/12/90	77	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-37	R9C-SCI	11/29/90	12/13/90	80	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
BLANK 3:	BLANK 3	11/30/90	12/13/90	NA	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-38,39	R9D-SCI-1,2	11/30/90	12/13/90	82	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-38,39 DUP	R9D-SCI-1,2	11/30/90	12/13/90	NA	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/13/90	75	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-43	R7C-SAL-1	11/30/90	12/13/90	78	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-44	R7-SAL-2	11/30/90	12/13/90	93	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-45	R5C-SAL-1	11/30/90	12/13/90	74	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-46	R5C-SAL-2	11/30/90	12/13/90	89	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-47	R4D-SAL-1	11/30/90	12/13/90	76	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-48	R4D-SAL-2	11/30/90	12/13/90	88	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-49	R7D-SAL-1	11/30/90	12/13/90	80	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
245-50	R7D-SAL-2	11/30/90	12/13/90	89	10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U

U Indicates analyte not detected at detection limit shown

B indicates analyte present in blank associated with that sample (one method blank was run on each date)

NA indicates not applicable

PAH RESULTS FOR WES PLANT SAMPLES

(Concentrations in ug/Kg Dry Weight, ppb)

Battelle Code	Sponsor Code	Date Ext'd	Date Anal'd	Chrysene	Dibenzo- (a,h)- anthracene	Fluor- anthene	Fluorene	Indeno- 1,2,3- Pyrene	2-Methyl- Naphthalene	Naph- thalene	Phenan- threne	Pyrene
BLANK 1	BLANK 1	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
245-1	R5B-SPA	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
245-2	R4B-SPA	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	30	63	38	10 U
245-3	R7A-SPA	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	28	54	37	10 U
245-4,5,6,	R2A-SPA-1,2,3	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	21	50 U	30	10 U
245-7	R1A-SPA	3/18/91	4/11/91	10 U	10 U	10 U	15	10 U	29	63	31	10 U
245-8,9	R2B-SPA-1,2	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	32	50 U	20	10 U
245-10	RID-SAL-1	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	20 U	50 U	48	10 U
245-11	RID-SAL-2	3/18/91	4/11/91	15	10 U	10 U	10 U	10 U	190	380	130	10 U
245-12,14	RIC-SAL-1,3	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	20	50	10	10 U
245-13,15	RIC-SAL-2,4	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	20 U	50 U	28	10 U
245-16,18	R2D-SAL-1,3	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	37	98	10 U	10 U
245-17,19	R2D-SAL-2,4	3/18/91	4/11/91	10 U	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
BLANK 2	BLANK 2	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	20 U	20 U	10 U	10 U
245-20,21	R13C-TYP-1,2	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	38	92	18	10 U
245-22,23	R13B-TYP-1,2	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	20 U	42	10 U	10 U
245-24,25	R13D-TYP-1,2	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	20 U	22	10 U	10 U
245-26	R13A-TYP	12/13/90	12/14/90	10 U	10 U	10 U	10 U	10 U	20 U	47	12	10
245-27	R10D-TYP	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	28	65	20	10 U
245-28	R10A-TYP	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	29	63	10 U	10 U
245-29	R10C-TYP	11/29/90	12/12/90	10 U	27	10 U	10 U	10 U	22	51	11	10 U
245-30,31	R10B-TYP-1,2	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	20 U	25	10 U	10 U
245-32	R7B-SPA	11/29/90	12/12/90	10 U	10 U	10 U	10 U	10 U	20 U	28	12	12
245-33	R1B-SPA	12/13/90	12/14/90	10 U	10 U	10 U	10 U	10 U	20 U	42	13	10 U
245-33, DUP	R1B-SPA, DUP	12/13/90	12/14/90	10 U	10 U	16	10 U	10 U	20 U	48	18	19
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/12/90	10 U	10 U	13	10 U	10 U	20 U	18	18	11
245-37	R9C-SCI	11/29/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	20 U	10 U	10 U
BLANK 3	BLANK 3	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	30 U	10 U	10 U
245-38,39	R9D-SCI-1,2	11/30/90	12/13/90	10 U	10 U	11	10 U	10 U	20 U	30 U	11	10 U
245-38,39 DUP	R9D-SCI-1,2	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	30 U	10	10 U
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/13/90	10 U	21	22	10 U	10 U	20 U	30 U	17	19
245-43	R7C-SAL-1	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	30 U	10 U	10 U
245-44	R7-SAL-2	11/30/90	12/13/90	10 U	10 U	13	10 U	10 U	20 U	49	21	13
245-45	R5C-SAL-1	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	30 U	10 U	10 U
245-46	R5C-SAL-2	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	30 U	10 U	10 U
245-47	R4D-SAL-1	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	30 U	10 U	10 U
245-48	R4D-SAL-2	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	41	11	10 U
245-49	R7D-SAL-1	11/30/90	12/13/90	10 U	10 U	11	10 U	10 U	20 U	30 U	10 U	12
245-50	R7D-SAL-2	11/30/90	12/13/90	10 U	10 U	10 U	10 U	10 U	20 U	31	11	10 U

U Indicates analyte not detected at detection limit shown

B indicates analyte present in blank associated with that sample

NA indicates not applicable

PAH RESULTS FOR WES PLANT SAMPLES

(Concentrations in ug/Kg Dry Weight, ppb)

SURROGATE PERCENT RECOVERIES

Battelle Code	Sponsor Code	Date Ext'd	Date Anal'd	Naph-	Acen-	Phenan-	B[a]P-	Fluorene-	Chrysene-
				d8	d10	d10?Q	d12	d10	d12
BLANK 1	BLANK 1	3/18/91	4/11/91	98%	NA	NA	NA	56%	110%
245-1	R5B-SPA	3/18/91	4/11/91	36%	NA	NA	NA	35%	37%
245-2	R4B-SPA	3/18/91	4/11/91	110%	NA	NA	NA	120%	110%
245-3	R7A-SPA	3/18/91	4/11/91	79%	NA	NA	NA	100%	85%
245-4,5,6,	R2A-SPA-1,2,3	3/18/91	4/11/91	100%	NA	NA	NA	110%	110%
245-7	R1A-SPA	3/18/91	4/11/91	120%	NA	NA	NA	130%	110%
245-8,9	R2B-SPA-1,2	3/18/91	4/11/91	76%	NA	NA	NA	92%	82%
245-10	RID-SAL-1	3/18/91	4/11/91	110%	NA	NA	NA	130%	130%
245-11	RID-SAL-2	3/18/91	4/11/91	110%	NA	NA	NA	120%	130%
245-12,14	RIC-SAL-1,3	3/18/91	4/11/91	36%	NA	NA	NA	35%	37%
245-13,15	RIC-SAL-2,4	3/18/91	4/11/91	96%	NA	NA	NA	100%	91%
245-16,18	R2D-SAL-1,3	3/18/91	4/11/91	94%	NA	NA	NA	100%	100%
245-17,19	R2D-SAL-2,4	3/18/91	4/11/91	22%	NA	NA	NA	20%	23%
BLANK 2	BLANK 2	11/29/90	12/12/90	30 %	50 %	73 %	120 %		
245-20,21	R13C-TYP-1,2	11/29/90	12/12/90	72 %	83 %	78 %	120 %		
245-22,23	R13B-TYP-1,2	11/29/90	12/12/90	73 %	85 %	80 %	120 %		
245-24,25	R13D-TYP-1,2	11/29/90	12/12/90	84 %	97 %	91 %	136 %		
245-26	R13A-TYP	12/13/90	12/14/90	52 %	63 %	62 %	91 %		
245-27	R10D-TYP	11/29/90	12/12/90	78 %	90 %	81 %	130 %		
245-28	R10A-TYP	11/29/90	12/12/90	60 %	68 %	63 %	95 %		
245-29	R10C-TYP	11/29/90	12/12/90	75 %	87 %	79 %	125 %		
245-30,31	R10B-TYP-1,2	11/29/90	12/12/90	69 %	83 %	77 %	120 %		
245-32	R7B-SPA	11/29/90	12/12/90	35 %	65 %	72 %	110 %		
245-33	R1B-SPA	12/13/90	12/14/90	46 %	65 %	63 %	95 %		
245-33, DUP	R1B-SPA, DUP	12/13/90	12/14/90	76 %	87 %	78 %	120 %		
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/12/90	46 %	51 %	50 %	73 %		
245-37	R9C-SCI	11/29/90	12/13/90	28 %	65 %	77 %	125 %		
BLANK 3	BLANK 3	11/30/90	12/13/90	31 %	44 %	63 %	110 %		
245-38,39	R9D-SCI-1,2	11/30/90	12/13/90	60 %	79 %	75 %	120 %		
245-38,39 DUP	R9D-SCI-1,2	11/30/90	12/13/90	59 %	78 %	76 %	120 %		
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/13/90	45 %	60 %	59 %	87 %		
245-43	R7C-SAL-1	11/30/90	12/13/90	56 %	81 %	78 %	120 %		
245-44	R7-SAL-2	11/30/90	12/13/90	73 %	86 %	83 %	130 %		
245-45	R5C-SAL-1	11/30/90	12/13/90	69 %	85 %	82 %	120 %		
245-46	R5C-SAL-2	11/30/90	12/13/90	23 %	70 %	84 %	130 %		
245-47	R4D-SAL-1	11/30/90	12/13/90	66 %	80 %	78 %	120 %		
245-48	R4D-SAL-2	11/30/90	12/13/90	67 %	85 %	81 %	120 %		
245-49	R7D-SAL-1	11/30/90	12/13/90	62 %	87 %	86 %	140 %		
245-50	R7D-SAL-2	11/30/90	12/13/90	62 %	85 %	82 %	130 %		

U indicates analyte not detected at detection limit shown

B indicates analyte present in blank associated with that sample

NA indicates not applicable

245PAH

PAH MATRIX SPIKE PERCENT RECOVERIES c# 245

Battelle Code	Sponsor Code	Date Ext'd	Date Anal'd	% Moist.	Acenaph- thene	Acenaph- thylene	Anthra- cene	Benzo[a] Anthra- cene	Benzo[b] Fluor- anthene	Benzo[k] Fluor- anthene	Benzo[a] pyrene	Benzo- (g,h,i)- perylene
Matrix spike	RSA-SPA-1,2,3	11/28/90	12/11/90	NA	105 %	107 %	114 %	119 %	111 %	103 %	104 %	109 %
Matrix Spike Dup	RSA-SPA-1,2,3	11/28/90	12/11/90	NA	114 %	114 %	126 %	129 %	117 %	114 %	114 %	119 %
Matrix spike	R9B-SCI-1,2,3	11/29/90	12/12/90	NA	106 %	97 %	111 %	117 %	107 %	101 %	100 %	91 %
Matrix Spike Dup	R9B-SCI-1,2,3	11/29/90	12/12/90	NA	103 %	103 %	114 %	120 %	108 %	102 %	98 %	88 %
Matrix spike	R9A-SCI, 1,2,3	11/30/90	12/13/90	NA	96 %	99 %	107 %	95 %	93 %	92 %	94 %	110 %
Matrix Spike Dup	R9A-SCI, 1,2,3	11/30/90	12/13/90	NA	91 %	91 %	101 %	100 %	91 %	89 %	87 %	92 %

245PAH

PAH MATRIX SPIKE PERCENT RECOVERIES

PAH MATRIX SPIKE PERCENT RECOVERIES

Battelle Code	Sponsor Code	Date Ext'd	Date Anal'd	Chrysene	Dibenzo- (a,h)- anthracene	Fluor- anthene	Fluorene	Indeno- 1,2,3- Pyrene	2-Methyl- Naphthalene	Naph- thalene	Phenan- threne	Pyrene
Matrix spike	RSA-SPA-1,2,3	11/28/90	12/11/90	99 %	105 %	113 %	111 %	110 %	NA	29 %	128 %	109 %
Matrix Spike Dup	RSA-SPA-1,2,3	11/28/90	12/11/90	108 %	129 %	129 %	134 %	123 %	NA	114 %	144 %	124 %
Matrix spike	R9B-SCI-1,2,3	11/29/90	12/12/90	103 %	118 %	143 %	128 %	101 %	NA	45 %	128 %	139 %
Matrix Spike Dup	R9B-SCI-1,2,3	11/29/90	12/12/90	104 %	112 %	123 %	124 %	102 %	NA	65 %	128 %	116 %
Matrix spike	R9A-SCI, 1,2,3	11/30/90	12/13/90	84 %	99 %	104 %	105 %	110 %	NA	84 %	102 %	101 %
Matrix Spike Dup	R9A-SCI, 1,2,3	11/30/90	12/13/90	86 %	75 %	94 %	101 %	94 %	NA	82 %	94 %	92 %

245PAH

PAH MATRIX SPIKE PERCENT RECOVERIES

SURROGATE PERCENT RECOVERIES

Battelle Code	Sponsor Code	Date Ext'd	Date Anal'd	Naph- d8	Acen- d10	Phenan- d10?Q	B[a]P- d12
Matrix spike	RSA-SPA-1,2,3	11/28/90	12/11/90	57 %	64 %	59 %	89 %
Matrix Spike Dup	RSA-SPA-1,2,3	11/28/90	12/11/90	23 %	49 %	50 %	78 %
Matrix spike	R9B-SCI-1,2,3	11/29/90	12/12/90	13 %	35 %	53 %	100 %
Matrix Spike Dup	R9B-SCI-1,2,3	11/29/90	12/12/90	36 %	63 %	73 %	110 %
Matrix spike	R9A-SCI, 1,2,3	11/30/90	12/13/90	54 %	79 %	77 %	140 %
Matrix Spike Dup	R9A-SCI, 1,2,3	11/30/90	12/13/90	59 %	81 %	78 %	110 %

PLANT BUTYLTIN RESULTS

(Concentrations in ug/kg Dry Weight, ppm)

Battelle Code	Sponsor Code	Date Extracted	TETRABUTYL TIN	TRIBUTYL TIN	DIBUTYL TIN	MONOBUTYL TIN	SURROGATE RECOVERY TRIPENYLTIN
245-1	R5B-SPA	12/21/90	4.1 U	4.5 U	3.8 U	3.8 U	76%
245-2	R4B-SPA	12/21/90	4.2 U	4.6 U	3.9 U	3.9 U	78%
245-3	R7A-SPA	12/21/90	4.1 U	4.4 U	3.8 U	3.8 U	74%
245-4,5,6,	R2A-SPA-1,2,3	12/21/90	2.3 U	2.5 U	2.2 U	2.2 U	65%
245-7	R1A-SPA	1/15/91	4.7 U	9.2	4.3 U	19.8	89%
245-8,9	R2B-SPA-1,2	12/21/90	3.1 U	3.4 U	2.9 U	2.9 U	62%
245-10	RID-SAL-1	1/15/91	3.2 U	7.4	2.9 U	21.1	89%
245-11	RID-SAL-2	12/21/90	4.1 U	4.5 U	3.9 U	3.9 U	70%
245-12,14	RIC-SAL-1,3	12/21/90	1.6 U	1.8 U	1.5 U	1.5 U	65%
245-13,15	RIC-SAL-2,4	12/21/90	4.5 U	4.9 U	4.3 U	4.2 U	66%
245-16,18	R2D-SAL-1,3	1/15/91	3.2 U	7.4 B	2.9 U	12.5 B	89%
245-17,19	R2D-SAL-2,4	12/21/90	2.8 U	3.0 U	2.6 U	2.6 U	76%
245-20,21	R13C-TYP-1,2	12/21/90	3.2 U	3.6 U	3.0 U	3.0 U	46%
245-22,23	R13B-TYP-1,2	2/28/91	14.7 B	6.8 B	4.1 B	5.5 B	82%
245-24,25	R13D-TYP-1,2	2/28/91	18.3 B	4.3 B	2.3 B	3.3 B	82%
245-26	R13A-TYP	2/28/91	13.1 B	8.4 B	4.4 B	7.0 B	89%
245-27	R10D-TYP	2/28/91	6.3 B	2.2 B	3.7 B	14.0 B	92%
245-28	R10A-TYP	2/28/91	11.4 B	4.7 B	2.5 B	9.5 B	85%
245-29	R10C-TYP	2/28/91	11.0 B	3.9 B	2.8 B	2.2 U	83%
245-30,31	R10B-TYP-1,2	2/28/91	6.1 B	5.7 B	3.0 B	4.1 B	87%
245-32	R7B-SPA	12/21/90	3.3 U	3.6 U	3.1 U	3.1 U	51%
245-33	R1B-SPA	12/21/90	3.3 U	3.7 U	3.1 U	3.1 U	56%
245-34,35,36 REP 1	R9B-SCI-1,2,3 REP 1	1/15/91	3.2 U	6.5 B	2.9 U	2.9 U	92%
245-37	R9C-SCI	1/15/91	3.8 U	8.4 B	3.6	5.0	93%
245-38,39	R9D-SCI-1,2	1/15/91	5.1 U	14.7 B	6.7	4.6 U	84%
245-40,41,42 REP 1	R9A-SCI-1,2,3 REP 1	1/15/91	6.1	8.3 B	4.6	4.3	85%
245-43	R7C-SAL-1	1/15/91	8.2	9.6 B	5.6	6.1	89%
245-44	R7-SAL-2	1/15/91	7.4 U	18.0 B	131.8	25.1	89%
245-45	R5C-SAL-1	1/15/91	2.9 U	6.0 B	2.7 U	18.1	85%
245-46	R5C-SAL-2	1/15/91	6.9 U	16.5 B	12.9	6.3 U	89%
245-47	R4D-SAL-1	1/15/91	3.2 U	7.0 B	2.9 U	17.6	89%
245-48	R4D-SAL-2	1/15/91	7.4 U	18.1 B	11.8	17.7	92%
245-49	R7D-SAL-1	1/15/91	5.8 U	12.6 B	13.2	5.3 U	91%
245-50	R7D-SAL-2	1/15/91	6.0 U	13.3 B	16.6	5.4 U	89%
PROCEDURAL BLANK		12/21/90	5.8 U	6.4 U	6.4 U	5.3 U	38%
PROCEDURAL BLANK		1/15/91	4.2 U	7.9	3.8 U	3.8 U	85%
PROCEDURAL BLANK		2/28/91	4.7	6.5	2.6	5.9	75%

B indicates analyte detected in blank. Note, blanks for specific samples are identified by corresponding "extraction date".

U indicates analyte not detected at detection limit shown

MATRIX SPIKE RECOVERIES

(Concentrations In ug/kg Dry Weight, ppm)

Sponsor Code	TETRABUTYL TIN	TRIBUTYL TIN	DIBUTYL TIN	MONOBUTYL TIN	SURROGATE RECOVERY TRIPENYLTIN
Sample Concentration: R9B-SCI-1,2,3 REP 1	12/21/90	3.2 U 893.0	6.5 893.0	2.9 U 893.0	2.9 U 893.0
Amount Spiked:					92%
Amount Recovered:	600.1	701.1	692.2	194.1	87%
Percent Recovery :	67%	78%	77%	22%	
Sample Concentration: R1A-SPA	1/15/91	4.7 U 1087.0	9.2 1087.0	4.3 U 1087.0	19.8 1087.0
Amount Spiked:					89%
Amount Recovered:	824.5	885.4	852.3	178.6	87%
Percent Recovery :	75%	81%	78%	15%	

PLANT PCB RESULTS

(Concentrations in ug/Kg Dry Weight, ppb)

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	% Molst.	Aroclor-1016	Aroclor-1221	Aroclor-1232	Aroclor-1242	Aroclor-1248	Aroclor-1254	Aroclor-1260	SURROGATE DEC
BLANK 1	BLANK 1	11/28/90	12/3/90	NA	100 U	92 %						
245-1	R5B-SPA	11/28/90	12/3/90	86	100 U	102 %						
245-2	R4B-SPA	11/28/90	12/3/90	85	100 U	68 %						
245-3	R7A-SPA	11/28/90	12/3/90	81	100 U	65 %						
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	80	100 U	56 %						
245-7	R1A-SPA	11/28/90	12/3/90	82	100 U	61 %						
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	85	100 U	82 %						
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	NA	100 U	72 %						
245-10	RID-SAL-1	11/28/90	12/4/90	73	100 U	79 %						
245-11	RID-SAL-2	11/28/90	12/4/90	91	100 U	153 %						
245-12,14	RIC-SAL-1,3	11/28/90	12/4/90	72	100 U	82 %						
245-13,15	RIC-SAL-2,4	11/28/90	12/4/90	89	100 U	86 %						
245-16,18	R2D-SAL-1,3	11/28/90	12/4/90	73	100 U	90 %						
245-17,19	R2D-SAL-2,4	11/28/90	12/4/90	85	100 U	144 %						
BLANK 2	BLANK 2	11/29/90	12/4/90	NA	100 U	78 %						
245-20,21	R13C-TYP-1,2	11/29/90	12/4/90	86	100 U	68 %						
245-22,23	R13B-TYP-1,2	11/29/90	12/4/90	88	100 U	99 %						
245-24,25	R13D-TYP-1,2	11/29/90	12/4/90	81	100 U	46 %						
245-26	R13A-TYP	11/29/90	12/4/90	90	100 U	67 %						
245-17	R10D-TYP	11/29/90	12/4/90	88	100 U	99 %						
245-28	R10A-TYP	11/29/90	12/4/90	85	100 U	51 %						
245-29	R10C-TYP	11/29/90	12/4/90	81	100 U	75 %						
245-30,31	R10B-TYP-1,2	11/29/90	12/4/90	82	100 U	67 %						
245-32	R7B-SPA	11/29/90	12/4/90	80	100 U	82 %						
245-33	R1B-SPA	11/29/90	12/5/90	86	100 U	65 %						
245-33	R1B-SPA	11/29/90	12/5/90	NA	100 U	77 %						
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	77	100 U	39 %						
245-37	R9C-SCI	11/29/90	12/5/90	80	100 U	40 %						
BLANK 3	BLANK 3	11/30/90	12/5/90	NA	100 U	63 %						
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	82	100 U	66 %						
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	NA	100 U	78 %						
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	75	100 U	66 %						
245-43	R7C-SAL-1	11/30/90	12/5/90	78	100 U	85 %						
245-44	R7-SAL-2	11/30/90	12/5/90	93	100 U	83 %						
245-45	R5C-SAL-2	11/30/90	12/5/90	74	100 U	83 %						
245-46	R5C-SAL-1	11/30/90	12/5/90	89	100 U	95 %						
245-47	R4D-SAL-1	11/30/90	12/5/90	76	100 U	77 %						
245-48	R4D-SAL-2	11/30/90	12/5/90	88	100 U	82 %						
245-49	R7D-SAL-1	11/30/90	12/6/90	80	100 U	70 %						
245-50	R7D-SAL-2	11/30/90	12/6/90	89	100 U	86 %						

U indicates analyte not detected above detection limit shown.

NA indicates not applicable

PESTICIDE RESULTS

(Concentrations in ug/Kg Dry Weight, ppb)

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	% Molst.	Aldrin	Alpha-BHC	Beta-BHC	Delta-BHC	Gamma-BHC	Chlordane	4,4'-DDD
BLANK 1	BLANK 1	11/28/90	12/3/90	NA	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-1	R5B-SPA	11/28/90	12/3/90	86	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-2	R4B-SPA	11/28/90	12/3/90	85	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-3	R7A-SPA	11/28/90	12/3/90	81	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	80	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-7	R1A-SPA	11/28/90	12/3/90	82	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	85	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	NA	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-10	R1D-SAL-1	11/28/90	12/4/90	73	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-11	R1D-SAL-2	11/28/90	12/4/90	91	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-12,14	R1C-SAL-1,3	11/28/90	12/4/90	72	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-13,15	R1C-SAL-2,4	11/28/90	12/4/90	89	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-16,18	R2D-SAL-1,3	11/28/90	12/4/90	73	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-17,19	R2D-SAL-2,4	11/28/90	12/4/90	85	20 U	20 U	20 U	20 U	20 U	30 U	20 U
BLANK 2	BLANK 2	11/29/90	12/4/90	NA	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-20,21	R13C-TYP-1,2	11/29/90	12/4/90	86	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-22,23	R13B-TYP-1,2	11/29/90	12/4/90	88	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-24,25	R13D-TYP-1,2	11/29/90	12/4/90	81	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-26	R13A-TYP	11/29/90	12/4/90	90	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-17	R10D-TYP	11/29/90	12/4/90	88	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-28	R10A-TYP	11/29/90	12/4/90	85	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-29	R10C-TYP	11/29/90	12/4/90	81	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-30,31	R10B-TYP-1,2	11/29/90	12/4/90	82	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-32	R7B-SPA	11/29/90	12/4/90	80	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-33	R1B-SPA	11/29/90	12/5/90	86	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-33	R1B-SPA	11/29/90	12/5/90	NA	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	77	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-37	R9C-SCI	11/29/90	12/5/90	80	20 U	20 U	20 U	20 U	20 U	30 U	20 U
BLANK 3	BLANK 3	11/30/90	12/5/90	NA	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	82	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	NA	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	75	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-43	R7C-SAL-1	11/30/90	12/5/90	78	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-44	R7-SAL-2	11/30/90	12/5/90	93	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-45	R5C-SAL-2	11/30/90	12/5/90	74	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-46	R5C-SAL-1	11/30/90	12/5/90	89	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-47	R4D-SAL-1	11/30/90	12/5/90	76	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-48	R4D-SAL-2	11/30/90	12/5/90	88	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-49	R7D-SAL-1	11/30/90	12/6/90	80	20 U	20 U	20 U	20 U	20 U	30 U	20 U
245-50	R7D-SAL-2	11/30/90	12/6/90	89	20 U	20 U	20 U	20 U	20 U	30 U	20 U

U indicates analyte not detected above detection limit shown.

NA indicates not applicable

PESTICIDE RESULTS

(Concentrations in ug/Kg Dry Weight, ppb)

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	4,4'-DDE	4,4'-DDT	Dieldrin	Endosulfan I	Endosulfan II	Sulfate	Endrin
BLANK 1	BLANK 1	11/28/90	12/3/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-1	R5B-SPA	11/28/90	12/3/90	20 U	20 U	20 U	20 U	20 U	20 U	670 U
245-2	R4B-SPA	11/28/90	12/3/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-3	R7A-SPA	11/28/90	12/3/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-7	R1A-SPA	11/28/90	12/3/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-10	RID-SAL-1	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-11	RID-SAL-2	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-12,14	RIC-SAL-1,3	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-13,15	RIC-SAL-2,4	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-16,18	R2D-SAL-1,3	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-17,19	R2D-SAL-2,4	11/28/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
BLANK 2	BLANK 2	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-20,21	R13C-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-22,23	R13B-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-24,25	R13D-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-26	R13A-TYP	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-17	R10D-TYP	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-28	R10A-TYP	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-29	R10C-TYP	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-30,31	R10B-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-32	R7B-SPA	11/29/90	12/4/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-33	R1B-SPA	11/29/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-33	R1B-SPA	11/29/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-37	R9C-SCI	11/29/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
BLANK 3	BLANK 3	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-43	R7C-SAL-1	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-44	R7-SAL-2	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-45	R5C-SAL-2	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-46	R5C-SAL-1	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-47	R4D-SAL-1	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-48	R4D-SAL-2	11/30/90	12/5/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-49	R7D-SAL-1	11/30/90	12/6/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U
245-50	R7D-SAL-2	11/30/90	12/6/90	20 U	20 U	20 U	20 U	20 U	20 U	20 U

U indicates analyte not detected above detection limit show

NA indicates not applicable

PESTICIDE RESULTS

(Concentrations in ug/Kg Dry Weight, ppb)

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	Endrin Aldehyde	Heptachlor	Heptachlor Epoxide	Methoxy-chlor	Toxaphene	SURROGATE DBC
BLANK 1	BLANK 1	11/28/90	12/3/90	20 U	20 U	20 U	30 U	200 U	92 %
245-1	R5B-SPA	11/28/90	12/3/90	20 U	20 U	20 U	30 U	200 U	102 %
245-2	R4B-SPA	11/28/90	12/3/90	20 U	20 U	20 U	30 U	200 U	68 %
245-3	R7A-SPA	11/28/90	12/3/90	20 U	20 U	20 U	30 U	200 U	65 %
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	20 U	20 U	20 U	30 U	200 U	56 %
245-7	R1A-SPA	11/28/90	12/3/90	20 U	20 U	20 U	30 U	200 U	61 %
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	82 %
245-8,9	R2B-SPA-1,2	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	72 %
245-10	RID-SAL-1	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	79 %
245-11	RID-SAL-2	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	153 %
245-12,14	RIC-SAL-1,3	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	82 %
245-13,15	RIC-SAL-2,4	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	86 %
245-16,18	R2D-SAL-1,3	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	90 %
245-17,19	R2D-SAL-2,4	11/28/90	12/4/90	20 U	20 U	20 U	30 U	200 U	144 %
BLANK 2	BLANK 2	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	78 %
245-20,21	R13C-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	68 %
245-22,23	R13B-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	99 %
245-24,25	R13D-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	46 %
245-26	R13A-TYP	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	67 %
245-17	R10D-TYP	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	99 %
245-28	R10A-TYP	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	51 %
245-29	R10C-TYP	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	75 %
245-30,31	R10B-TYP-1,2	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	67 %
245-32	R7B-SPA	11/29/90	12/4/90	20 U	20 U	20 U	30 U	200 U	82 %
245-33	R1B SPA	11/29/90	12/5/90	20 U	20 U	20 U	30 U	200 U	65 %
245-33	R1B-SPA	11/29/90	12/5/90	20 U	20 U	20 U	30 U	200 U	77 %
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	20 U	20 U	20 U	30 U	200 U	39 %
245-37	R9C-SCI	11/29/90	12/5/90	20 U	20 U	20 U	30 U	200 U	40 %
BLANK 3	BLANK 3	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	63 %
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	66 %
245-38,39	R9D-SCI-1,2	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	78 %
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	66 %
245-43	R7C-SAL-1	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	85 %
245-44	R7-SAL-2	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	83 %
245-45	R5C-SAL-2	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	83 %
245-46	R5C-SAL-1	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	95 %
245-47	R4D-SAL-1	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	77 %
245-48	R4D-SAL-2	11/30/90	12/5/90	20 U	20 U	20 U	30 U	200 U	82 %
245-49	R7D-SAL-1	11/30/90	12/6/90	20 U	20 U	20 U	30 U	200 U	70 %
245-50	R7D-SAL-2	11/30/90	12/6/90	20 U	20 U	20 U	30 U	200 U	86 %

U indicates analyte not detected above detection limit show

NA indicates not applicable

PESTICIDE MATRIX SPIKE RECOVERIES

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	% Molst.	Aldrin	Alpha-BHC	Beta-BHC	Delta-BHC	Gamma-BHC	Chlordane	4,4'-DDD
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	NA	71 %	NA	NA	NA	NA	NA	NA
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	NA	88 %	NA	NA	NA	NA	NA	NA
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	NA	79 %	NA	NA	NA	NA	NA	NA
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	NA	102 %	NA	NA	NA	NA	NA	NA
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	NA	105 %	NA	NA	NA	NA	NA	NA
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	NA	99 %	NA	NA	NA	NA	NA	NA

PESTICIDE MATRIX SPIKE RECOVERIES

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	4,4'-DDE	4,4'-DDT	Dieldrin	Endo-sulfan I	Endo-sulfan II	Sulfate	Endrin
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	NA	NA	48 %	NA	NA	NA	NA
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90	NA	NA	58 %	NA	NA	NA	NA
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	NA	NA	47 %	NA	NA	NA	NA
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90	NA	NA	64 %	NA	NA	NA	NA
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	NA	NA	84 %	NA	NA	NA	NA
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90	NA	NA	62 %	NA	NA	NA	NA

PESTICIDE MATRIX SPIKE RECOVERIES

Client Sample ID	Sponsor Code	Date Ext'd	Date Anal'd	Endrin Aldehyde	Heptachlor	Heptachlor Epoxide	Methoxy-chlor	Toxaphene	DBC
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90		NA	NA	NA	200 NA	NA
245-4,5,6	R2A-SPA-1,2,3	11/28/90	12/3/90		NA	NA	NA	200 NA	67 %
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90		NA	NA	NA	200 NA	46 %
245-34,35,36	R9B-SCI-1,2,3	11/29/90	12/5/90		NA	NA	NA	200 NA	70 %
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90		NA	NA	NA	200 NA	66 %
245-40,41,42	R9A-SCI-1,2,3	11/30/90	12/5/90		NA	NA	NA	200 NA	65 %

BUTYLTINS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(Concentrations In ug/kg dry weight)

MSL Code	Sponsor Code	Tripentyl % Surrogate	Pentylbutyl % Internal	Tetra	Tributyl	Dibutyl	Monobutyl
277- 1-R	SED09-CB	75.47	123.0	1.9 U	3.2	3.3	5.8
277- 2	SED07-CM	98.55	131.6	2.9	2.0	9.6	2.1
277- 3-R	SED01-MR	74.94	133.6	1.3 U	2.3	1.4 U	1.3 U
277- 4-R	SED05-CM	73.13	137.3	1.2 U	3.1	1.7	1.2 U
277- 5-R	SED10-CB	75.68	134.1	1.5 U	3.6	1.6 U	4.7
277- 6-R	SED13-CF	67.43	142.8	0.9 U	1.8	0.9 U	0.9 U
277- 7-R	SED08-CM	75.64	122.7	2.0	2.3	1.4 U	1.3 U
277- 8-R	SED14-BR	73.00	140.4	1.3 U	3.5	1.8	2.4
277- 9-R	SED11-CB	75.35	142.5	0.9 U	33.4	0.9 U	0.9 U
277- 10-R	SED04-CM	86.65	127.5	1.4 U	3.1	2.0	2.3
277- 11	SED02-CM	148.81	142.4	0.5	2.6	3.6	17.0
277- 12-R	SED03-CM	83.58	133.6	3.0	2.6	1.4 U	2.9
277- 13-R	SEDWR09-CM	83.68	135.9	0.8 U	1.3	0.8 U	0.7 U
277- 14	08A-SAL	128.66	145.5	2.4	4.5	2.2	53.5
277- 15	14C-SAL	100.94	155.1	2.4	4.8	2.2	35.1
277- 16	08C-SAL	98.79	134.9	2.0	3.5	11.1	24.6
277- 17	04C-SAL	119.99	140.7	3.2	6.0	19.0	64.3
277- 18	03C-SAL	91.91	126.7	2.2	3.1	6.6	15.6
277- BLK-2		83.66	136.6	3.3	4.8	12.1	24.7
277- 19R	11C-SCI	80.40	144.4	2.2 U	4.1	2.1 U	4.4
277- 20-R	11A-SCI	76.59	118.9	4.1 U	5.6	5.6	3.7 U
277- 21	11B-SCI	129.64	136.8	5.5	5.2	2.6	9.5
277- 22-R	03B-SPA	96.66	111.0	3.6 U	8.3	3.7	5.1
277- 23	04A-SPA	152.9	64.4	2.7	5.2	2.5	NA
277- 24	14A-SCI	178.00	44.0	1.2	2.2	1.1	NA
277- 25-R	14D-SAL	81.97	117.4	3.1 U	4.4	3.0 U	5.6
277- 26-R	02C-SAL	88.17	113.4	9.7	6.5	3.5 U	7.1
277- 27	05D-SAL	74.01	127.2	54.7	35.8	2.3	5.3
277- 28	03D-SAL	74.05	126.2	3.3	4.8	4.4	7.1
227- BLK-4		67.67	133.7	2.2 U	3.9	2.1 U	10.2
277- 29	08D-SAL	61.68	122.9	2.3 U	4.0	2.8	2.1
277- 30	08B-SAL	63.86	136.4	3.1 U	5.3	3.1 U	2.9 U
277- 31-R	08B-SAL	90.86	115.6	3.6 U	5.6	3.6 U	6.4
277- BLANK		75.31	123.8	3.8 U	5.3	3.7 U	8.7
277- 32	03A-SPA	62.55	121.0	2.1 U	2.9	2.1 U	1.9 U
277- 33	05A-SPA	70.21	125.6	2.2 U	5.2	2.2 U	2.0 U
277- 34	R08-SN01	85.92	112.1	1.6 U	3.3	5.0	1.5 U

BUTYLTINS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(Concentrations in ug/kg dry weight)

MSL Code	Sponsor Code	Tripentyl % Surrogate	Pentylbutyl % Internal	Tetra	Tributyl	Dibutyl	Monobutyl
277- 34 DUP	R08-SN01	80.00	128.8	1.4 U	3.5	4.2	1.7
277- 35	R08-SN02	80.15	134.2	0.6 U	1.4	0.9	1.6
277- 36	R13-CBR1	71.18	132.5	14.6	40.7	30.1	11.8
277- 37	R01-MOR1	69.60	122.2	3.9 U	34.9	9.3	7.8
277- 38	R01-MOR2	131.20	52.3	5.0 U	38.3	5.0 U	4.6 U

U Indicates not detected at detection limit shown

MATRIX SPIKE RESULTS

277-1-C SPIKE Percent Recovery	69.36	144.4	168.0 35%	219.6 46%	243.0 51%	37.6 7.7%
277-5-C SPIKE Percent Recovery	67.29	142.5	149.6 34%	184.6 41%	31.5 7%	123.4 28%
277-8-C SPIKE Percent Recovery	82.71	125.4	138.4 35%	203.7 51%	260.7 66%	44.8 11%
227-BLANK SPIKE Percent Recovery	76.42	177.2	294.4 58%	301.6 59%	327.5 65%	221.5 43%
277-36 SPIKE Percent Recovery	73.86	128.1	1004.6 60%	1153.7 69%	841.8 50%	63.5 4%

METALS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(concentrations in mg/kg dry weight)

MSL Code	Rep	Sponsor ID	Rep	Ag	As	Cd	(a) Cr	Cu	Hg	NI	(a) Pb	(b) Se	Zn
				AA	XRF	AA	ANXRF	XRF	CVAA	XRF	ANXRF	AA/XRF	XRF
SEDIMENT													
277-1	REP 1	SED09-CB	REP 1	0.448	19.3	0.28	183.0	68.5	0.383	107.7	85.6	0.41	142.2
277-1	REP 2	SED09-CB	REP 2	0.446	20.7	0.28	168.0	72.4	0.394	106.6	84.6	0.39	145.5
277-2		SED07-CM		0.355	10.6	0.33	195.0	67.5	0.469	119.8	33.8	0.33	157.5
277-3		SED01-MR		1.418	23.7	0.33	174.0	71.6	0.515	102.0	36.3	0.33	137.2
277-4		SED05-CM		0.660	14.4	0.26	179.0	67.6	0.419	125.9	34.1	0.25	158.4
277-5		SED10-CB		0.359	17.2	0.56	126.0	67.9	0.321	93.3	47.8	0.91	135.0
277-6		SED13-CF		0.234	5.36	0.55	110.0	24.2	0.059	32.2	14.0	0.14 U	161.7
277-7		SED08-CM		0.023	5.29	0.41	224.0	35.9	0.074	72.2	20.9	0.14 U	88.5
277-8		SED14-BR		0.206	16.9	0.36	193.0	77.3	0.362	122.1	32.5	0.25	164.7
277-9		SED11-CB		0.350	15.3	0.22	181.0	50.3	0.283	83.3	13.7	0.16	89.8
277-10		SED04-CM		0.143	13.4	0.31	214.0	72.6	0.439	135.5	35.7	0.17	160.1
277-11		SED02-CM		0.372	18.5	0.32	219.0	90.6	0.469	125.4	36.8	0.33	158.9
277-12		SED03-CM		0.479	18.2	0.41	179.0	70.1	0.166	145.2	33.0	0.42	166.1
277-13		SEDWR09-CM		0.194	9.9	0.22	256.0	28.6	0.164	72.7	13.2	0.17	77.8
PLANTS													
277-14	REP 1	08A-SAL		0.003 U	1.3 U	0.13	0.4	9.7	0.024	1.7 U	0.23	1.10 U	27.3
277-14	REP 2	14C-SAL		0.003 U	0.96 U	0.13	0.4	9.2	0.023	2.33	0.34	1.10 U	29.2
277-15	REP 1	14C-SAL		0.003	0.92 U	0.17	3.6	10.1	0.034	3.78	0.99	0.70 U	30.8
277-15	REP 2	14C-SAL		-	-	-	3.9 *	-	-	-	1.27 *	-	-
277-16		08C-SAL		0.007	0.85 U	0.21	0.4	8.7	0.030	1.48	0.92	0.66 U	36.0
277-17		04C-SAL		0.007	1.0 U	0.29	5.9	19.1	0.038	6.29	1.42	0.79 U	45.7
277-18		03C-SAL		0.003 U	1.0	0.05	1.8	8.0	0.016	3.31	0.66	0.65 U	26.6
277-19		11C-SCI		0.003 U	0.79 U	0.16	0.7	15.3	0.018	4.47	0.49	0.56 U	88.7
277-20		11A-SCI		0.003 U	0.87	0.17	2.7	31.1	0.050	6.70	0.87	0.62 U	89.9
277-21		11B-SCI		0.003 U	0.89 U	0.24	4.0	17.4	0.044	9.39	1.03	0.65 U	133.0
277-22		03B-SPA		0.135	1.04	0.12	7.7	13.9	0.025	9.29	1.84	0.64 U	84.9
277-23		04A-SPA		0.107	1.82	0.07	2.5	8.9	0.014	2.05	0.60	0.68 U	60.9
277-24		14A-SCI		0.034	0.79 U	0.08	3.3	7.7	0.038	3.47	1.18	0.58 U	48.4
277-25		14D-SAL		0.009	0.95 U	0.07	1.7	11.4	0.019	1.85	0.71	0.71 U	29.8
277-26		02C-SAL		0.003	0.91 U	0.16	1.8	10.8	0.019	2.47	0.61	2.20 U	40.0
277-27		05D-SAL		0.014	0.88 U	0.06	4.1	11.5	0.018	4.49	0.86	0.66 U	44.3
277-28		03D-SAL		0.009	0.86 U	0.08	2.6	12.0	0.021	5.27	0.93	0.64 U	25.8
277-29		08D-SAL		0.003	0.83 U	0.15	0.4	8.9	0.025	0.93 U	0.38	0.65 U	36.3
277-30		08B-SAL		0.003	0.99 U	0.10	0.5	8.8	0.016	1.47	0.49	0.77 U	57.4
277-31		11D-SCI		0.005	0.84 U	0.13	1.9	13.8	0.028	5.81	0.76	0.61 U	59.3
277-32		03A-SPA		0.217	1.27	0.06	7.2	13.7	0.022	8.76	1.39	0.63 U	98.0
277-33		05A-SPA		0.165	0.99	0.08	8.5	11.4	0.027	9.1	2.04	0.65 U	65.5

METALS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(concentrations in mg/kg dry weight)

MSL Code	Rep	Sponsor ID	Rep	Ag	As	Cd	(a) Cr	Cu	Hg	NI	(a) Pb	(b) Se	Zn
				AA	XRF	AA	AA/XRF	XRF	CVAA	XRF	AA/XRF	AA/XRF	XRF

TISSUES

277-34	R08-SN01	0.347	11.62	1.03	2.2	93.6	0.180	10.2	1.15	1.33	401.0
277-34	R08-SN01	0.360	9.22	1.03	2.1	74.3	0.172	8.5	1.43	1.04	309.0
277-35	R08-SN02	0.121	2.5	0.34	1.2	23.5	0.055	4.5	0.82	1.47	131.4
277-36	R13-CBR1	1.03	10.79	3.34	4.3	164.1	0.469	5.78	1.89	3.98	273.0
277-37	R01-MOR1	0.819	8.76	3.53	4.0	23.1	0.398	7.74	1.71	4.19	71.7
277-38	R01-MOR2	0.914	8.93	3.45	3.3	20.5	0.304	5.33	1.39	3.52	71.1
Blank		0.019	N/A	0.01 U	0.8	N/A	0.001 U	N/A	N/A	0.14 U	N/A
Blank		0.007	N/A	0.01 U	0.8	N/A	0.001 U	N/A	0.17 U	0.14 U	N/A

U Indicates not detected at detection limit shown

N/A indicates not applicable

PESTICIDES IN SED., PLANT & TISSUE & TISSUE

Sponsor: SIMMER (McGUFFIE)

		(CONCENTRATIONS IN UG/KG WET WEIGHT) 3HT											
		% Moist.	% Moist.	Aldrin	Alpha- BHC	Beta- BHC	Delta- BHC	Gamma- BHC	Chlor- dane	4,4- DDD	4,4- DDDE	4,4- DDDT	Dieldrin
SEDIMENT													
SEDIMENT	METHOD BLANK	N/A	N/A	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 1	SED09-CB	60	60	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.6	3.0 U	3.0 U
277- 2	SED07-CM	55	55	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 3	SED01-MR	47	47	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 4	SED05-CM	45	45	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 5	SED10-CB	72	72	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U
277- 6	SED13-CF	33	33	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 7	SED08-CM	38	38	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 8	SED14-BR	34	34	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 9	SED11-CB	32	32	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 10	SED04-CM	54	54	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 11	SED02-CM	49	49	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 12	SED03-CM	57	57	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
277- 13	SEDWR09-CM	19	19	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U
PLANTS													
PLANT	METHOD BLANK	N/A	N/A	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 14	08A-SAL	89	89	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 15	14C-SAL	89	89	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 16	08C-SAL	86	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 17	04C-SAL	90	90	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	3.0 U
277- 18	03C-SAL	84	84	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 19	11C-SCI	90	90	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 20	11A-SCI	87	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 21	11B-SCI	86	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 22	03B-SPA	87	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 23	04A-SPA	87	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 24	14A-SCI	74	74	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 25	14D-SAL	81	81	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 26	02C-SAL	84	84	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 27	05D-SAL	87	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 28	03D-SAL	85	85	2.0 U	2.3	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 29	08D-SAL	87	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 30	08B-SAL	86	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 31	11D-SCI	88	88	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	3.0 U
277- 32	03A-SPA	86	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U
277- 33	05A-SPA	84	84	2.0 U	3.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U

PESTICIDES IN SED., PLANT & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN ug/kg WET WEIGHT)

	% Moist.	Endo- sultan I	Endo- sultan I	Endosulfan Sulfate	Endrin	Endrin Aldehyde	Hepta- chlor	Heptachlor Epoxide	Methoxy- chlor	Toxa- phene	SURROGATE DBC	PERCENT RECOVERY
SEDIMENT												
SEDIMENT METHOD BLANK	N/A	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	37
277- 1 SED09-CB	60	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	57
277- 2 SED07-CM	55	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	63
277- 3 SED01-MR	47	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	73
277- 4 SED05-CM	45	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	9
277- 5 SED10-CB	72	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	5.0 U	250 U	32
277- 6 SED13-CF	33	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	38
277- 7 SED08-CM	38	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	59
277- 8 SED14-BR	34	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	87
277- 9 SED11-CB	32	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	97
277- 10 SED04-CM	54	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	121
277- 11 SED02-CM	49	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	83
277- 12 SED03-CM	57	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	87
277- 13 SEDWR09-CM	19	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	3.0 U	200 U	92
PLANTS												
PLANT METHOD BLANK	N/A	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	63
277- 14 08A-SAL	89	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	71
277- 15 14C SAL	89	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	132
277- 16 08C-SAL	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	89
277- 17 04C-SAL	90	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	122
277- 18 03C-SAL	84	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	11
277- 19 11C-SCI	90	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	67
277- 20 11A-SCI	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	106
277- 21 11B-SCI	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	113
277- 22 03B-SPA	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	121
277- 23 04A-SPA	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	48
277- 24 14A-SCI	74	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	32
277- 25 14D-SAL	81	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	103
277- 26 02C-SAL	84	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	87
277- 27 05D-SAL	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	122
277- 28 03D-SAL	85	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	176
277- 29 08D-SAL	87	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	138
277- 30 08B-SAL	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	154
277- 31 11D-SCI	88	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	79
277- 32 03A-SPA	86	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	87
277- 33 05A-SPA	84	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	2.0 U	100 U	75

PESTICIDES IN SED., PLANT & TISSUE & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

GHT

	% Moist.	% Moist.	Aldrin	Alpha- BHC	Beta- BHC	Delta- BHC	Gamma- BHC	Chlor- dane	4,4- DDD	4,4- DDOE	4,4- DDDT	Dieldrin
--	-------------	-------------	--------	---------------	--------------	---------------	---------------	----------------	-------------	--------------	--------------	----------

TISSUE

TISSUE METHOD BLANK		N/A	N/A	10 U	10 U							
277- 34 R08-SN01		66	66	10 U	10 U							
277- 34 R08-SN01		66	66	10 U	10 U							
277- 35 R08-SN02		35	35	10 U	10 U							
277- 36 R13-CBR1		92	92	10 U	10 U	12 U	24 U	10 U	10 U	10 U	115 U	30 U
277- 37 R01-MOR1		85	85	10 U	10 U							
277- 38 R01-MOR2		88	88	10 U	10 U							

U indicates not detected at detection limits shown limits shown

PESTICIDES IN SED., PLANT & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

	% Moist.	Endo- sulfan I	Endo- sulfan I	Endosulfan Sulfate	Endrin	Endrin Aldehyde	Hepta- chlor	Heptachlor Epoxide	Methoxy- chlor	Toxa- phene	PERCENT RECOVERY SURROGATE DBC
--	-------------	-------------------	-------------------	-----------------------	--------	--------------------	-----------------	-----------------------	-------------------	----------------	---

TISSUE

TISSUE METHOD BLANK	N/A	10 U	67								
277- 34 R08-SN01	66	10 U	111								
277- 34 R08-SN01	66	10 U	157								
277- 35 R08-SN02	35	10 U	110								
277- 36 R13-CBR1	92	10 U	10 U	10 U	18	10 U	42	10 U	10 U	10 U	119
277- 37 R01-MOR1	85	10 U	130								
277- 38 R01-MOR2	88	10 U	70								

U indicates not detected at detection limits shown

PCBs IN SED., PLANT & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

	% Moist.	Aroclor- 1016	Aroclor- 1221	Aroclor- 1232	Aroclor- 1242	Aroclor- 1248	Aroclor- 1254	Aroclor- 1260	SURROGATE DBC	PERCENT RECOVERY
SEDIMENT										
SEDIMENT METHOD BLANK	N/A	30 U	37							
277- 1 SED09-CB	60	30 U	57							
277- 2 SED07-CM	55	30 U	63							
277- 3 SED01-MR	47	30 U	150	30 U	73					
277- 4 SED05-CM	45	30 U	9							
277- 5 SED10-CB	72	50 U	32							
277- 6 SED13-CF	33	30 U	38							
277- 7 SED08-CM	38	30 U	59							
277- 8 SED14-BR	34	30 U	87							
277- 9 SED11-CB	32	30 U	97							
277- 10 SED04-CM	54	30 U	120	30 U	121					
277- 11 SED02-CM	49	30 U	83	30 U	83					
277- 12 SED03-CM	57	30 U	210	30 U	87					
277- 13 SEDWR09-CM	19	30 U	75	30 U	92					
PLANTS										
PLANT METHOD BLANK	N/A	20 U	63							
277- 14 08A-SAL	89	20 U	71							
277- 15 14C-SAL	89	20 U	132							
277- 16 08C-SAL	86	20 U	89							
277- 17 04C-SAL	90	20 U	122							
277- 18 03C-SAL	84	20 U	11							
277- 19 11C-SCI	90	20 U	67							
277- 20 11A-SCI	87	20 U	106							
277- 21 11B-SCI	86	20 U	113							
277- 22 03B-SPA	87	20 U	121							
277- 23 04A-SPA	87	20 U	48							
277- 24 14A-SCI	74	20 U	32							
277- 25 14D-SAL	81	20 U	103							
277- 26 02C-SAL	84	20 U	87							
277- 27 05D-SAL	87	20 U	122							
277- 28 03D-SAL	85	20 U	176							
277- 29 08D-SAL	87	20 U	138							
277- 30 08B-SAL	86	20 U	154							
277- 31 11D-SCI	88	20 U	79							
277- 32 03A-SPA	86	20 U	87							
277- 33 05A-SPA	84	20 U	75							

PCBs IN SED., PLANT & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

% Moist.	Aroclor- 1016	Aroclor- 1221	Aroclor- 1232	Aroclor- 1242	Aroclor- 1248	Aroclor- 1254	Aroclor- 1260	PERCENT RECOVERY SURROGATE DBC
-------------	------------------	------------------	------------------	------------------	------------------	------------------	------------------	---

TISSUE

TISSUE METHOD BLANK	N/A	100 U	67						
277- 34 R08-SN01	66	100 U	111						
277- 34 R08-SN01	66	100 U	157						
277- 35 R08-SN02	35	100 U	110						
277- 36 R13-CBR1	92	100 U	119						
277- 37 R01-MOR1	85	100 U	130						
277- 38 R01-MOR2	88	100 U	70						

U indicates not detected at detection limits shown

PCB and Pesticide Matrix Spike Recoveries

Battelle Code	Sponsor Codes	Aldrin	Dieldrin	Aroclor- 1254	Surrogate DBC
277- 12	SED03-CM	61%	113%	79%	
277- 12	SED03-CM	62%	88%	ND	
277- 33	05A-SPA	57%	59%	189%	165%
277- 33	05A-SPA	58%	50%	144%	129%
277- 36	R13-CBR1	107%	80%	83%	116%
277- 36	R13-CBR1	83%	83%	100%	103%

PCBs IN SED., PLANT & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

% Moist.	Aroclor- 1016	Aroclor- 1221	Aroclor- 1232	Aroclor- 1242	Aroclor- 1248	Aroclor- 1254	Aroclor- 1260	PERCENT RECOVERY SURROGATE DBC
-------------	------------------	------------------	------------------	------------------	------------------	------------------	------------------	---

TISSUE

TISSUE METHOD BLANK	N/A	100 U	67						
277- 34 R08-SN01	66	100 U	111						
277- 34 R08-SN01	66	100 U	157						
277- 35 R08-SN02	35	100 U	110						
277- 36 R13-CBR1	92	100 U	119						
277- 37 R01-MOR1	85	100 U	130						
277- 38 R01-MOR2	88	100 U	70						

U indicates not detected at detection limits shown

PAHS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

Battelle Code	Sponsor Code	% Moist.	Dibenzo- (a,h)- anthracene	Fluor- anthene	Fluorene	Indeno- 1,2,3- Pyrene	2-Methyl- Naphthalene	Naph- thalene	Phenan- threne	Pyrene
SEDIMENT										
277- 1	SED09-CB	60%	14	74	10 U	51	30	63	20	89
277- 2	SED07-CM	55%	15	120	10 U	87	12	26	45	160
277- 3	SED01-MR	47%	19	190	10 U	99	30	61	94	240
277- 4	SED05-CM	45%	10 U	10 U	10 U	10 U	15	34	10 U	10 U
277- 5	SED10-CB	72%	30	260	10 U	100	48	97	76	240
277- 6	SED13-CF	33%	10 U	49	10 U	14	17	35	20	46
277- 7	SED08-CM	38%	10 U	18	10 U	11	10 U	56	10 U	20
277- 8	SED14-BR	34%	10 U	28	10 U	17	30	59	13	33
277- 9	SED11-CB	32%	10 U	10 U	10 U	10 U	10 U	20 U	10 U	10 U
277- 10	SED04-CM	54%	11	110	10 U	77	25	50	42	140
277- 11	SED02-CM	49%	10	94	10 U	59	27	53	36	120
277- 12	SED03-CM	57%	10 U	54	10 U	43	35	64	25	72
277- 13	SEDWR09-CM	19%	69	490	72	320	20	37	460	630
PLANTS										
277- 14	08A-SAL	89%	10 U	10 U	10 U	10 U	30	90	16	10 U
277- 15	14C-SAL	89%	10 U	10 U	10 U	10 U	32	97	17	10 U
277- 16	08C-SAL	86%	10 U	10 U	10 U	10 U	25	68	20	10 U
277- 17	04C-SAL	90%	10 U	10 U	10 U	10 U	25	73	13	10 U
277- 18	03C-SAL	84%	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
277- 19	11C-SCI	90%	10 U	10 U	10 U	10 U	20 U	50 U	10	10 U
277- 20	11A-SCI	87%	10 U	10 U	10 U	10 U	24	60	18	10 U
277- 21	11B-SCI	86%	10 U	10 U	10 U	10 U	27	76	18	10 U
277- 22	03B-SPA	87%	10 U	10 U	10 U	10 U	29	88	14	10 U
277- 23	04A-SPA	87%	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
277- 24	14A-SCI	74%	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
277- 25	14D-SAL	81%	10 U	10 U	10 U	10 U	24	61	16	10 U
277- 26	02C-SAL	84%	10 U	10 U	10 U	10 U	24	59	22	10 U
277- 27	05D-SAL	87%	10 U	10 U	10 U	10 U	37	120	17	10 U
277- 28	03D-SAL	85%	10 U	10 U	10 U	10 U	28	83	15	10 U
277- 29	08D-SAL	87%	10 U	10 U	10 U	10 U	28	89	15	10 U
277- 30	08B-SAL	86%	10 U	10 U	10 U	10 U	20	60	12	10 U
277- 31	11D-SCI	88%	10 U	10 U	10 U	10 U	25	62	14	10 U
277- 32	03A-SPA	86%	10 U	10 U	10 U	10 U	24	68	14	10 U
277- 33	05A-SPA	84%	10 U	10 U	10 U	10 U	25	68	17	10 U

PAHS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

Battelle Code	Sponsor Code	% Moist.	Acenaph- thene	Acenaph- thylenne	Anthra- cene	Benzo[a] Anthra- cene	Benzo[b] Fluor- anthene	Benzo[k] Fluor- anthene	Benzo[a] pyrene	Benzo- (g,h,i)- perylene	Chrysene
SEDIMENT											
277- 1	SED09-CB	60%		10 U	10 U	17	56	83	67	62	65
277- 2	SED07-CM	55%		10 U	10 U	16	67	82	72	86	100
277- 3	SED01-MR	47%		12	15	38	100	96	82	130	110
277- 4	SED05-CM	45%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	100
277- 5	SED10-CB	72%		19	120	97	150	211	150	130	10 U
277- 6	SED13-CF	33%		10 U	10 U	10 U	29	18	20	22	110
277- 7	SED08-CM	38%		10 U	10 U	10 U	10 U	15	11	11	300
277- 8	SED14-BR	34%		10 U	10 U	10 U	11	18	13	15	27
277- 9	SED11-CB	32%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	16
277- 10	SED04-CM	54%		10 U	10 U	15	47	67	50	80	10 U
277- 11	SED02-CM	49%		10 U	10 U	15	41	58	44	63	88
277- 12	SED03-CM	57%		10 U	10 U	10 U	22	40	26	39	53
277- 13	SEDWR09-CM	19%		27	86	230	290	250	250	410	350
PLANTS											
277- 14	08A-SAL	89%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 15	14C-SAL	89%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 16	08C-SAL	86%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 17	04C-SAL	90%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 18	03C-SAL	84%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 19	11C-SCI	90%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 20	11A-SCI	87%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 21	11B-SCI	86%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 22	03B-SPA	87%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 23	04A-SPA	87%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 24	14A-SCI	74%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 25	14D-SAL	81%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 26	02C-SAL	84%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 27	05D-SAL	87%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 28	03D-SAL	85%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 29	08D-SAL	87%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 30	08B-SAL	86%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 31	11D-SCI	88%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 32	03A-SPA	86%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 33	05A-SPA	84%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U

277PAH

PAHS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

Battelle Code	Sponsor Code	% Moist.	Acenaph- thene	Acenaph- thylene	Anthra- cene	Benzo[a] Anthra- cene	Benzo[b] Fluor- anthene	Benzo[k] Fluor- anthene	Benzo[a] pyrene	Benzo- (g,h,i)- perylene	Chrysene
TISSUE											
277- 34	R08-SN01	66%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 34 DUP	R08-SN01	66%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 35	R08-SN02	35%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 36	R13-CBR1	92%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 37	R01-MOR1	85%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
277- 38	R01-MOR2	88%		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
BLANK	BLANK	N/A		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U
BLANK	BLANK	N/A		10 U	10 U	10 U	10 U	10 U	10 U	10 U	10 U

U indicates not detected at detection limit shown

277PAH

PAHS IN SEDIMENTS, PLANTS & TISSUE

Sponsor: SIMMER (McGUFFIE)

(CONCENTRATIONS IN UG/KG WET WEIGHT)

Battelle Code	Sponsor Code	% Molst.	Dibenzo- (a,h)- anthracene	Fluor- anthene	Fluorene	Indeno- 1,2,3- Pyrene	2-Methyl- Naphthalene	Naph- thalene	Phenan- threne	Pyrene
<u>TISSUE</u>										
277- 34	R08-SN01	66%	10 U	10 U	10 U	10 U	30 U	60 U	10 U	10 U
277- 34 DUP	R08-SN01	66%	10 U	10 U	10 U	10 U	30 U	60 U	10 U	10
277- 35	R08-SN02	35%	10 U	11	10 U	10 U	30 U	60 U	10 U	10 U
277- 36	R13-CBR1	92%	10 U	10 U	10 U	10 U	30 U	220	10 U	10 U
277- 37	R01-MOR1	85%	10 U	10 U	10 U	10 U	45	120	37	26
277- 38	R01-MOR2	88%	10 U	10 U	10 U	10 U	30 U	61	14	10 U
BLANK	BLANK	N/A	10 U	10 U	10 U	10 U	20 U	50 U	10 U	10 U
BLANK	BLANK	N/A	10 U	10 U	10 U	10 U	30 U	60 U	10 U	10 U

U indicates not detected at detection limit shown

REPORT DOCUMENTATION PAGE

*Form Approved
OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) November 2000		2. REPORT TYPE Final Report		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Field Survey of Contaminant Concentrations in Existing Wetlands in the San Francisco Bay Area				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) C. R. Lee, D. L. Brandon, J. W. Simmers, H. E. Tatem, R. A. Price, and S. P. Miner				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Engineer Research and Development Center Environmental Laboratory 3909 Halls Ferry Road Vicksburg, MS 39180-6199				8. PERFORMING ORGANIZATION REPORT NUMBER ERDC/EL SR-00-15	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Engineer District, San Francisco 333 Market St. San Francisco, CA 94105-2197				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The importance of wetlands to the productivity of estuaries has been realized in the San Francisco Bay Area. A heightened public interest has emerged to restore wetland acreage that has dwindled away over the past 50 years. Dredged material was thought to be of potential value in wetland creation or restoration. This report presents the results of a field survey of existing wetland sites in the San Francisco Bay Area. Dominant plants, of existing wetland sites in the San Francisco Bay Area. Dominant plants, animals (where present) and wetland soil from selected marine and estuarine wetlands were sampled and analyzed for contaminants. These data will be used to establish a wetland reference database. Sediment biological and chemical test results concentrations will be compared to the reference database to evaluate its potential use in wetland creation.					
15. SUBJECT TERMS Animal tissue concentrations Heavy models PCBs Wetlands Field survey PAHs Plant tissue concentrations					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT UNCLASSIFIED	b. ABSTRACT	c. THIS PAGE UNCLASSIFIED		118	19b. TELEPHONE NUMBER (include area code)