CRREL

REPORT 76-24

Vanadium and other elements in Greenland ice cores

Cover: The 1973 GISP field camp at Milcent, Greenland. (Photograph by C.C. Langway, Jr.)

CRREL Report 76-24

Vanadium and other elements in Greenland ice cores

M.M. Herron, C.C. Langway, Jr., H.V. Weiss,
J.P. Hurley, R.J. Kerr and J.H. Cragin

July 1976

Prepared for

NATIONAL SCIENCE FOUNDATION

By

CORPS OF ENGINEERS, U.S. ARMY

COLD REGIONS RESEARCH AND ENGINEERING LABORATORY

HANOVER, NEW HAMPSHIRE

Approved for public release; distribution unlimited.

*For conversion of SI metric units to U.S./British
customary units of measurement consult ASTM
Standard E380, Metric Practice Guide, published
by the American Society for Testing and Materials,
1916 Race St., Philadelphia, Pa. 19103.*

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER CRREL Report 76-24	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) VANADIUM AND OTHER ELEMENTS IN GREENLAND ICE CORES		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) M.M. Herron, C.C. Langway, Jr., H.V. Weiss, J.P. Hurley, R.J. Kerr and J.H. Cragin		8. CONTRACT OR GRANT NUMBER(s) NSF Project OPP 750-6750
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Army Cold Regions Research and Engineering Laboratory Hanover, New Hampshire 03755		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Division of Polar Programs National Science Foundation, Washington, D.C.		12. REPORT DATE July 1976
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Annual accumulations Ice Chemical analysis Ice cores Dating Seasonal variations Greenland Vanadium		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Chemical analysis of surface snows and deeper ice core samples from Milcent, Greenland, indicates a marine origin for Na and Cl and a terrestrial origin for Al, Mn and V. Pre-1900 enrichment factors, based on average crustal composition, are high for Zn and Hg and appear to be related to their volatility. A comparison of pre-1900 and 1971-1973 concentrations of V and Hg shows no decided increase from industrial production; however, the abundance of Zn (relative to Al) increased three-fold during this time period. The chemical composition of ancient ice is extremely useful in interpreting modern aerosols.		

PREFACE

This report was prepared by M.M. Herron, graduate student in the Department of Chemistry, San Diego State University; by Dr. C.C. Langway, Jr., Chairman of the Department of Geological Sciences, State University of New York at Buffalo; by Dr. H.V. Weiss, Research Chemist, Dr. J.P. Hurley, Radiation Physicist, and R.J. Kerr, Physicist, of the Naval Undersea Center, San Diego, California, and by J.H. Cragin, Chemist, of the Snow and Ice Branch, Research Division, U.S. Army Cold Regions Research and Engineering Laboratory. The research was funded by the National Science Foundation's Division of Polar Programs Project OPP 750-6750.

Technical reviewers of the manuscript were Dr. I.K. Iskandar and D. Leggett of CRREL.

The authors gratefully thank E.D. Goldberg and M. Koide for laboratory use.

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

VANADIUM AND OTHER ELEMENTS IN GREENLAND ICE CORES

M.M. Herron, C.C. Langway, Jr., H.V. Weiss,
J.P. Hurley, R.J. Kerr and J.H. Cragin

INTRODUCTION

The atmospheric concentrations of trace elements, particularly those which may include an anthropogenic fraction, have been the subject of much recent study (Zoller et al. 1973, Chester and Stoner 1974, Zoller et al. 1974, Duce et al. 1975). The crustal enrichment factor EF , used to identify possible pollutants, is defined as

$$EF_X = \frac{(X/Al)_{\text{SAMPLE}}}{(X/Al)_{\text{CRUST}}}$$

where $(X/Al)_{\text{SAMPLE}}$ and $(X/Al)_{\text{CRUST}}$ are the concentration ratios of the element X to aluminum in the sample and in average crustal material, respectively. An element whose aerosol originates primarily from crustal weathering processes should have an EF value near one. A significantly greater EF indicates input from local or additional sources, selective volatilization at the source (Zoller et al. 1974), selective transport or selective deposition from the atmosphere.

One method of distinguishing naturally high enrichment factors from man's contribution to the atmospheric burden is the analysis of chemical impurities within ice cores from the Greenland ice sheet. Pollutant lead and sulfur have been identified in modern Greenland snows by comparison with concentrations in pre-1900 ice strata (Murozumi et al. 1969, Koide and Goldberg 1971, Weiss et al. 1974, Cragin et al. 1975). Mercury was identified as a possible pollutant in northern Greenland ice (Weiss et al. 1971a), though work in southern Greenland has disputed this (Weiss et al. 1975). Extreme variations in

mercury concentrations in four northern Greenland samples were interpreted as suggesting that the natural mercury burden was highly variable (Carr and Wilkniiss 1972).

This paper presents the results of a study designed to determine the elemental concentrations of Na, Cl, Al, Mn, V, Zn, and Hg in Greenland ice and snow over the past several hundred years, and to help determine the significance of modern aerosol analyses.

SAMPLES

During the 1973 field season of the Greenland Ice Sheet Program a field camp was established at Milcent (70°18'N, 44°35'W). A continuous 400-m x 12-cm-diam ice core was taken at the main camp and surface snow samples were collected at four pits 2 to 3 km from the camp (pit 1 was always upwind). The individual who collected the samples wore a cleanroom coat, surgical face mask, hair covering, and powder-free polyethylene gloves throughout the collection, and about 10 cm of snow was trimmed from the pit walls with a precleaned stainless steel shovel before the samples were taken. One-year channel samples, as determined by stratigraphic features, were obtained with a polyethylene scoop, placed in 10-liter polyethylene containers, bagged, and shipped while frozen to the U.S. All bottles and field implements were precleaned by leaching in 1% HNO₃, rinsing five times with distilled, deionized water (DW), and rinsing twice with double-distilled, deionized water (DDW).

In the laboratory, the surface samples were melted in a microwave oven in their original 10-liter containers and immediately transferred to hot nitric acid treated (*HNAT*) 1-liter linear polyethylene bottles. These bottles had been filled with reagent grade HNO_3 , heated to 80°–90°C for periods greater than one hour, and rinsed five times with *DW*, twice with *DDW* and twice with sample meltwater. This treatment was proven effective in preventing adsorption of ^{48}V and ^{203}Hg for at least 32 days in separate 400-ml aliquots of the pit 4 sample. The spike concentrations were 6.8 ng ^{48}V /kg (5400 counts per minute, cpm) and < 1 ng ^{203}Hg /kg (2400 cpm).

The ice core samples were cleaned by the "dry" cleaning procedure (Langway et al. 1974), which involves extensive rinsing of the core with *DDW* in a Class 100 clean area. Three deep samples, not used for mercury concentration analysis, were stored in conventional 1-liter polyethylene bottles prepared by leaching with 7% Ultrex HNO_3 for several days at room temperature, followed by the usual rinsing procedure. A concentration of 10 ng ^{48}V /kg was added to a 250-ml aliquot of the pit 4 sample in one of the 1-liter bottles and quantitatively recovered after 21 days storage.

Zinc concentrations were measured in the unconcentrated samples using flameless atomic absorption with stopped nitrogen flow during atomization. Eppendorf pipette tips used for sample injection were found to give erroneously high concentrations until they were given the hot nitric acid treatment. Mercury concentrations were measured in 1-liter aliquots of the meltwater by neutron activation following the procedure of Williams et al. (1974). Before radiochemical Hg yields could be determined, the surface samples were destroyed in a laboratory fire and a yield of 0.9 was assumed. Aliquots containing 5 ml of the same meltwater were analyzed for Na and Cl by instrumental neutron activation analysis (*INAA*).

The remainder of the samples, 600 ml to 2 liters in volume, were evaporated to 20 ml in a precleaned quartz flask in a clean air station. They were transferred to 50-ml Teflon beakers in a glass enclosure under filtered N_2 , evaporated to 5 ml, and transferred to *HNAT* polyethylene irradiation vials (Weiss et al. 1971b). A spike of 0.2 ng ^{48}V added prior to the pre-concentration was quantitatively recovered. The samples were then analyzed for V, Al, and Mn by *INAA*. All counting was done with a 75 cm² Princeton Gamma-Tech Ge(Li) detector coupled to a Nuclear Data

Model 4420 computerized 4096 channel analyzer, and peaks were integrated by computer.

RESULTS AND DISCUSSION

The full data for Milcent snow and deeper ice are presented in Table I. Each ice core sample represents exactly two year's accumulation as determined by oxygen isotope ratios.*

The average chlorine to sodium ratio of the samples in Table I is 1.8, which matches the Cl/Na ratio of bulk seawater and suggests a marine origin for these two elements. Based on average seawater composition, the contribution of sea spray to the other elements is minimal.

The mercury concentrations are the highest yet reported for Greenland snow and ice, and there is no indication of significant Hg input in modern times. One interpretation is that the hot nitric acid treatment of the linear polyethylene bottles prevented adsorption on the container walls and resulted in more representative concentrations. Sample contamination is not considered to be very likely since samples from Point Barrow, Alaska, which were processed and analyzed identically to the Milcent samples, showed consistently low concentrations. The storage containers for the study of Weiss et al. (1971a) were prepared using a cold nitric acid leaching (Murozumi et al. 1969), and the samples were stored unacidified for three years in liquid state prior to analysis for Hg. In the study of Weiss et al. (1974) a cold acid leach was used and samples were analyzed 12 hours after melting (Weiss and Bertine 1973). Bottle preparation was not discussed by Carr and Wilkniss (1972).

Mean crustal enrichment factors using Al as the reference element are presented in Table II. The enrichment factors are calculated on the basis of the mean crustal abundance of Taylor (1964). Also given in Table II are enrichment factors for aerosol collections from Chester and Stoner (1974) for Atlantic Ocean "Westerlies" samples, from Duce et al. (1975) for Atlantic Ocean samples, and from Zoller et al. (1974) for South Pole aerosol samples.

* Personal communication, W. Dansgaard, University of Copenhagen.

Table I. Elemental concentrations in Milcent snow and ice samples (all concentrations in $\mu\text{g/kg}$).

Sample	Depth (m)	Age	Na	Cl	Al	V	Mn	Zn	Hg
Pit 1	0-1.36	1972-3	13	19	13	0.022	0.215	0.215	0.429*
Pit 1	1.36-2.27	1971-2	15	25	7	0.018	0.097	0.248	0.370*
Pit 1	1.52-2.75	1971-2	9	26	7	0.008	0.104	0.270	0.418*
Pit 2	0-1.51	1972-3	11	24	6	0.021	0.125	0.211	0.290*
Pit 3	0-1.37	1972-3	14	26	9	0.016	0.122		0.578*
Pit 4	0-1.38	1972-3	12	14	18	0.010	0.171	0.174	0.881*
1588	109.94-111.10	1803-5	9	22	3	BD†	0.075	0.038	
2757	169.43-170.42	1687-9	19	25	3	BD†	0.056	0.025	
4467	250.09-251.00	1600**	14	23	7	0.019	0.051	0.070	
1002	77.92-78.41	1864							0.716
1485	104.49-104.95	1815							0.318
2471	155.25-155.69	1717							0.823
2896	176.23-176.72	1675							0.263
3314	196.50-196.86	1631							0.445

* Assumes radiochemical yield of 0.9.

† Below detection limit of 0.007 $\mu\text{g/kg}$.

** Estimated age.

Table II. Enrichment factors at Milcent, based on average crustal abundances of Taylor (1964).

Sample	Enrichment factors				
	Al	V	Mn	Zn	Hg
Avg 1600-1805	1.0	$< 1.8 \pm 0.1$	1.5 ± 0.8	12 ± 3	$62,000 \pm 30,000^*$
Avg 1971-1973	1.0	1.1 ± 0.6	1.3 ± 0.4	32 ± 15	$53,000 \pm 11,000$
South Pole†	1.0	1.4	1.4	69	
Atlantic Ocean 30°N**	1.0	17	2.6	110	
Atlantic Westerlies ††	1.0	2.5	1.4	11	

* Assumes average Al concentration of 8.5 $\mu\text{g/kg}$ in the Milcent core.

† From Zoller et al. (1974).

** From Duce et al. (1975).

†† From Chester and Stoner (1974).

The crustal enrichment factors for V and Mn in Milcent ice are near unity and do not show any increase in modern deposits. This suggests a common origin due to crustal weathering. Duce et al. (1975) reported a vanadium *EF* of 17 for aerosol samples collected at 30°N latitude over the Atlantic Ocean, attributable to the burning of V-enriched fossil fuels. Zoller et al. (1974), on the other hand, found a vanadium *EF* of 1.4 for aerosol samples at the South Pole. The fact

that fossil fuel-derived V does not appear to reach the polar regions may be related to the relatively low volatility of V compounds.

The pre-1900 zinc *EF* in Milcent ice is 12 ± 3 , increasing to 32 ± 15 for 1971-1973 snows. This indicates not only that zinc is selectively volatilized, transported to and/or precipitated in Greenland, but also that the atmospheric load of zinc has increased in

modern times. The fossil fuel mobilization of Zn and V has been estimated at 7 and 12×10^9 g/year, respectively (Bertine and Goldberg 1971). Yet the Milcent evidence suggests that industrial Zn, and not V, reaches the Greenland ice sheet.

CONCLUSION

At present, it appears that elemental enrichment factors in pre-1900 Greenland ice may be explained adequately in terms of crustal abundances and relative volatilities. Thus Hg and Zn have high EF 's while those of V and Mn are near unity. Apparently the presence or absence of a pollution fraction in modern snows is also affected by relative volatilities. Efforts to expand the list of enrichment factors of ancient and modern precipitation are presently underway.

LITERATURE CITED

- Bertine, K.K. and E.D. Goldberg (1971) Fossil fuel combustion and the major sedimentary cycle. *Science*, vol. 173, p. 233-5.
- Carr, R.A. and P.E. Wilkniss (1972) Mercury in the Greenland ice sheet: Further data. *Science*, vol. 181, p. 843-4.
- Chester, R. and J.H. Stoner (1974) The distribution of Mn, Fe, Cu, Ni, Co, Ga, Cr, V, Ba, Sr, Sn, Zn and Pb in some soil-sized particulates from the lower troposphere over the world ocean. *Marine Chemistry*, vol. 2, p. 1273-85.
- Cragin, J.H., M.M. Herron and C.C. Langway, Jr. (1975) The chemistry of 700 years of precipitation at Dye 3, Greenland. CRREL Research Report 341. AD A014970.
- Duce, R.A., G.L. Hoffman and W.H. Zoller (1975) Atmospheric trace metals at remote northern and southern sites: pollution or natural? *Science*, vol. 187, p. 59-61.
- Koide, M. and E.D. Goldberg (1971) Atmospheric sulfur and fossil fuel combustion. *Journal of Geophysical Research*, vol. 76, p. 7689-96.
- Langway, C.C., Jr., M. Herron and J.H. Cragin (1974) Chemical profile of the Ross Ice Shelf at Little America V, Antarctica. *Journal of Glaciology*, vol. 13, p. 431-5.
- Murozumi, M., T.J. Chow and C. Patterson (1969) Chemical concentrations of pollutant lead aerosols, terrestrial dusts and sea salts in Greenland and Antarctica snow strata. *Geochimica et Cosmochimica Acta*, vol. 33, p. 1247-94.
- Taylor, S.R. (1964) Abundance of chemical elements in the continental crust: A new table. *Geochimica et Cosmochimica Acta*, vol. 28, p. 1273-85.
- Weiss, H.V., M. Koide and E.D. Goldberg (1971a) Mercury in a Greenland Ice Sheet: Evidence of recent input by man. *Science*, vol. 174, p. 692-4.
- Weiss, H.V., M. Koide and E.D. Goldberg (1971b) Selenium and sulfur in a Greenland Ice Sheet. *Science*, vol. 172, p. 261-3.
- Weiss, H.V. and K.K. Bertine (1973) Simultaneous determination of manganese, copper, arsenic, cadmium, antimony, and mercury in glacial ice by radioactivation. *Analytica Chimica Acta*, vol. 65, p. 253-9.
- Weiss, H.W., K.K. Bertine, M. Koide and E.D. Goldberg (1975) The chemical composition of a Greenland glacier. *Geochimica et Cosmochimica Acta*, vol. 39, p. 1-10.
- Williams, P.M., K.J. Robertson, K. Chew and H.V. Weiss (1974) Mercury in the south polar seas and in the northeast Pacific Ocean. *Marine Chemistry*, vol. 2, p. 287-99.
- Zoller, W.H., G.E. Gordon, E.S. Gladney and A.G. Jones (1973) The sources and distribution of vanadium in the atmosphere. *Advances in Chemistry Series 123*, American Chemical Society, Washington, D.C., p. 31-47.
- Zoller, W.H., E.S. Gladney and R.A. Duce (1974) Atmospheric concentrations and sources of trace metals at the South Pole. *Science*, vol. 183, p. 198-200.